

Status report of ttH analysis at 500 GeV

11th Apr. 2015

The 41st general meeting of the ILC physics working group




Yuji Sudo

Kyushu University

Introduction

- We are working on ttH study
 - $M_H=125$ GeV.
 - Polarization : $(P_{e^-}, P_{e^+})=(-0.8,+0.3)$
 - Integrated luminosity 500 fb^{-1}
 - ILD full simulation
- ttbar cross section is increased around ttbar threshold by ttbar bound-state effect
- ttH cross section is enhanced
- ttZ cross section is also increased

We can directly measure the top-Yukawa coupling via tth channel at $\sqrt{s} = 500$ GeV.



Updates

- Change Lepton ID method: cut base → BDT
- Increase MC statistics
- add $ttH \rightarrow 2l2\nu + 4b$ jets channel
- re-optimize event selection cut

Lepton ID efficiency

- Lepton ID efficiency with TMVA BDT (cut base) lepton selection

$ttH \rightarrow 2l2\nu + 4 \text{ jets}$

| (%) | elec | muon | taue | taum | tauh1 | tauh3 | bjet | ljet |
|------------------------|-----------------|------------------|------------------|------------------|------------------|-----------------|------------------|------|
| Identified as a lepton | 92.4 (90.91) | 96.53 (95.35) | 73.51 (63.09) | 74.63 (65.78) | 56.56 (47.08) | 49.3 (38.82) | 0.016 (0.017) | N/A |

ttZ

| (%) | elec | muon | taue | taum | tauh1 | tauh3 | bjet | ljet |
|------------------------|----------------|------------------|------------------|------------------|------------------|------------------|----------------|---------------|
| Identified as a lepton | 92.5 (89.8) | 96.61 (95.71) | 72.97 (63.38) | 73.41 (65.56) | 55.64 (45.81) | 48.27 (38.12) | 0.11 (0.14) | 0.6 (0.66) |

- Lepton ID efficiency and miss ID efficiency are improved by BDT method
- ID efficiency is almost same to ttZ and $ttH \rightarrow 2l2\nu + 4 \text{ jets}$.

Expected # of events @ 500fb⁻¹

- $\sqrt{s} = 500$ GeV, $M_h = 125$ GeV, $(P_{e^-}, P_{e^+}) = (-0.8, +0.3)$
- Production cross section
- Branching ratio

| Process | σ (fb) |
|------------------------------|---------------|
| $e^-e^+ \rightarrow tth$ | 0.485 |
| $e^-e^+ \rightarrow ttZ$ | 1.974 |
| $e^-e^+ \rightarrow ttg(bb)$ | 1.058 |
| $e^-e^+ \rightarrow tbW$ | 979.8 |

| Decay mode | Branching ratio |
|-------------------------|-----------------|
| $h \rightarrow bb$ | 0.577 |
| $tt \rightarrow bqqbqq$ | 0.457 |
| $tt \rightarrow blvbqq$ | 0.438 |
| $tt \rightarrow blvblv$ | 0.105 |

- Expected # of signals and Backgrounds(@500fb⁻¹)

| | | | |
|-----------------------------|--------------|---------|--------|
| ttH(tt→6j, H→bb) | 63.9 | | |
| ttH(tt→lv4j, H→bb) | 61.3 | ttZ | 987 |
| ttH(tt→lvlv2j, H→bb) | 14.6 | ttg(bb) | 529 |
| ttH(tt→all, H(nobb)) | 102.6 | tbW | 489902 |

Event selection lv+6 jets channel

- Isolated Lepton ID

- ✓ Exact one Isolated lepton

$$Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos\theta)}{E_{\text{cm}}^2}$$

- jet clustering : Durham algorithm

- forced 6 jets clustering is applied to ttH→6jets channel

- ✓ “ $Y_{65} > 0.0016$ ” + “ $Y_{65} \leq 0.0016 \ \&\& \ Y_{54} > 0.006$ ”

- ✓ b jet candidate ≥ 4 (b likeness $\geq 0.85, 0.8, 0.6, 0.2$)

- reject event with very forward jets

- ✓ $|\cos\theta_{\text{jet}}| \leq 0.99$

- Missing Momentum > 20 GeV

- Jet paring, $\chi^2 < 30.5$

- Leading 2 jets energy sum < 197 GeV

- smallest 2 jets energy sum > 66 GeV

- $M_{jjj}, M_{jlv} > 140$ GeV (mass of top candidate)



- $95 < M_{jj} < 160$ GeV (2 jets mass of higgs candidate)

Event selection lv+6 jets channel

- Isolated Lepton ID
 - ✓ Exact one Isolated lepton
- jet clustering : Durham algorithm

$$Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos \theta)}{E_{cm}^2}$$

- forced 6 jets clustering is applied to ttH → 6jets channel
 - ✓ “ $Y_{65} > 0.0016$ ” + “ $Y_{65} \leq 0.0016 \ \&\& \ Y_{54} > 0.006$ ”



Event selection lv+6 jets channel

- Isolated Lepton ID

✓ Exact one Isolated lepton

- jet clustering : Durham algorithm

$$Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos\theta)}{E_{cm}^2}$$

- forced 6 jets clustering is applied to ttH → 6jets channel

✓ “ $Y_{65} > 0.0016$ ” + “ $Y_{65} \leq 0.0016 \ \&\& \ Y_{54} > 0.006$ ”

✓ b jet candidate ≥ 4 (b likeness $\geq 0.85, 0.8, 0.6, 0.2$)

- reject event with very forward jet

✓ $|\cos\theta_{jet}| \leq 0.99$

• Missing Momentum > 20 GeV


• Jet paring, $\chi^2 < 30.5$

• Leading 2 jets energy sum < 197 GeV

• smallest 2 jets energy sum > 66 GeV



• $M_{jjj}, M_{jlv} > 140$ GeV (3 jets mass)

• $95 < M_{jj} < 160$ GeV (2 jets mass)



Event selection lv+6 jets channel

- Isolated Lepton ID



✓ $|\cos\theta_{jet}| \leq 0.99$

- Missing Momentum > 20 GeV
- Jet paring, $\chi^2 < 30.5$
- Leading 2 jets energy sum < 197 GeV
- smallest 2 jets energy sum > 66 GeV
- $M_{jjj}, M_{jlv} > 140$ GeV (3 jets mass of top candidate)
- $95 < M_{jj} < 160$ GeV (2 jets mass of higgs candidate)

Event selection lv+6 jets channel

- Isolated Lepton ID

✓ Exact one Isolated lepton

- jet clustering : Durham algorithm

$$Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos\theta)}{E_{cm}^2}$$

- forced 6 jets clustering is applied to ttH → 6jets channel

✓ “ $Y_{65} > 0.0016$ ” + “ $Y_{65} \leq 0.0016$ ”

✓ b jet candidate ≥ 4 (b likeness > 0.5)

- reject event with very forward jet

✓ $|\cos\theta_{jet}| \leq 0.99$

- Missing Momentum > 20 GeV


- Jet paring, $\chi^2 < 30.5$

- Leading 2 jets energy sum < 197 GeV

- smallest 2 jets energy sum > 66 GeV

- $M_{jjj}, M_{jlv} > 140$ GeV (3 jets mass of top candidate)


- $95 < M_{jj} < 160$ GeV (2 jets mass of higgs candidate)



Result of Event selection

higgs candidate M_{jj} (lv6jet)

- $\sqrt{s} = 500 \text{ GeV}$, $L=500 \text{ fb}^{-1}$, $(P_{e^-}, P_{e^+})=(-0.8, +0.3)$




| Process | # of evt |
|--|--------------|
| $t\bar{t}H$ (not Signal) | 0.25 |
| $t\bar{t}Z$ | 5.19 |
| $t\bar{t}bb$ | 2.04 |
| tbW | 8.39 |
| bkgd total | 15.88 |
| $t\bar{t}H \rightarrow lv6jet$ | 10.26 |

Result of Event selection


higgs candidate $M_{jj}(8jet, 2l2v4bjet)$

$ttH \rightarrow 8jet$



| Process | # of evt |
|--|--------------|
| ttH (not Signal) | 0.42 |
| ttZ | 7.17 |
| ttbb | 2.59 |
| tbW | 19.24 |
| bkgd total | 29.43 |
| ttH \rightarrow 8jet | 14.37 |

$ttH \rightarrow 2l2v+4jet$



| Process | # of evt |
|--|-------------|
| ttH (not Signal) | 0.06 |
| ttZ | 1.41 |
| ttbb | 0.54 |
| tbW | 1.88 |
| bkgd total | 3.91 |
| ttH \rightarrow 2l2v4jet | 2.62 |

tth ($h \rightarrow bb$) significance

$$S/\sqrt{S+B}$$

| (P_e, P_{e^+}) | (-0.8,+0.3) | | (0.8,-0.3) | |
|----------------------------|-------------|------|------------|------|
| Lumi. (fb^{-1}) | 500 | 1600 | 500 | 1600 |
| 8 jets | 2.17 | 3.89 | 1.40 | 2.53 |
| lv + 6 jets | 2.00 | 3.58 | 1.29 | 2.32 |
| 2l2v + 4 jets | 1.02 | 1.83 | 0.72 | 1.31 |

- (P_e, P_{e^+})=(-0.8,+0.3) polarization makes better result.
-- number of signal is important.

measurement precision at different \sqrt{s}

combined result of $ttH \rightarrow 8\text{jets}$, $lv+6\text{jets}$ and $2l2v+4b$ jets channels

$M_H = 125$ GeV, $(P_{e^-}, P_{e^+}) = (-0.8, +0.3)$

$\sigma_{ttH} = 0.485$ fb @ 500 GeV

$ttH (H \rightarrow bb) 500 \text{ fb}^{-1}$

$\sqrt{s} : S/\sqrt{S+B} : |\Delta y_t/y_t| \%$

500 : 3.13 : 16.6

520 : 5.26 : 9.88

550 : 7.72 : 6.73


$ttH (H \rightarrow bb) 1600 \text{ fb}^{-1}$

$\sqrt{s} : S/\sqrt{S+B} : |\Delta y_t/y_t| \%$

500 : 5.60 : 9.28

520 : 9.45 : 5.50

550 : 13.9 : 3.73



* Same MC samples are used for all analysis. MC sample are generated at 500 GeV.

* We just change production cross section of signal and backgrounds for each energy.

Summary

- lepton ID method: cut base \rightarrow BDT
- Counting analysis of ttH ($H \rightarrow bb$)
- Significance (stat. only) at 500 GeV
ttH ($H \rightarrow bb$) $S/\sqrt{S+B}$ ($|\Delta y_t/y_t|$)
500 fb⁻¹ : 3.13 (16.6 %)
1600fb⁻¹ : 5.60 (9.28 %)
- @ $\sqrt{s} = 520$ GeV, 500 fb⁻¹ : $S/\sqrt{S+B} = 5.26$
 $|\Delta y_t/y_t| = 9.88 \%$
- @ $\sqrt{s} = 480$ GeV, direct measurement of top Yukawa coupling is impossible.


Backup

ttH (H→bb) analysis

- interference term is negligible
- counting analysis with cut based event selection

In this analysis, higgs decays into two b jets

- **4 b jets** out of 4-8 jets (b tagging: LCFIPlus)
- Use Kt clustering only for removing low Pt background



Event selection

- **signal topology**
 - ✓ **number of jets**
 - ✓ **# of Isolated Lepton**
 - ✓ **# b jet candidates ≥ 4**
- **kinematics**
 - ✓ energy cut of leading jets
 - ✓ energy cut of low energy jets
 - ✓ Missing momentum > 20 GeV (4, 6jtes mode)
- **detector acceptance**
 - $|\text{Jet } \cos\theta| \leq 0.99$
- **jet paring**
 - ✓ **reconstructed mass**
 - ✓ M_{jj} of top candidates
 - ✓ M_{jj} of higgs candidate

Event Selection 8 jets

- Isolated Lepton ID with BDT

- ✓ require no Isolated lepton

$$Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos\theta)}{E_{\text{cm}}^2}$$

- Jet clustering : Durham algorithm

- forced 8 jet clustering for ttH→8jets channel

- ✓ “ $Y_{87} > 0.00038$ ” + “ $Y_{87} \leq 0.00038 \ \&\& \ Y_{76} > 0.004$ ”

- ✓ b candidate jets ≥ 4 (b likeness $\geq 0.85, 0.8, 0.6, 0.2$)

- reject events with very forward jets

- ✓ $|\text{Jet } \cos\theta| \leq 0.99$

- Jet paring, $\chi^2 < 13.3$

- Leading 2 jets energy sum < 188 GeV


- smallest 3 jets energy sum > 60 GeV

- $M_{jjj} > 140$ GeV (top candidate 3 jet mass)


- $95 < M_{jj} < 160$ (GeV) (range of higgs candidate M_{jj})

some variables used event selection ($ttH \rightarrow 8\text{jets}$)

$Y_{87} > 0.00038$






$Y_{87} \leq 0.00038 \ \&\& \ Y_{76} > 0.004$



Leading 2 jets
energy sum < 188 (GeV)

smallest 3 jets
energy sum > 60 GeV

$M_{jjj} > 140$ GeV



Event selection lv+6 jets channel

- Isolated Lepton ID

- ✓ Exact one Isolated lepton

$$Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos\theta)}{E_{\text{cm}}^2}$$

- jet clustering : Durham algorithm

- forced 6 jets clustering is applied to ttH→6jets channel

- ✓ “ $Y_{65} > 0.0016$ ” + “ $Y_{65} \leq 0.0016 \ \&\& \ Y_{54} > 0.006$ ”

- ✓ b jet candidate ≥ 4 (b likeness $\geq 0.85, 0.8, 0.6, 0.2$)

- reject event with very forward jets

- ✓ $|\cos\theta_{\text{jet}}| \leq 0.99$

- Missing Momentum > 20 GeV

- Jet paring, $\chi^2 < 30.5$

- Leading 2 jets energy sum < 197 GeV

- smallest 2 jets energy sum > 66 GeV


- $M_{\text{jjj}}, M_{\text{jl}v} > 140$ GeV (3 jets mass of top candidate)

- $95 < M_{\text{jj}} < 160$ GeV (2 jets mass of higgs candidate)


some variables used event selection

($ttH \rightarrow lv + 6\text{jets}$)


$Y_{65} > 0.0016$




$Y_{65} \leq 0.0016 \ \&\& \ Y_{54} > 0.006$




Missing Momentum > 20 GeV




Leading 2 jets
energy sum < 197 GeV



smallest 2 jets
energy sum > 66 GeV



$M_{jjj} > 140 \text{ GeV}$



Event Selection 2l2n+4 jets

- Isolated Lepton ID with BDT

- ✓ require exact two Isolated leptons

- Jet clustering : Durham algorithm $Y_{ij} = \frac{2\min\{E_i^2, E_j^2\}(1 - \cos\theta)}{E_{\text{cm}}^2}$

- forced 6 jet clustering for ttH→4jets channel

- ✓ “ $Y_{43} > 0.002$ ”

- ✓ 4 b jets (b likeness $\geq 0.85, 0.8, 0.6, 0.2$)

- reject events with very forward jets

- ✓ $|\text{Jet } \cos\theta| \leq 0.99$

- Missing momentum > 20 GeV

- Jet paring, $\chi^2 < 12.5$

- Leading jet energy < 112 (Gev)


- smallest jet energy > 38 (GeV)

- $100 < M_{jj} < 155$ (GeV) (range of higgs candidate M_{jj})


some variables used event selection

$(ttH \rightarrow 2l2\nu+4b)$


$Y_{43} > 0.002$




Missing momentum > 20 GeV



Leading jet energy < 112 (GeV)



smallest jet energy > 38 (GeV)



Jet pairing, χ^2 Cut (8 jets mode)

- $\sqrt{s} = 500\text{GeV}$ is near by threshold of the ttH production

- P_{higgs} should be small
- Dijet angle becomes large

→ Angle information between higgs candidate jets is effective to choose correct jet pair.

- try all combination and choose a pair with minimum χ^2 value

$$\chi^2 = \left(\frac{\Delta\text{angle}(j_1, j_2) - \Delta\text{angle}(\text{higgs } jj)}{\sigma_{\Delta\text{angle}(\text{higgs } jj)}} \right)^2 + \left(\frac{m_{j_3 j_4 j_5} - M_{\text{top}}}{\sigma_{M_{\text{top}}}} \right)^2 + \left(\frac{m_{j_4 j_5} - M_W}{\sigma_{M_W}} \right)^2 + \left(\frac{m_{j_6 j_7 j_8} - M_{\text{top}}}{\sigma_{M_{\text{top}}}} \right)^2 + \left(\frac{m_{j_7 j_8} - M_W}{\sigma_{M_W}} \right)^2$$

require b likeness ≥ 0.2 to j_1, j_2, j_3, j_6

- Reference values are made from reconstructed jets which are matched with MC information
 - $M_{\text{top}} = 171.5\text{GeV}$
 - $\sigma_{M_{\text{top}}} = 16.8\text{ GeV}$
 - $M_W = 80.5\text{GeV}$
 - $\sigma_{M_W} = 9.9\text{ GeV}$
 - $\text{angle}(jj) = 2.448$
 - $\sigma_{\text{angle}(jj)} = 0.277$

higgs and top pairing, χ^2 Cut (6 jets mode)

Angle information between higgs candidate jets is effective to choose correct jet pair.

$$\chi^2 = \left(\frac{\Delta angle(j_1, j_2) - \Delta angle(higgs jj)}{\sigma_{\Delta angle(higgs jj)}} \right)^2 + \left(\frac{m_{j_3 j_4 j_5} - M_{top}}{\sigma_{M_{top}}} \right)^2 + \left(\frac{m_{j_4 j_5} - M_W}{\sigma_{M_W}} \right)^2 + \left(\frac{m_{j_6 l \nu} - M_{top}}{\sigma_{M_{top}}} \right)^2$$

A W mass is reconstructed with Isolated lepton and Missing P

- try all combination and choose a pair with minimum χ^2 value

require b likeness ≥ 0.2 to j_1, j_2, j_3, j_6

- Reference values are made from reconstructed jets which are matched with MC information
 - $M_{top} = 171.5 \text{ GeV}$
 - $\sigma_{M_{top}} = 16.8 \text{ GeV}$
 - $M_W = 80.5 \text{ GeV}$
 - $\sigma_{M_W} = 9.9 \text{ GeV}$
 - $\text{angle}(jj) = 2.448$
 - $\sigma_{\text{angle}(jj)} = 0.277$

higgs and top pairing, χ^2 Cut (4 jets mode)

$$\chi^2 = \left(\frac{\Delta angle(j_1, j_2) - \Delta angle(higgs jj)}{\sigma_{\Delta angle(higgs jj)}} \right)^2$$

Angle information between higgs candidate jets is used to choose a jet pair.

try all combination and choose a pair with minimum χ^2 value

- Reference values are made from reconstructed jets which are matched with MC information
 - angle(jj) = 2.448
 - sigma angle(jj) = 0.277

event selection

MC stat.

tth, ttz, ttbb: 100k~200k events

tbW(DBD samples): 10k~100k events

Lepton ID

- muon selection



- electron selection



- tau (e)



- tau(muon)



- tau (1-prong)



- tau(3-prong)

Mjj shape estimation of tbW events in 4 b jets category

- tbW event: 6 fermion events including ttbar, single top etc..
- It is difficult to estimate Mjj shape of tbW events in 4b jets category, because of low statistics of tbW MC events
- At first, compare Mjj shapes of ttZ event in 2/4b jets category
- check event shape of variables which are used event selection. (ttZ, tbW)
- ✓ use Mjj shape of tbW event in 2b jets category
higgs candidate $M_{jj} \geq 80$ GeV

ttH \rightarrow 8jets, 500 fb⁻¹

| \sqrt{s} | $: S/\sqrt{S+B}$ | $: \Delta g_t/g_t \%$ |
|------------|------------------|-------------------------|
| 480 | : 0.59042 | : 88.0717 |
| 490 | : 1.34733 | : 38.5949 |
| 500 | : 2.1717 | : 23.9444 |
| 510 | : 2.96893 | : 17.5147 |
| 520 | : 3.70795 | : 14.0239 |
| 530 | : 4.38629 | : 11.8551 |
| 540 | : 4.993 | : 10.4146 |
| 550 | : 5.50563 | : 9.44488 |
| 560 | : 5.94584 | : 8.74562 |
| 570 | : 6.33231 | : 8.21186 |
| 580 | : 6.6747 | : 7.79062 |
| 590 | : 6.97431 | : 7.45594 |
| 600 | : 7.24148 | : 7.18085 |

$ttH \rightarrow l\nu + 6\text{jets}, 500 \text{ fb}^{-1}$

$\sqrt{s} \quad : S/\sqrt{S+B} : |\Delta g_t/g_t| \%$

480 : 0.57541 : 90.3704

490 : 1.27453 : 40.7994

500 : 2.00677 : 25.9123

510 : 2.69726 : 19.2788

520 : 3.32768 : 15.6265

530 : 3.90086 : 13.3304

540 : 4.41032 : 11.7905

550 : 4.83863 : 10.7469

560 : 5.2051 : 9.9902

570 : 5.52592 : 9.4102

580 : 5.80966 : 8.95062

590 : 6.05733 : 8.58464

600 : 6.27784 : 8.2831

$ttH \rightarrow 2l2n + 4b \text{ jets}, 500 \text{ fb}^{-1}$

$\sqrt{s} : S/\sqrt{S+B} : |\Delta g_t/g_t| \%$

480 : 0.29792 : 174.538

490 : 0.654923 : 79.3986

500 : 1.0257 : 50.697

510 : 1.37375 : 37.8526

520 : 1.69079 : 30.7549

530 : 1.97869 : 26.28

540 : 2.23437 : 23.2728

550 : 2.4491 : 21.2322

560 : 2.6327 : 19.7516

570 : 2.7933 : 18.6159

580 : 2.93529 : 17.7154

590 : 3.05914 : 16.9982

600 : 3.16936 : 16.4071

$ttH \rightarrow 8\text{jets}, 1600 \text{ fb}^{-1}$

| \sqrt{s} | $: S/\sqrt{S+B}$ | $: \Delta g_t/g_t \%$ |
|------------|------------------|-------------------------|
| 480 | : 1.04733 | : 49.6498 |
| 490 | : 2.40384 | : 21.632 |
| 500 | : 3.89502 | : 13.3504 |
| 510 | : 5.34728 | : 9.72456 |
| 520 | : 6.70016 | : 7.76101 |
| 530 | : 7.94612 | : 6.54407 |
| 540 | : 9.06307 | : 5.73757 |
| 550 | : 10.008 | : 5.19582 |
| 560 | : 10.8202 | : 4.80585 |
| 570 | : 11.5334 | : 4.50863 |
| 580 | : 12.1657 | : 4.27433 |
| 590 | : 12.7189 | : 4.08841 |
| 600 | : 13.2122 | : 3.93575 |

$ttH \rightarrow \ln + 6 \text{ jets}, 1600 \text{ fb}^{-1}$

$\sqrt{s} : S/\sqrt{S+B} : |\Delta g_t/g_t| \%$

480 : 1.02932 : 50.5185

490 : 2.27995 : 22.8076

500 : 3.58981 : 14.4854

510 : 4.825 : 10.7772

520 : 5.95273 : 8.73549

530 : 6.97807 : 7.45192

540 : 7.88943 : 6.5911

550 : 8.6556 : 6.00767

560 : 9.31117 : 5.58469

570 : 9.88506 : 5.26046

580 : 10.3926 : 5.00355

590 : 10.8357 : 4.79896

600 : 11.2301 : 4.63039

$ttH \rightarrow 2l2n + 4b \text{ jets}, 1600 \text{ fb}^{-1}$

$\sqrt{s} : S/\sqrt{S+B} : |\Delta g_t/g_t| \%$

480 : 0.53295 : 97.5699

490 : 1.17156 : 44.3852

500 : 1.83483 : 28.3405

510 : 2.45744 : 21.1603

520 : 3.02458 : 17.1925

530 : 3.53958 : 14.691

540 : 3.99696 : 13.0099

550 : 4.38109 : 11.8692

560 : 4.70951 : 11.0415

570 : 4.99681 : 10.4066

580 : 5.25081 : 9.90324

590 : 5.47236 : 9.50229

600 : 5.66952 : 9.17186