


whizard tutorial for physics at lepton colliders

Junping Tian (KEK)



tutorial set up

- simplest way: get a copy of the disk image of virtual machine (~6 GB); get it installed in your virtualbox; then you will already have all the needed libraries and also examples for this tutorial.

#if get that virtual machine started successfully, first thing you may would like to do is **change the keyboard layout**: System —>Preference—>Keyboard—>layout, set default your favourite (currently Japanese layout)

virtual machine: /home/ilc/tianjp/public/tutorial/whizard/
whizard_VM/whizard195_SL67.vdi

tutorial set up

- if you are on kekcc —> much simpler

```
cd /home/ilc/tianjp/public/tutorial/whizard/  
whizard_tutorial_MGS2015
```

everything is in “README”

tutorial set up

- alternatively: make available following libs

```
##### prerequisites to install whizard#####
```

```
## following libraries are necessary
```

```
# 1. GNU Make
```

```
# 2. PERL 5
```

```
# 3. Fortran 95
```

```
# 4. Fortran 77
```

```
# 5. C compiler
```

```
# 6. Objective Caml (e.g. ocaml-3.11.2)
```

```
## following libraries are only needed for more detailed analysis
```

```
# 1. Pythia (for fragmentation)
```

```
# 2. STDHEP (for output events)
```

```
# 3. Tauola (for proper decay of tau regarding polarisations)
```

```
# 4. Latex (for plotting Feynman Diagrams)
```

```
# 5. MetaPost (for plotting histograms)
```

```
# 6. PDFLIB (for parton distribution functions, from CERNLIB)
```

http://www-jlc.kek.jp/~tianjp/whizard_tutorial_MGS2015.tar.gz

get tutorial started

▶ `cd $HOME / whizard_tutorial_MGS2015 ; ls ;`

```
README          example02/      packages/  
example00/      example03/      processes/  
example01/      example04/      setup_for_whizard.sh
```

▶ `emacs README ;`

▶ `source setup_for_whizard.sh ;`

example01

$$e^+ e^- \rightarrow \mu^+ \mu^- H$$

- ▶ `tar zxvf packages/whizard-1.95.tgz -C example01 ;`
- ▶ `cd example01 ; ./configure ;`
- ▶ `cp ../processes/whizard.example01.prc conf/whizard.prc ;`
- ▶ `make prg install ;`

example01

$$e^+ e^- \rightarrow \mu^+ \mu^- H$$

- ▶ `cd results ;`
- ▶ `cp ../../processes/whizard.example01.in whizard.in ;`
- ▶ `cp ../../processes/whizard.example01.cut5 whizard.cut5`
- ▶ `./whizard ;`
 - `# find cross sections and summary in "whizard.out"`
 - `# 10K events should have been saved to "whizard.evt"`
 - `# visualize the Feynman diagrams in this process`
 - ▶ `make channels ; evince "whizard-channels.ps" ;`
 - `# visualize some kinematic distributions`
 - ▶ `make plots ; evince "whizard-plots.ps" ;`

example01

$$e^+ e^- \rightarrow \mu^+ \mu^- H$$


WHIZARD phase space channels

Process: e2e2h_o ($e^- e^+ \rightarrow \mu^- \mu^+ H$)

Color code: resonance, t-channel, radiation, infrared, collinear, external/off-shell


Grove 1

Multiplicity: 2
Resonances: 1
Log-enhanced: 0
t-channel: 0



Grove 2

Multiplicity: 3
Resonances: 0
Log-enhanced: 1
t-channel: 0



example01

$$e^+ e^- \rightarrow \mu^+ \mu^- H$$


WHIZARD data analysis

November 25, 2015

Process: e2e2h_o ($e^- e^+ \rightarrow \mu^- \mu^+ H$)

$$\sqrt{s} = 500.0 \text{ GeV} \quad \int \mathcal{L} = 5103. \text{ fb}^{-1}$$

#evt/bin



$$\sigma_{\text{tot}} = 1.9598 \pm 0.306 \times 10^{-02} \text{ fb} \quad [\pm 0.16 \%] \quad n_{\text{evt, tot}} = 10000$$

$$\sigma_{\text{cut}} = 1.9598 \pm 0.196 \times 10^{-01} \text{ fb} \quad [\pm 1.00 \%] \quad n_{\text{evt, cut}} = 10000 \quad [100.00 \%]$$

example01

$$e^+ e^- \rightarrow \mu^+ \mu^- H$$


WHIZARD data analysis

November 25, 2015

Process: e2e2h_o ($e^- e^+ \rightarrow \mu^- \mu^+ H$)

$$\sqrt{s} = 500.0 \text{ GeV} \quad \int \mathcal{L} = 5103. \text{ fb}^{-1}$$

#evt/bin



$$\sigma_{\text{tot}} = 1.9598 \pm 0.306 \times 10^{-02} \text{ fb} \quad [\pm 0.16 \%] \quad n_{\text{evt, tot}} = 10000$$

$$\sigma_{\text{cut}} = 1.9598 \pm 0.196 \times 10^{-01} \text{ fb} \quad [\pm 1.00 \%] \quad n_{\text{evt, cut}} = 10000 \quad [100.00 \%]$$

example01: ingredients to run a process by whizard

- define a process in “whizard.prc”
- configure model parameters in “whizard.in”
- histogram and cut in “whizard.cut5”

all located in the “example01 / results /”

define a process in “whizard.prc”

```
# The selected model (O'Mega)
model SM_CKM

# Processes for example01
# (Methods: chep=CompHEP, mad=MadGraph, omega=O'Mega)
# (Options: s=selected diagrams, number=QCD order [Madgraph])
# f=fudged width [O'Mega]
#
# Tag In Out Method Option
#=====
#####
#####
#
e2e2h_o e1,E1 e2,E2,h omega w:c,c
```

all the particle names are defined in “\${Model}.mdl”, e.g.

“example01 / conf / models / SM_CKM.mdl”;

particle: e1,e2,e3,n1,n2,n3,u,d,c,s,t,b

anti-particle: E1,E2,E3,N1,N2,N3,U,D,C,S,T,B

W+,W-,Z,h,a,g

WHIZARD configuration file

The selected model

model SM

Processes

Methods: chep=CompHEP, mad=MadGraph, omega=0'Mega, test=trivial)

Options: s selected diagrams (CompHEP/MadGraph)

r restricted intermediate state (0'Mega)

c apply exact color algebra (0'Mega)

n:XXX coupling order (MadGraph)

w:XXX width scheme (0'Mega)

p transfer polarization (test)

u unit matrix element (test)

#

Tag In Out Method Option

#=====

ee_c	e1,E1	e1,E1	chep	
ee_m	e1,E1	e1,E1	mad	
ee_o	e1,E1	e1,E1	omega	
ww_c	e1,E1	W+,W-	chep	
ww_m	e1,E1	W+,W-	mad	
ww_o	e1,E1	W+,W-	omega	
zh_c	e1,E1	Z,H	chep	
zh_m	e1,E1	Z,H	mad	
zh_o	e1,E1	Z,H	omega	
nnh_c	e1,E1	n1,N1,H	chep	
nnh_m	e1,E1	n1,N1,H	mad	
nnh_o	e1,E1	n1,N1,H	omega	
nnbb_m	e1,E1	n1,N1,b,B	mad	
nnbb_o	e1,E1	n1,N1,b,B	omega	

configure process parameters in "whizard.in"

```
&process_input
```

```
process_id = "e2e2h_o"
```

```
sqrts = 500
```

```
luminosity = 10
```

```
/
```

```
&integration_input
```

```
/
```

```
&simulation_input
```

```
n_events = 10000
```

```
/
```

```
&diagnostics_input
```

```
/
```

```
&parameter_input
```

```
Mmu = 0
```

```
MH = 125
```

```
wH = 0.0043
```

```
/
```

```
&beam_input
```

```
/
```

```
&beam_input
```

```
/
```

specify process, ecm, lumi, etc.

control integration accuracy, seed, etc.

set no. evts, evt. format, fragmentation, etc.

set time limit, etc.

set particle mass, coupling constant, etc.

set beam spectrum, polarisation, ISR, etc.

see complete config in example04

histogram and cut in “whizard.cut5”

e2e2h_o	e1,E1	e2,E2,h	omega
binary code	8 16	1 2 4	

```
# Analysis configuration file for the process ee -> mu mu h
process e2e2h_o
histogram M of 3 within 70 110 nbin 40
and
histogram E of 4 within 100 300 nbin 50
and
histogram PT of 3 within 0 300 nbin 50
and
histogram CTA of 3 within -1 1 nbin 50
and
histogram CTA of 1 within -1 1 nbin 50
and
histogram CTA of 2 within -1 1 nbin 50
```

cut PT of 4 within 0 150

cut M of 3 within 80 100 or 180 200 or 500 99999

example00

compare results using matrix elements
by **Omega, ComHEP, MadGraph**

- ▶ `tar zxvf packages / whizard-1.95.tgz -C example00 ;`
- ▶ `cd example00 ;`
- ▶ `cp ../processes / config.example00.site config.site ;`
- ▶ `./configure ;`
- ▶ `make test ; make test-QED ; make test-QCD`

there's a typo in README; order 1.3 and 1.4 should be swapped

example00

compare results using matrix elements
by **Omega**, **ComHEP**, **MadGraph**

$e^+e^- \rightarrow \nu\bar{\nu}H$

```
!=====  
! Summary (all processes):  
!-----  
! Process ID      Integral[fb]  Error[fb]   Err[%]      Frac[%]  
!-----  
nnh_c            8.4711467E+01  3.52E-01    0.42        33.35  
nnh_m            8.4435645E+01  3.40E-01    0.40        33.24  
nnh_o            8.4890431E+01  3.53E-01    0.42        33.42  
!-----  
sum              2.5403754E+02  6.04E-01    0.24        100.00  
!=====
```

$e^+e^- \rightarrow \mu^+\mu^-$

```
Running WHIZARD for process ee_e2: e1,E1 -> e2,E2  
ee_e2_o          3.4821868E+02  3.20E+00    0.92        33.33  
ee_e2_m          3.4821868E+02  3.20E+00    0.92        33.33  
ee_e2_c          3.4821868E+02  3.20E+00    0.92        33.33
```

$u\bar{u} \rightarrow t\bar{t}$

```
Running WHIZARD for process uu_u3: u,U -> t,T  
**uu_u3_o        1.7958449E+04  6.86E+01    0.38        33.33  
uu_u3_m          1.7958450E+04  6.86E+01    0.38        33.33  
uu_u3_c          1.7958450E+04  6.86E+01    0.38        33.33
```

example02

$$e^+e^- \rightarrow \nu_e\bar{\nu}_e H$$

$$e^+e^- \rightarrow e^+e^- H$$

- ▶ `tar zxvf packages/whizard-1.95.tgz -C example02 ;`
- ▶ `cd example02 ; ./configure ;`
- ▶ `cp ../processes/whizard.example02.prc conf/whizard.prc ;`
- ▶ `make prg install ;`
- ▶ `cd results ;`
- ▶ `cp ../../processes/whizard.example02.in whizard.in ;`
- ▶ `cp ../../processes/whizard.example02.cut5 whizard.cut5`
- ▶ `./whizard ;`
- ▶ `make channels ; evince "whizard-channels.ps" ;`
- ▶ `make plots ; evince "whizard-plots.ps" ;`

example02: select Feynman diagrams in “whizard.prc”

$$e^+ e^- \rightarrow \nu_e \bar{\nu}_e H$$

$$e^+ e^- \rightarrow e^+ e^- H$$

```
# Tag          In          Out          Method  Option
#=====
#####
#
n1n1h_o       e1,E1    n1,N1,h      omega    w:c,c
n1n1h_s_o     e1,E1    n1,N1,h      omega    w:c,c,r:3+4~Z
n1n1h_t_o     e1,E1    n1,N1,h      omega    w:c,c,r:1+3~W- && 2+4~W+

e1e1h_o       e1,E1    e1,E1,h      omega    w:c,c
e1e1h_s_o     e1,E1    e1,E1,h      omega    w:c,c,r:3+4~Z
e1e1h_t_o     e1,E1    e1,E1,h      omega    w:c,c,r:1+3~Z && 2+4~Z
```

#use option “r” to restrict the intermedia state

#note the particle number in “r” is different with binary code

example02: select Feynman diagrams in “whizard.prc”

$$e^+ e^- \rightarrow \nu_e \bar{\nu}_e H$$


WHIZARD phase space channels

Process: n1n1h_o ($e^- e^+ \rightarrow \nu_e \bar{\nu}_e H$)

Color code: resonance, t-channel, radiation, infrared, collinear, external/off-shell


Grove 1

Multiplicity: 2
Resonances: 1
Log-enhanced: 0
t-channel: 0



Grove 2

Multiplicity: 3
Resonances: 0
Log-enhanced: 2
t-channel: 2



#be careful when use restriction; do not violate gauge invariance, neither ignore interference


example02

$$e^+ e^- \rightarrow \nu_e \bar{\nu}_e H$$

Process: n1n1h_o ($e^- e^+ \rightarrow \nu_e \bar{\nu}_e H$)

$$\sqrt{s} = 500.0 \text{ GeV} \quad \int \mathcal{L} = 57.35 \text{ fb}^{-1}$$


#evt/bin



$$\sigma_{\text{tot}} = 77.810 \pm 0.323 \text{ fb} \quad [\pm 0.42 \text{ \%}] \quad n_{\text{evt, tot}} = 4521$$

$$\sigma_{\text{cut}} = 77.810 \pm 1.16 \text{ fb} \quad [\pm 1.49 \text{ \%}] \quad n_{\text{evt, cut}} = 4521 \quad [100.00 \text{ \%}]$$

homework (A): use whizard to reproduce following plot



example03: on-shell and off-shell Higgs decay

$$e^+e^- \rightarrow ZH$$

$$e^+e^- \rightarrow ZH \rightarrow Z(W^+W^-)$$



```
# Tag          In      Out          Method  Option
#=====
#####
#
zh_o           e1,E1    Z,h          omega    w:c,c
wwz_h_o       e1,E1    W+,W-,Z     omega    w:c,c,r:3+4~h
```

#same procedure to run as example01 / 02; see README

example03: on-shell and off-shell Higgs decay


$$e^+e^- \rightarrow ZH$$

$$e^+e^- \rightarrow ZH \rightarrow Z(W^+W^-)$$



#off-shell contribution / on-shell $\sim 3\%$

homework (B): $e^+e^- \rightarrow WWZ$ including both on-shell and off-shell Higgs contribution



example04: full ILC setup

more automatic, detailed instruction see example04/README

```
# The selected model (O'Mega)
model SM_CKM
```

```
# Tag          In          Out          Method  Option
#=====
```

```
#
alias f u:d:s:c:b:e1:e2:e3:n1:n2:n3
alias F U:D:S:C:B:E1:E2:E3:N1:N2:N3
```

```
# Tree ffh process
ffh_o      e1,E1    f,F,h      omega    w:c,c
ffhh_o     e1,E1    f,F,h,h   omega    w:c,c
```

#note features of using alias to include all possible processes

example04: full ILC setup

#e.g. enable beam spectrum

```
&process_input  
process_id = ""  
sqrts = 500  
luminosity = 1000  
polarized_beams = T  
structured_beams = T  
beam_recoil = T  
/
```

```
&beam_input  
particle_name = 'e1'  
polarization = 1.0 0.0  
USER_spectrum_on = T  
USER_spectrum_mode = 21  
ISR_on = T  
ISR_alpha = 0.0072993  
ISR_m_in = 0.000511  
EPA_on = F  
EPA_alpha = 0.0072993  
EPA_m_in = 0.000511  
EPA_mX = 4.  
EPA_Q_max = 4.  
/
```

example04: full ILC setup

#e.g. tune fragmentation parameters, specify Higgs decay branching ratios, set output event format (e.g. STDHEP)

```
&simulation_input
n_events = 10000
write_events_raw = F
keep_beam_remnants = T
fragment = T
fragmentation_method = 3
! OPAL tune
pythia_parameters = "PMAS(25,1)=125.0; PMAS(25,2)=0.0043;
MSTJ(41)=2; MSTU(22)=20; MSTJ(28)=2;
PARJ(21)=0.40000; PARJ(41)=0.11000; PARJ(42)=0.52000; PARJ(81)=0.25000;
PARJ(82)=1.90000; MSTJ(11)=3; PARJ(54)=-0.03100; PARJ(55)=-0.00200;
PARJ(1)=0.08500; PARJ(3)=0.45000; PARJ(4)=0.02500; PARJ(2)=0.31000;
PARJ(11)=0.60000; PARJ(12)=0.40000; PARJ(13)=0.72000; PARJ(14)=0.43000;
PARJ(15)=0.08000; PARJ(16)=0.08000; PARJ(17)=0.17000; MSTP(3)=1;
MWID(25)=2;
BRAT(212)=0.00044;BRAT(213)=0.0268;BRAT(214)=0.578;BRAT(219)=0.000221;
BRAT(220)=0.0637;BRAT(222)=0.0856;BRAT(223)=0.0023;BRAT(224)=0.00155;
BRAT(225)=0.0267;BRAT(226)=0.216"
! MDME(219,1)=0; To suppress h-> mu mu decay
write_events = T
write_events_format = 20
bytes_per_file = 500000000
/
```

#based on STDHEP events, use similar analysis method as what used for MadGraph based analysis, or SGV/ilcsoft which are commonly used by ILC community

homework (C): plot recoil mass using $Z \rightarrow \mu\mu$ by taking into account realistic beam spectrum

