





*Jet origin identification:  
AI enhanced reconstruction for  
Higgs factory*




Manqi Ruan

# Yields $\sim$ Xsec \* Lumi \* Time



- 4 Million Higgs (10 years)
- ~ 1 Giga W (1 year) + 4 Tera Z (2 years)
- Upgradable: Top factory (500 k ttbar)

# CEPC Physics study



Chinese Physics C Vol. 43, No. 4 (2019) 043002

## Precision Higgs physics at the CEPC\*

Fenfen An(安芬芬)<sup>4,23</sup> Yu Bai(白羽)<sup>6</sup> Chunhui Chen(陈春晖)<sup>23</sup> Xin Chen(陈新)<sup>5</sup> Zhenxing Chen(陈振兴)<sup>8</sup> Joao Guimaraes da Costa<sup>1</sup> Zhenwei Cui(崔振威)<sup>3</sup> Yaquan Fang(方亚泉)<sup>4,6,34,35</sup> Chengdong Fu(付成栋)<sup>1</sup> Jun Gao(高俊)<sup>23</sup> Yanyan Gao(高艳彦)<sup>22</sup> Yunning Gao(高原宁)<sup>2</sup> Shaofeng Ge(葛绍峰)<sup>13,29</sup> Jiayin Gu(顾嘉茵)<sup>13,29</sup> Fangyi Guo(郭方懿)<sup>4,4</sup> Jun Guo(郭军)<sup>19</sup> Tao Han(韩涛)<sup>31</sup> Shuang Han(韩爽)<sup>4</sup> Hongjian He(何红建)<sup>11,18</sup> Xianke He(何显柯)<sup>16</sup> Xiaogang He(何小刚)<sup>11,16,20</sup> Jifeng Hu(胡耀峰)<sup>16</sup> Shih-Chieh Hsu(徐士杰)<sup>12</sup> Shan Jin(金山)<sup>8</sup> Maoqiang Jing(荆茂强)<sup>3,5</sup> Susmita JyotiShamti<sup>33</sup> Ryuta Kinoshita<sup>4</sup> Chia-Ming Kuo(郭家铭)<sup>31</sup> Peizhuo Lai(赖培筑)<sup>3</sup> Boyang Li(李博扬)<sup>3</sup> Congqiao Li(李聪乔)<sup>3</sup> Gang Li(李刚)<sup>4,34,35</sup> Haifeng Li(李海峰)<sup>12</sup> Liang Li(李亮)<sup>19</sup> Shu Li(李数)<sup>11,19</sup> Tong Li(李通)<sup>12</sup> Qiang Li(李强)<sup>3</sup> Hao Liang(梁浩)<sup>4,6</sup> Zhijun Liang(梁志均)<sup>4</sup> Libo Liao(廖立波)<sup>4</sup> Bo Liu(刘波)<sup>4,23</sup> Jianbei Liu(刘建北)<sup>3</sup> Tao Liu(刘涛)<sup>14</sup> Zhen Liu(刘真)<sup>28,36,6</sup> Xinchou Lou(娄辛丑)<sup>4,43,14</sup> Lianliang Ma(马连良)<sup>12</sup> Bruce Mellado<sup>13,18</sup> Xin Mo(莫欣)<sup>4</sup> Mila Pandurovic<sup>16</sup> Jianming Qian(钱剑明)<sup>34,9</sup> Zhaonai Qian(钱卓妮)<sup>19</sup> Nikolaos Rempotis<sup>22</sup> Manqi Ruan(阮曼奇)<sup>40</sup> Alex Schry<sup>32</sup> Lianyou Shan(单连友)<sup>4</sup> Jingyuan Shi(史静远)<sup>4</sup> Xin Shi(史欣)<sup>4</sup> Shufang Su(苏淑芳)<sup>27</sup> Dayong Wang(王大勇)<sup>3</sup> Jun Wang(王隽)<sup>4</sup> Liantao Wang(王连涛)<sup>27,7</sup> Yifang Wang(王贻芳)<sup>4,6</sup> Yuqian Wei(魏或琦)<sup>4</sup> Yue Xu(许悦)<sup>3</sup> Haijun Yang(杨海军)<sup>30,31</sup> Ying Yang(杨迎)<sup>4</sup> Weiming Yao(姚伟明)<sup>2</sup> Dan Yu(于丹)<sup>4</sup> Kaiji Zhang(张凯冀)<sup>4,49</sup> Zhaoru Zhang(张照茹)<sup>4</sup> Mingrui Zhao(赵明锐)<sup>2</sup> Xiaohu Zhao(赵祥虎)<sup>4</sup> Ning Zhou(周宁)<sup>19</sup>

White papers +  
~300 Journal/AxXiv citables

Qingdao 266237, China  
<sup>1</sup>PRISMA Cluster of Excellence & Mainz Institute of Theoretical Physics, Johannes Gutenberg-Universität Mainz, Mainz 55128, Germany  
<sup>2</sup>Department of Physics, Hong Kong University of Science and Technology, Hong Kong  
<sup>3</sup>Kevin IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan  
<sup>4</sup>Vasco Institute of Nuclear Sciences, University of Belgrade, Belgrade 11000, Serbia  
<sup>5</sup>School of Physics and Institute for Collider Particle Physics, University of the Witwatersrand, Johannesburg 2050, South Africa

Received 9 November 2018, Revised 21 January 2019, Published online 4 March 2019

\* Supported by the National Key Program for S&T Research and Development (2016YFA040400); CAS Center for Excellence in Particle Physics; Yifang Wang's Science Studio of the Ten Thousand Talents Project; the CAS-SAFEA International Partnership Program for Creative Research Teams (H21518183); BHEP Innovation Grant (Y454517072); Key Research Program of Frontier Sciences, CAS (SQZDYDY-SW-SL3002); Chinese Academy of Science Special Grant for Large Scientific Project (113111KY5B20170005); the National Natural Science Foundation of China(11675203); the Hundred Talent Programs of Chinese Academy of Science (Y151540U1); the National 1000 Talents Program of China; Front Research Alliance, LLC (DE-AC02-07CH11359); the NSF(PHY1610074); by the Maryland Center for Fundamental Physics (MCFP); Tsinghua University Initiative Scientific Research Program; and the Beijing Municipal Science and Technology Commission project(Z181100004218003).

- 1) E-mail: fangyi@ihep.ac.cn
- 2) E-mail: jiang@ust.hk
- 3) E-mail: li.gang@mail.ihep.ac.cn
- 4) E-mail: zhanghs@imsl.edu
- 5) E-mail: qiong@imsl.edu
- 6) E-mail: manqi.ruan@ihep.ac.cn
- 7) E-mail: liantao@uchicago.edu
- 8) E-mail: zhangk1@ihep.ac.cn

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Article funded by SCOAP<sup>3</sup> and published under licence by Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

Scientific Significance quantified by CEPC physics studies, via full simulation/phenomenology studies:

- Higgs: Precisions exceed HL-LHC ~ 1 order of magnitude.
- EW: Precision improved from current limit by 1-2 orders.
- Flavor Physics, sensitive to NP of 10 TeV or even higher.
- Sensitive to varies of NP signal.
- ...





Table 2.1: Precision of the main parameters of interests and observables at the CEPC, from Ref. [1] and the references therein, where the results of Higgs are estimated with a data sample of 20 ab<sup>-1</sup>. The HL-LHC projections of 3000 fb<sup>-1</sup> data are used for comparison. [2]

Observable	Higgs		W, Z and top		
	HL-LHC projections	CEPC precision	Observable	Current precision	CEPC precision
$M_H$	20 MeV	3 MeV	$M_W$	9 MeV	0.5 MeV
$\Gamma_H$	20%	1.7%	$\Gamma_W$	49 MeV	2 MeV
$\sigma(ZH)$	4.2%	0.26%	$M_{top}$	760 MeV	$\mathcal{O}(10)$ MeV
$B(H \rightarrow bb)$	4.4%	0.14%	$M_Z$	2.1 MeV	0.1 MeV
$B(H \rightarrow cc)$	-	2.0%	$\Gamma_Z$	2.3 MeV	0.025 MeV
$B(H \rightarrow gg)$	-	0.81%	$R_b$	$3 \times 10^{-3}$	$2 \times 10^{-4}$
$B(H \rightarrow WW^*)$	2.8%	0.53%	$R_c$	$1.7 \times 10^{-2}$	$1 \times 10^{-3}$
$B(H \rightarrow ZZ^*)$	2.9%	4.2%	$R_\mu$	$2 \times 10^{-3}$	$1 \times 10^{-4}$
$B(H \rightarrow \tau^+\tau^-)$	2.9%	0.42%	$R_\tau$	$1.7 \times 10^{-2}$	$1 \times 10^{-4}$
$B(H \rightarrow \gamma\gamma)$	2.6%	3.0%	$A_\mu$	$1.5 \times 10^{-2}$	$3.5 \times 10^{-5}$
$B(H \rightarrow \mu^+\mu^-)$	8.2%	6.4%	$A_\tau$	$4.3 \times 10^{-3}$	$7 \times 10^{-5}$
$B(H \rightarrow Z\gamma)$	20%	8.5%	$A_b$	$2 \times 10^{-2}$	$2 \times 10^{-4}$
$B(\text{upper}(H \rightarrow \text{inv.}))$	2.5%	0.07%	$N_\nu$	$2.5 \times 10^{-3}$	$2 \times 10^{-4}$

# Performance requirements

- To reconstruct all kinds of Physics Object
  - Identification & Measurements
  - Objects:
    - Lepton, Photons, Kaon,
    - pi-0, Tau, Lambda, Kshort,
    - Heavy flavor hadrons,
    - **Jets**
    - Missing energy/momentum
    - Exotics...
- Massive Four in Standard Model:
  - Z & W: ~ 70% goes to a pair of jets
  - Higgs: ~90% final state with jets (ZH events)
  - Top:  $t \rightarrow W + b$



- Requirements:
  - **1-1 correspondence**  
Excellent pattern. Reco. & Object id
  - Larger acceptance, Excellent intrinsic resolutions, Extremely stable...
- Be addressed by detector design, technology, and reconstruction algorithm

# Jet origin id

Hao Liang, Yongfeng Zhu, Yuzhi Che, Yuexin Wang, Huiling Qu, Cen Zhou, etc

PHYSICAL REVIEW LETTERS **132**, 221802 (2024)

Eur. Phys. J. C (2024) 84:152  
<https://doi.org/10.1140/epjc/s10052-024-12475-5>

THE EUROPEAN  
PHYSICAL JOURNAL C



Regular Article - Experimental Physics

## Jet-Origin Identification and Its Application at an Electron-Positron Higgs Factory

Hao Liang<sup>1,2,\*</sup>, Yongfeng Zhu<sup>3,\*</sup>, Yuexin Wang<sup>1,4</sup>, Yuzhi Che<sup>1,2</sup>, Manqi Ruan<sup>1,2,†</sup>,  
Chen Zhou<sup>3,‡</sup> and Huiling Qu<sup>5,§</sup>

<sup>1</sup>Institute of High Energy Physics, Chinese Academy of Sciences, 19B Yuquan Road, Shijingshan District, Beijing 100049, China

<sup>2</sup>University of Chinese Academy of Sciences, 19A Yuquan Road, Shijingshan District, Beijing 100049, China

<sup>3</sup>State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

<sup>4</sup>China Center of Advanced Science and Technology, Beijing 100190, China

<sup>5</sup>CERN, EP Department, CH-1211 Geneva 23, Switzerland

(Received 16 October 2023; revised 26 April 2024; accepted 1 May 2024; published 31 May 2024)

To enhance the scientific discovery power of high-energy collider experiments, we propose and realize the concept of jet-origin identification that categorizes jets into five quark species ( $b, c, s, u, d$ ), five antiquarks ( $\bar{b}, \bar{c}, \bar{s}, \bar{u}, \bar{d}$ ), and the gluon. Using state-of-the-art algorithms and simulated  $\nu\bar{\nu}H, H \rightarrow jj$  events at 240 GeV center-of-mass energy at the electron-positron Higgs factory, the jet-origin identification simultaneously reaches jet flavor tagging efficiencies ranging from 67% to 92% for bottom, charm, and strange quarks and jet charge flip rates of 7%–24% for all quark species. We apply the jet-origin identification to Higgs rare and exotic decay measurements at the nominal luminosity of the Circular Electron Positron Collider and conclude that the upper limits on the branching ratios of  $H \rightarrow s\bar{s}, u\bar{u}, d\bar{d}$  and  $H \rightarrow sb, db, uc, ds$  can be determined to  $2 \times 10^{-4}$  to  $1 \times 10^{-3}$  at 95% confidence level. The derived upper limit for  $H \rightarrow s\bar{s}$  decay is approximately 3 times the prediction of the standard model.

## ParticleNet and its application on CEPC jet flavor tagging

Yongfeng Zhu<sup>1,a</sup>, Hao Liang<sup>2,3</sup>, Yuexin Wang<sup>2,3</sup>, Huiling Qu<sup>4</sup>, Chen Zhou<sup>1,b</sup>, Manqi Ruan<sup>2,3,c</sup>

<sup>1</sup> State Key Laboratory of Nuclear Physics and Technology, School of Physics, Peking University, Beijing 100871, China

<sup>2</sup> Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

<sup>3</sup> University of Chinese Academy of Sciences (UCAS), Beijing 100049, China



<sup>4</sup> EP Department, CERN, 1211 Geneva 23, Switzerland

Received: 15 November 2023 / Accepted: 23 January 2024  
© The Author(s) 2024

<https://arxiv.org/abs/2310.03440>



<https://arxiv.org/abs/2309.13231>

# Geo. & Tools



- **Jet origin identification: 11 categories (5 quarks + 5 anti quarks + gluon)**
  - Jet Flavor Tagging + Jet Charge measurements + s-tagging + gluon tagging...
- Full Simulated vvH, Higgs to two jets sample at CEPC baseline configuration: CEPC-v4 detector, reconstructed with **Arbor + ParticleNet (Deep Learning Tech.)**
- 1 Million samples each, 60/20/20% for training, validation & test

# Particle Net: IO




Variable	Definition
$\Delta\eta$	difference in pseudorapidity between the particle and the jet axis
$\Delta\phi$	difference in azimuthal angle between the particle and the jet axis
$\log p_T$	logarithm of the particle's $p_T$
$\log E$	logarithm of the particle's energy
$\log \frac{p_T}{p_T(jet)}$	logarithm of the particle's $p_T$ relative to the jet $p_T$
$\log \frac{E}{E(jet)}$	logarithm of the particle's energy relative to the jet energy
$\Delta R$	angular separation between the particle and the jet axis ( $\sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$ )
d0	transverse impact parameter of the track
d0err	uncertainty associated with the measurement of the d0
z0	longitudinal impact parameter of the track
z0err	uncertainty associated with the measurement of the z0
charge	electric charge of the particle
isElectron	if the particle is an electron
isMuon	if the particle is a muon
isChargedKaon	if the particle is a charged Kaon
isChargedPion	if the particle is a charged Pion
isProton	if the particle is a proton
isNeutralHadron	if the particle is a neutral hadron
isPhoton	if the particle is a photon

Table 3. The input variables used in ParticleNet for jet flavor tagging at the CEPC.

- Input: measurable information of all reconstructed jet particles
- Output: 10(11)-likelihoods to different categories



# 11-dim migration behavior

- Let the jet be identified as the category with highest likelihood:
- Pid: ideal Pid – three categories
  - Lepton identification
  - **Charged Kaon identification**
  - Neutral Kaon identification
- Patterns:
  - ~ Diagonal at quark sector...
  - $P(g \rightarrow q) < P(q \rightarrow g)$ ...
  - Light jet id...





# Performance with different PID scenarios & $H \rightarrow ss$ measurements




Flavor tagging: type that maximize  $\{L_q + L_{q\text{-bar}}, L_g\}$

If quark jet: jet charge  $\sim$  compare  $\{L_q, L_{q\text{-bar}}\}$

Remark: current jet flavor tagging efficiency & jet charge flip rates are projections of the 11-dim arrays produced by Jet origin id

# Benchmark analyses: Higgs rare/FCNC



Improved by ~3 times

Improved by 1-2 orders of magnitudes



Presumably... firstly quantified

TABLE I: Summary of background events of  $H \rightarrow b\bar{b}/c\bar{c}/gg, Z,$  and  $W$  prior to flavor-based event selection, along with the expected upper limits on Higgs decay branching ratios at 95% CL. Expectations are derived based on the background-only hypothesis.



	Bkg. ( $10^3$ )			Upper limit ( $10^{-3}$ )						
	$H$	$Z$	$W$	$s\bar{s}$	$u\bar{u}$	$d\bar{d}$	$sb$	$db$	$uc$	$ds$
$\nu\bar{\nu}H$	151	20	2.1	0.81	0.95	0.99	0.26	0.27	0.46	0.93
$\mu^+\mu^-H$	50	25	0	2.6	3.0	3.2	0.5	0.6	1.0	3.0
$e^+e^-H$	26	16	0	4.1	4.6	4.8	0.7	0.9	1.6	4.3
Comb.	-	-	-	0.75	0.91	0.95	0.22	0.23	0.39	0.86

- [28] J. Duarte-Campderros, G. Perez, M. Schlaffer, and A. Soffer. Probing the Higgs–strange-quark coupling at  $e^+e^-$  colliders using light-jet flavor tagging. *Phys. Rev. D*, 101(11):115005, 2020.
- [50] Alexander Albert et al. Strange quark as a probe for new physics in the Higgs sector. In *Snowmass 2021*, 3 2022.
- [59] J. de Blas et al. Higgs Boson Studies at Future Particle Colliders. *JHEP*, 01:139, 2020.
- [60] Jorge De Blas, Gauthier Durieux, Christophe Grojean, Jiayin Gu, and Ayan Paul. On the future of Higgs, electroweak and diboson measurements at lepton colliders. *JHEP*, 12:117, 2019.

# Performance V.S. Jet Kinematics





# Performance @ Z and Higgs





- *M10 instead of M11*

# V.S. Hadronization models



# Recent update at more benchmarks



- From Jet Flavor Tagging to Jet Origin ID:

- $\nu\nu H, H \rightarrow c\bar{c}$ : 3%  $\rightarrow$  1.7% (**Preliminary**)
- $V_{cb}$ : 0.75%  $\rightarrow$  0.45% ( $\mu\nu qq$  channel.  $e\nu qq$ : 0.6%, combined 0.4%)

# Updated result on $\sin^2 \theta_{eff}^l$ measurement

**Table 2.** Sensitivity  $S$  of different final state particles.


$\sqrt{s}/\text{GeV}$	$S$ of $A_{FB}^{e/\mu}$	$S$ of $A_{FB}^d$	$S$ of $A_{FB}^u$	$S$ of $A_{FB}^s$	$S$ of $A_{FB}^c$	$S$ of $A_{FB}^b$
70	0.224	4.396	1.435	4.403	1.445	4.352
75	0.530	5.264	2.598	5.269	2.616	5.237
92	1.644	5.553	4.200	5.553	4.201	5.549
105	0.269	4.597	1.993	4.598	1.994	4.586
115	0.035	3.956	1.091	3.958	1.087	3.942
130	0.027	3.279	0.531	3.280	0.520	3.261

**Table 3.** Cross section of process  $e^+e^- \rightarrow f\bar{f}$  calculated using the ZFITTER package. Values of the fundamental parameters are set as  $m_Z = 91.1875 \text{ GeV}$ ,  $m_t = 173.2 \text{ GeV}$ ,  $m_H = 125 \text{ GeV}$ ,  $\alpha_s = 0.118$  and  $m_W = 80.38 \text{ GeV}$ .

$\sqrt{s}/\text{GeV}$	$\sigma_\mu/\text{mb}$	$\sigma_d/\text{mb}$	$\sigma_u/\text{mb}$	$\sigma_s/\text{mb}$	$\sigma_c/\text{mb}$	$\sigma_b/\text{mb}$
70	0.039	0.032	0.066	0.031	0.058	0.028
75	0.039	0.047	0.073	0.046	0.065	0.043
92	1.196	5.366	4.228	5.366	4.222	5.268
105	0.075	0.271	0.231	0.271	0.227	0.265
115	0.042	0.135	0.122	0.135	0.118	0.132
130	0.026	0.071	0.068	0.071	0.066	0.069

Verify the RG behavior... using  
~1 month of data taking

**Expected statistical uncertainties on  $\sin^2 \theta_{eff}^l$  measurement.**  
(Using one-month data collection, ~ **4e12/24 Z events** at Z pole)




$\sqrt{s}$	$b$	$c$	$s$
70	$1.6 \times 10^{-5}$	$3.2 \times 10^{-5}$	$2.2 \times 10^{-5}$
75	$1.3 \times 10^{-5}$	$1.8 \times 10^{-5}$	$1.8 \times 10^{-5}$
92	$1.6 \times 10^{-6}$	$2.2 \times 10^{-6}$	$2.2 \times 10^{-6}$
105	$1.0 \times 10^{-5}$	$2.4 \times 10^{-5}$	$1.4 \times 10^{-5}$
115	$1.9 \times 10^{-5}$	$6.8 \times 10^{-5}$	$2.7 \times 10^{-5}$
130	$3.9 \times 10^{-5}$	$2.3 \times 10^{-4}$	$5.4 \times 10^{-5}$

# B-charge flip rate: Bs oscillations


Opposite side

- p charged Leptons with impact param.
- p charged Kaons with impact param.
- p charged pions with impact param.
- p protons with impact param. ?







# B-charge flip rate: Bs oscillations



- Flip rate  $\sim$  15%, Eff. Tagging power  $>$  40%



# Fast/Full Simulation

Z  $\rightarrow$   $\mu\mu$  (91.2 GeV)









- Delphes  $\sim$  Perfect PFA (1 – 1 correspondence.. )


# Arbor + AI: @ Boson Mass Resolution



# BMR: impact on critical measurements





# BMR decomposition @ CDR baseline



- 1<sup>st</sup>, Ultimate Precision  $\sim 2.8$  with CDR baseline
- 3<sup>rd</sup>, HCAL
- 2<sup>nd</sup>, HCAL resolution dominant the uncertainties from intrinsic detector resolution: *need better HCAL*
- 3<sup>rd</sup> Leading contribution: Confusion from shower Fragments (fake particles), *need better Pattern Reco.*

# Preliminary: Identify & veto charged shower fragments using AI





Trained at 12E4 events,

Test & Applied at 4E4 events



**score > 0.75**  
**efficiency ~83%**  
**purity ~95%**

# ... At Bosons ...



# 1-1 correspondence: preliminary

nCluHit != 0 & E > 1 GeV & |cosθ| < 0.9



- Next step: to improve the neutral hadron reco & to optimize the detector configuration





# Summary

- Higgs factory: immense science merit...
- Jet origin id: efficiently separate different species of colored SM particle
  - A “game changer” and opens new horizon for precise flavor studies at all future experiments
- Significantly impact on physics
  - Higgs: improve  $H \rightarrow ss, uu, dd, sb, uc, sd, db$  by 3-100 times, and  $H \rightarrow cc$  by 2 times
  - Flavor: Improve  $V_{cb}$  precision by  $\sim 50\%$ , effective tagging power for b-jet  $> 40\%$ ...
  - EW: Weak mixing angle...
  - QCD: Fragmentation relevant - **Road Map wanted**: towards better hadronization models + experimental validation (from both current data + GigaZ + TeraZ) + applications
  - NP: ...
- AI @ PFA: significantly reduce the confusing... and towards 1-1 correspondence reco.
- Long term version: 'see' gluon + quarks, as we see photon + leptons


# Back up

# V.S. Hadronization models




# b-jet: leading b-hadrons & flip rates


Percentage of b hadrons by Whizard & Herwig




Charge Flip Rate  $\omega$  of b hadrons by Whizard & Herwig



Difference in Percentage of b hadrons between Whizard and Herwig




Difference in Charge Flip Rate  $\omega$  of b hadrons between Whizard and Herwig




# c-jet: leading c-hadrons & flip rates


Percentage of c hadrons by Whizard & Herwig




Charge Flip Rate ω of c hadrons by Whizard & Herwig



Difference in Percentage of c hadrons between Whizard and Herwig




Difference in Charge Flip Rate ω of c hadrons between Whizard and Herwig




# s-jet: leading s-hadrons & flip rates


Percentage of s hadrons by Whizard & Herwig




Charge Flip Rate  $\omega$  of s hadrons by Whizard & Herwig




Difference in Percentage of s hadrons between Whizard and Herwig




Difference in Charge Flip Rate  $\omega$  of s hadrons between Whizard and Herwig




## M11 2 with charged hadron



## M11 3 with charged hadron and $K_L$ $K_S$



# Arbor PFA: Towards one-to-one correspondence (Totoro)





# Arbor

## Tree topology of particle shower

THE EUROPEAN  
PHYSICAL JOURNAL C



Eur. Phys. J. C (2018) 78:426  
<https://doi.org/10.1140/epjc/s10052-018-5876-z>

Special Article - Tools for Experiment and Theory

### Reconstruction of physics objects at the Circular Electron Positron Collider with Arbor

Manqi Ruan<sup>1,a</sup>, Hang Zhao<sup>1</sup>, Gang Li<sup>1</sup>, Chengdong Fu<sup>1</sup>, Zhigang Wang<sup>1</sup>, Xinchou Lou<sup>6,7,8</sup>, Dan Yu<sup>1,2</sup>, Vincent Boudry<sup>2</sup>, Henri Videau<sup>2</sup>, Vladislav Balagura<sup>2</sup>, Jean-Claude Brient<sup>2</sup>, Peizhu Lai<sup>3</sup>, Chia-Ming Kuo<sup>3</sup>, Bo Liu<sup>1,4</sup>, Fenfen An<sup>1,4</sup>, Chunhui Chen<sup>4</sup>, Soeren Prell<sup>4</sup>, Bo Li<sup>5</sup>, Imad Laketineh<sup>5</sup>

<sup>1</sup> Institute of High Energy Physics, Beijing, China

<sup>2</sup> Laboratoire Leprince-Ringuet, Ecole Polytechnique, Palaiseau, France

<sup>3</sup> Department of Physics and Center of high energy and high field physics, National Central University, Taoyuan City, Taiwan


<sup>4</sup> Iowa State University, Ames, USA

<sup>5</sup> Institut de Physique Nucleaire de Lyon, Lyon, France

<sup>6</sup> Institute of High Energy Physics, Chinese Academy of Sciences, Beijing, China



<sup>7</sup> Physics Department, University of Texas at Dallas, Richardson, TX, USA

<sup>8</sup> University of Chinese Academy of Sciences (UCAS), Beijing, China



20 GeV Klong reconstructed @ ILD Calo  
Curves indicating expected particle  
trajectories (from MC-truth)

# Validation: Arbor Branch Length Vs MC Truth



Arbor: successfully **tag** sub-shower structure

*Samples: Particle gun event at ILD HCAL (readout granularity  $1\text{cm}^2$  & layer thickness  $2.65\text{cm}$ )*

*Length:*

*Charged MCParticle: spatial distance between generation/end points*

*Arbor branch: sum of distance between neighboring cells*

$Z \rightarrow 2 \text{ muon},$   
 $H \rightarrow 2 b$   
 $\sim 2\%$


$Z \rightarrow 2 \text{ jet},$   
 $H \rightarrow 2 \text{ tau}$   
 $\sim 5\%$

$ZH \rightarrow 4 \text{ jets}$   
 $\sim 50\%$



$Z \rightarrow 2 \text{ muon}$   
 $H \rightarrow WW^* \rightarrow eevv$   
 $\sim 1\%$



CMS Experiment at LHC, CERN  
Data recorded: Thu Jan 1 01:00:00 1970 CEST  
Run/Event: 1 / 1201  
Lumi section: 13



# V.S. Multiplicity










- ...many patterns need further understanding & towards further optimization...*



# PID studies with beamtest data

Xin Xia (IHEP)

- FD characteristics of different beam particles
  - Imaging capability of high granularity calorimeter ( )



Oct 19 - Nov 2, 2022


Apr 26 - May 10, 2023

May 17 - 31, 2023





SPS H8 beamline

SPS H2 beamline

PS T9 beamline



# Lepton: isolated






*BDT method using 4 classes of 24 input discrimination variables.*

Test performance at: Electron =  $E_{\text{likeness}} > 0.5$  ;  
 Muon =  $Mu_{\text{likeness}} > 0.5$

Single charged reconstructed particle, for  $E > 2$  GeV:  
 lepton efficiency  $> 99.5\%$  && Pion mis id rate  $\sim 1\%$

<https://link.springer.com/article/10.1140/epjc/s10052-017-5146-5>  
 CEPC-DocDB-id:148, Eur. Phys. J. C (2017) 77: 591

# Lepton: inside jet





Compared the single particle sample, the jet lepton (at  $Z \rightarrow b\bar{b}$  sample at  $\sqrt{s} = 91.2$  GeV) Performance will be slightly degraded – Due to the limited clustering performance (splitting & contamination).

At the same working point, the efficiency can be reduced by up to 3%; while mis-id rate increases up to 1%. Marginal Impact on Flavor Physics measurements as  $B_c \rightarrow \tau \nu$ .



# Kaon




Highly appreciated in flavor physics @ CEPC Z pole  
 TPC dEdx + ToF of 50 ps

At inclusive Z pole sample:

Conservative estimation gives efficiency/purity of 91%/94% (2-20 GeV, 50% degrading +50 ps ToF)  
 Could be improved to 96%/96% by better detector/DAQ performance (20% degrading + 50 ps ToF)



# Pid performance



	factor	1.	1.2	1.5	2.
dE/dx	$\epsilon_K$ (%)	95.97	94.09	91.19	87.09
dE/dx	$pur_{K^\pm}$ (%)	81.56	78.17	71.85	61.28
dE/dx & TOF	$\epsilon_K$ (%)	98.43	97.41	95.52	92.3
dE/dx & TOF	$pur_{K^\pm}$ (%)	97.89	96.31	93.25	87.33




3% of dE/dx & dN/dx + 50 ps ToF: eff/purity of Kaon reco > 95%

# Neutral Particle id: Very Preliminary




- Fast Sim Prediction: BMR: 2.9  $\rightarrow$  2.6
  - Need excellent CALO + ToF  $\sim$  o(10 ps)
  - Need high efficiency neutral hadron reco (1-1 correspondence)

# 2-body decay particles and tau leptons




Eur. Phys. J. Plus (2020) 135:274

Page 9 of 14 274




$\pi_0$ : 60/30 GeV  
with 5/10 mm cell.

Kshort, Lambda,  
Phi, Tau, D meson...



(a)  $K_S^0$



(b)  $\Lambda$

Fig. 7 All reconstructed mass distributions of  $K_S^0$  and  $\Lambda$ . They are fitted with double-sided crystal ball functions

08/07/2004