





# Jet flavor identification for FCCee

Michele Selvaggi  
(CERN)


Credits to: Andrea Del Vecchio, Dolores Garcia, Laurent Forthomme,  
Franco Bedeschi, Michele Selvaggi, Loukas Gouskos  
[EPJ C 82 646 (2022) [link](#)]

# Physics motivation

- Flavour tagging essential for the  $e^+e^-$  program, e.g.:
  - **Higgs Sector:**
    - (HL-)LHC can access 3<sup>rd</sup> gen. couplings and a few of 2<sup>nd</sup> generation
    - Future  $e^+e^-$ : Measure Higgs particle properties and interactions in challenging decay modes
      - E.g.  $cc$ , 1<sup>st</sup> gen quarks/fermions,  $gg$  [?]
  - **Top quark physics [if  $E_{CM}$  sufficient]**
    - Precise determination of top properties [mass, width, Yukawa]
  - **QCD Physics**
    - strong coupling ( $a_s$ ), event shapes ..
    - modelling of hadronization, MC tuning, ...
  - .....



# Basics of flavour tagging (b/c)



- Large lifetime
  - $b$  ( $c$ ) lifetime  $\sim 1$  ps ( $\sim 0.1$ ps)
  - $b$  ( $c$ ) decay length:  $\sim 500$   $\mu\text{m}$
  - (2-3) mm for  $\sim 50$  GeV boost
  
- Displaced vertices/tracks
  - Large impact parameters
  - Tertiary vertices when B hadron decays to C hadron
  
- Large track multiplicity
  - $\sim 5$  ( $\sim 2$ ) charged tracks/decay
  
- Presence of non-isolated  $e/\mu$ 
  - $\sim 20$  (10)% in B (C) decays

## Detector constraints:

Need power pixel/tracking detectors


- Good spatial resolution
- As little material as possible
- Precise track alignment

# Input variables

- Comparison of input distributions for different jet flavors


## Constituent relative energy

FCC-ee simulation (Delphes)



## Impact parameter ( $d_0$ )


FCC-ee simulation (Delphes)




# Basics of flavour tagging (strange)

[2003.09517]


Momentum weighted fraction:



Strange  $p_T = 45$  GeV



Down  $p_T = 45$  GeV



- Large Kaon content

- Charged Kaon as track:

- K/pi separation
  - TOF
  - $dE/dx/dNdx$

- Neutral Kaons:





- $K_S \rightarrow \pi\pi$ 
  - Displaced 2 track vertex
  - 4 photons
- $K_L$ 
  - TOF vs n ?

**Detector constraints:**

- timing detectors
- charged energy loss (gas/silicon)
- cherenkov detectors

# Particle ID: dN/dx and ToF

- Count number of **primary ionization** clusters along track path
- ToF results in good K/ $\pi$  separation at low-momenta
- Modules added in Delphes





# Designing a Graph-based tagger

- **Jet representation:** critical for powerful jet tagging algorithms
  - **In theory:** A spray of particles produced by the hadronization of  $q$  and  $g$
  - **Experimentally:** A cone of reconstructed particles in the detector
- Reminder: Current and future experiments have / will have a **PF-based** event reconstruction
  - **Output:** mutually exclusive list of particles
    - Rich set of info/particle
      - Energy/momentum, position
      - Displacement, particle type
      - timing
      - ...
- **Until recently:** Jet taggers based on human-inspired higher-level observables
  - Inputs to cut-based or simple ML-based algorithms
- Move to **particle-based jet tagging:** i.e. exploit directly the full list of jet constituents (ReconstructedParticles) and **new advances in ML**

[O(50) properties/particle]  
x [~50-100 particles/jet]  
~O(1000) inputs/jet



# Full list of input variables


Variable	Description
Kinematics	
$E_{\text{const}}/E_{\text{jet}}$	energy of the jet constituent divided by the jet energy
$\theta_{\text{rel}}$	polar angle of the constituent with respect to the jet momentum
$\phi_{\text{rel}}$	azimuthal angle of the constituent with respect to the jet momentum
Displacement	
$d_{xy}$	transverse impact parameter of the track
$d_z$	longitudinal impact parameter of the track
$\text{SIP}_{2\text{D}}$	signed 2D impact parameter of the track
$\text{SIP}_{2\text{D}}/\sigma_{2\text{D}}$	signed 2D impact parameter significance of the track
$\text{SIP}_{3\text{D}}$	signed 3D impact parameter of the track
$\text{SIP}_{3\text{D}}/\sigma_{3\text{D}}$	signed 3D impact parameter significance of the track
$d_{3\text{D}}$	jet track distance at their point of closest approach
$d_{3\text{D}}/\sigma_{d_{3\text{D}}}$	jet track distance significance at their point of closest approach
$C_{ij}$	covariance matrix of the track parameters
Identification	
$q$	electric charge of the particle
$m_{\text{t.o.f.}}$	mass calculated from time-of-flight
$dN/dx$	number of primary ionisation clusters along track
<code>isMuon</code>	if the particle is identified as a muon
<code>isElectron</code>	if the particle is identified as an electron
<code>isPhoton</code>	if the particle is identified as a photon
<code>isChargedHadron</code>	if the particle is identified as a charged hadron
<code>isNeutralHadron</code>	if the particle is identified as a neutral hadron



# ParticleNet(-ee)

H. Qu and LG  
 PRD 101 056019 (2020)  
 F. Bedeschi, M. Selvaggi, LG  
 EPJ C 82 646 (2022)

- Jet representation: “*Point Cloud*” → “*Particle Clouds*”
  - Treat the jet as an unordered set of particles
- Algorithm design: Graph Neural Networks
  - Particle cloud represented as a graph
    - Each particle: **node** of the graph; Connections between particles: the **edges**
- Follow a hierarchical learning approach
  - First learn local structures → then move to more global ones





# ParticleNet in FCCSW


## Sample Generation for training

- Generation of the samples in EDM4hep (whole event reconstructions, features for training not explicit)
- FCCAnalyses (wrapper RDataFrame)
  - Per-event → per-jet structure
  - 2 stages. 1: read edm4hep and extract features. 2: produce n-tuples one per class.
  - final dataset: 5/7 classes and  $10^6$  events per class
  - trained on gpus (A100 )


## Inference

- Inference in FCCAnalyses:
  - load ONNX training files
  - Extract hard vertex and perform jet clustering
  - Extract jet constituents and compute observables
  - Evaluate NN → output: **one probability per category**

# Training the model



# Inference





# Inference with FCCAnalyses

```
## get local file, else download from url
weaver_preproc = get_file_path(url_preproc, local_preproc)
weaver_model = get_file_path(url_model, local_model)

from addons.ONNXRuntime.python.jetFlavourHelper import JetFlavourHelper
from addons.FastJet.python.jetClusteringHelper import ExclusiveJetClusteringHelper
```

Loading model parameters

```
## define jet clustering parameters
jetClusteringHelper = ExclusiveJetClusteringHelper(collections["PFParticles"], njets, tag)
```

JetClusteringHelper

Jet clustering

```
## run jet clustering
df = jetClusteringHelper.define(df)
```

```
## define jet flavour tagging parameters
```

```
jetFlavourHelper = JetFlavourHelper(
    collections,
    jetClusteringHelper.jets,
    jetClusteringHelper.constituents,
    tag,
)
```

JetFlavourHelper

Obtain input parameters

```
## define observables for tagger
df = jetFlavourHelper.define(df)
```


```
## tagger inference
df = jetFlavourHelper.inference(weaver_preproc, weaver_model, df)
```

```
return df
```

Run inference


# ParticleNet@FCCee: b/c tagging

## b-tagging



WP	Eff (b)	Mistag (g)	Mistag (ud)	Mistag (c)
Loose	90%	2%	0.1%	2%
Medium	80%	0.7%	<0.1%	0.3%

## c-tagging





WP	Eff (c)	Mistag (g)	Mistag (ud)	Mistag (b)
Loose	90%	7%	7%	4%
Medium	80%	2%	0.8%	2%

# ParticleNet@FCCee: s/g tagging

## strange-tagging

## gluon-tagging




WP	Eff (s)	Mistag (g)	Mistag (ud)	Mistag (c)	Mistag (b)
Loose	90%	20%	40%	10%	1%
Medium	80%	9%	20%	6%	0.4%

WP	Eff (g)	Mistag (ud)	Mistag (c)	Mistag (b)
Loose	90%	25%	7%	2.5%
Medium	80%	15%	5%	2%


# Impact of detector configurations

## Strange tagging [PID]



- dN/dx brings most of the gain
- additional gain w/ TOF (30ps)
  - TOF (3ps): marginal improvement
  - dN/dX + TOF(30ps) ~ perfect PID

## c-tagging [PIX layers]



- Additional pixel layer 1 cm from beam pipe vs 1.5 cm:
  - improved BKG rejection in c-tagging
  - marginal/no improvement in b-tagging




# Robustness

- ParticleNet-ee trained using *Pythia 8* samples
  - tested on *Pythia 8* [solid lines]
  - tested on *WZ-Pythia6* [dashed lines]


## b-tagging

FCC-ee simulation (IDEA)



## gluon -tagging


FCC-ee simulation (IDEA)




- Modest dependence on choice of generator
- More parton showers coming up (Herwig, Sherpa...)

# Tagger update(up and down)

## Up -tagging



## Down-tagging



- Up vs Down discrimination seems possible thanks to jet charge
- 30% bkg eff at 50% signal (better than random coin toss)

# Summary

- Powerful jet flavour identification important for the  $e^+e^-$  physics program
- Sophisticated jet tagging algorithms developed for  $e^+e^-$  experiments
  - Striking improvement in tagging performance compared to previous tools
    - allows us to explore more of the detector and event reconstruction potential
  - Integrated in FCCSW [data preparation, training, validation, inference, analysis] and used in FCCee physics analyses
- Still room for improvement / other ideas to try:
  - secondary tasks, secondary vertexing regression
  - new higher order graph architectures
  - improve explainability
  - resilience to modelling (more generators)
  - calibration (Z pole  $\rightarrow$  ZH threshold extrapolation)

