



1

2 **ILD sensitivity to light scalar production with invisible decays**

3

Kamil Zembaczyński*, Aleksander Filip Żarnecki*

4

* *Univeristy of Warsaw*

5

While the main goal of the electron-positron Higgs factory is to measure precisely the Higgs boson properties, production of new exotic light scalars is still not excluded by the existing experimental data, provided their coupling to the gauge bosons is sufficiently suppressed. In this contribution sensitivity of the future Higgs factory experiment to exotic light scalar particle production in association with a Z boson, assuming invisible scalar decays is discussed. This work was carried out in the framework of the ILD concept group as a contribution to the New exotic scalars (EXscalar) focus topic.

6

7

8

9

10

11

12 1 Analysis framework

13 Presented study is based on the framework developed previously for the search of light exotic scalar production with decays into two taus [1]. Considered is scalar production in the scalar-strahlung process, $e^+e^- \rightarrow ZS$, with hadronic Z decays (for highest sensitivity) and S decays into invisible final state (eg. dark sector). Samples of background and signal events were generated with WHIZARD 3.1.2 [2] using built-in SM_CKM model. For signal events Higgs boson mass was varied in the model and only its decay into four neutrinos was allowed. For the background, all relevant 4 and 2-fermion final states were considered including contribution from SM-like Higgs boson as well as from processes with beamstrahlung or EPA photons in the initial state. ISR and luminosity spectra for ILC running at 250 GeV were taken into account. H-20 running scenario [3, 4] for ILC was assumed with $\pm 80\%$ and $\pm 30\%$ polarisation for e^- and e^+ beam respectively. Fast simulation of detector response was performed with Delphes ILCgen model [5, 6].

23 2 Pre-selection

24 Two stage pre-selection was used to select events consistent with the expected signal signature. First, rejected were all events with isolated leptons or photons reconstructed in the detector (including also forward calorimeters, LumiCal and BeamCal). All reconstructed objects were then clustered into two jets (using Durham algorithm in exclusive mode). Selected for the analysis were events with reconstructed di-jet invariant mass in the ± 20 GeV window around the Z mass and the missing transverse momentum greater than 10 GeV. Distributions of the considered variables and the applied pre-selection kinematic cuts are presented in Figure 1(a) and Figure 1(b). Figure 1(c) presents invariant mass recoiling against Z boson candidate before and after the cuts. Pre-selection significantly improves signal to background ratio and thus makes training ML classifiers faster and more precise.




Figure 1: Distributions of reconstructed variables for background (black) and signal (blue) events for scalar mass of 50 GeV and $e_L^- e_R^+$ polarisation: (a) di-jet invariant mass with pre-selection acceptance region indicated, (b) missing transverse momentum with pre-selection cut indicated and (c) recoil mass before pre-selection (dashed lines) and after pre-selection cuts (solid lines).

33 3 Reconstruction and classification

34 Following input variables were used for event classification: di-jet (Z candidate) invariant mass and energy, missing transverse momentum, cosine of the Z polar angle, angle between two jets, recoil mass (corresponding to S mass for signal events), y_{23} and y_{34} variables from clustering algorithm. For most efficient event classification, separate Boosted Decision Tree [7] classifiers were trained for each beam polarisation configuration, resulting in four BDTs for each considered scalar mass. Background sample with $q\bar{q}l\nu$ in the final state was found to be the dominant background source.

40 4 Results

41 The expected limits on exotic scalar production cross section, calculated from the template fit to the BDT
 42 response distributions, are presented in figure Figure 2(a). One should note that $e_R^- e_L^+$ running is expected
 43 to result significantly better limits than $e_L^- e_R^+$ running with the same luminosity. This is expected from the
 44 suppression of the $W^+ W^-$ production, which contribute to the main background channel. Similar limits are
 45 obtained with 900 fb^{-1} collected with preferred polarization configuration as for 2 ab^{-1} with unpolarized beam.
 46 Combined analysis of the four polarisation configurations, according to H-20 running scenario, results in about
 47 20% improvement with respect to the same integrated luminosity with the unpolarized beams.




Figure 2: Expected 95% C.L. cross section limits on the light scalar production cross section times invisible branching ratio for ILC running at 250 GeV, assuming H-20 running scenario. (a) Comparison of combined limits with limits obtained for different polarisation configurations. (b) Comparison of combined limits with and without polarisation and with and without systematic effects.

48 To test possible impact of systematical effect on the extracted cross section limits nine additional nuisance
 49 parameters were introduced in the template fit procedure: four describing luminosity uncertainties of the
 50 data samples (for four polarization settings) and five for the uncertainties in the theoretical predictions (for
 51 $\bar{q}q$, $W^- W^+$, ZZ , $e^\pm \gamma$ and $\gamma\gamma$ processes). Impact of systematic uncertainties is significant, as indicated in
 52 Figure 2(b), especially for the region of scalar masses around W^\pm and Z boson masses. The impact of
 53 systematic effects is also slightly bigger for unpolarized beam than for polarised one, as expected.

54 5 Conclusion

55 Sensitivity of the ILD detector operating at the 250 GeV ILC for discovering new exotic scalar particles was
 56 studied for the hypothesis of invisible new scalar decays. Hadronic decays of the Z boson produced in as-
 57 sociation with the new scalar were considered for maximum sensitivity. The expected 95% C.L. limits on the
 58 scalar production cross section times the invisible branching ratio, relative to the SM production cross section,
 59 were calculated in the mass range between 20 and 120 GeV. Results correspond to about order of magnitude
 60 increase in sensitivity with respect to the model independent study [8], assuming invisible decays of the new
 61 scalar dominate. Running with polarized beams, according to H-20 running scenario, results in about 20%
 62 limit improvement compared to running with unpolarised beam. Impact of systematic uncertainties was found
 63 to be significant and should be studied in more detail.

64 **6 References**

- 65 [1] B. Brudnowski, K. Zembaczyński, A. F. Żarnecki,
66 *Prospects for light exotic scalar measurements at the e^+e^- Higgs factory*,
67 International Workshop on Future Linear Colliders, 2024, arXiv: 2409.19761 [hep-ph].
- 68 [2] J. Reuter et al., *Modern Particle Physics Event Generation with WHIZARD*,
69 J. Phys. Conf. Ser. **608** (2015), ed. by L. Fiala, M. Lokajicek, N. Tumova 012063,
70 DOI: 10.1088/1742-6596/608/1/012063, arXiv: 1410.4505 [hep-ph].
- 71 [3] T. Barklow et al., *ILC Operating Scenarios* (2015), arXiv: 1506.07830 [hep-ex].
- 72 [4] P. Bambade et al., *The International Linear Collider: A Global Project* (2019),
73 arXiv: 1903.01629 [hep-ex].
- 74 [5] J. de Favereau et al., DELPHES 3,
75 *DELPHES 3, A modular framework for fast simulation of a generic collider experiment*,
76 JHEP **02** (2014) 057, DOI: 10.1007/JHEP02(2014)057, arXiv: 1307.6346 [hep-ex].
- 77 [6] A. F. Żarnecki, *ILCDelphes*, <https://github.com/zarnecki/ILCDelphes>, 2020.
- 78 [7] T. Chen, C. Guestrin, *XGBoost: A Scalable Tree Boosting System*, Proceedings of the 22nd ACM
79 SIGKDD International Conference on Knowledge Discovery and Data Mining, KDD '16, ACM, 2016,
80 DOI: 10.1145/2939672.2939785,
81 URL: <http://dx.doi.org/10.1145/2939672.2939785>.
- 82 [8] Y. Wang, International Large Detector Concept Group,
83 *Search for Light Scalars Produced in Association with a Z boson at the 250 GeV stage of the ILC*,
84 PoS **ICHEP2018** (2019) 630, DOI: 10.22323/1.340.0630.