

# Off-Shell and Interference Effects for SUSY Particle Production

Jürgen Reuter

University of Freiburg



Hagiwara/Kilian/Krauss/Ohl/Plehn/Rainwater/JR/Schumann **PRD 73** (2006), 055005;  
JR et al., hep-ph/0512012

DESY, June 1st, 2007

# SUSY Precision measurements

Motivation for SUSY: see Tuesday's Symposium

Analysis Goal:

- ▶ Mass measurements to get the spectrum
- ▶ Access spin of all new particles: angular/spin correlations
- ▶ Coupling measurements: verify SUSY by the structure of couplings

Precise predictions for SUSY processes:

**background** to other (more difficult) SUSY processes

Precise parameter values: Reverse the renormalization-group evaluation and **get a handle on GUT parameters** ( $\Rightarrow$  P. Zerwas' talk)

$\Rightarrow$  **SPA project**    <http://spa.desy.de/spa>

# Classification of corrections to (SUSY) processes

Corrections to the SUSY processes fall into six categories:

- ▶ **Loop corrections to SUSY production and decay processes**

Kilian/JR/Robens, EPJ C48 (2006), 389, see T. Robens' talk

- ▶ nonfactorizable, maximally resonant photon exchange between production and decay

- ▶ **real radiation of photons** [gluons]

Kilian/JR/Robens, EPJ C48 (2006), 389, see T. Robens' talk

- ▶ **off-shell kinematics for the signal process** see also Berdine/Rainwater/Kauer, 2007

- ▶ **irreducible background from all other SUSY processes**

- ▶ **reducible, experimentally indistinguishable SM background processes**

# Classification of approximations in (SUSY) processes

Some generic SUSY process:

$$e^+e^- \rightarrow b\bar{b}e^+e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0 \quad \text{66478 diagrams.} \quad (\text{It's just } e^+e^- \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_2^0!)$$

- ▶ Entanglement of different signal diagrams ( $e^+e^- \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0, \tilde{b}_i \tilde{b}_j, \tilde{e}_i \tilde{e}_j$ )
- ▶ **Need for cuts** to disentangle those (experimentally/simulation)
- ▶ Add SM backgrounds ( $e^+e^- \rightarrow b\bar{b}e^+e^- \nu_i \bar{\nu}_i$ )
- ▶ **Much more complicated processes for LHC, and even also for ILC**

Process  $A_1 A_2 \rightarrow P^{(*)} \rightarrow F_1 F_2$ , 3 different levels:

<b>Narrow width</b>	$\sigma(A_1 A_2 \rightarrow P) \times \text{BR}(P \rightarrow F_1 F_2)$
<b>Breit-Wigner</b>	$\sigma(A_1 A_2 \rightarrow P) \times \frac{M_P^2 \Gamma_P^2}{(s - M_P^2)^2 + \Gamma_P^2 M_P^2} \times \text{BR}(P \rightarrow F_1 F_2)$
<b>Full matrix element</b>	$\sigma(A_1 A_2 \rightarrow F_1 F_2)$

last level not featured by ISAJET, PYTHIA, HERWIG, SUSYGEN

# The generator generator O'Mega $\Omega$ / Whizard



## Matrix Element Generator O'Mega:

Ohl, 2000/01; M.Moretti/Ohl/JR, 2001; JR, 2002

Optimized helicity amplitudes: avoiding all redundancies

## Multi-purpose Event Generator Whizard:

Ohl, 1996; Kilian, 2000; Kilian/Ohl/JR, 2007

- Multi-Channel adaptive Monte-Carlo integration
- Generator generator for arbitrary multi-particle processes
- Well-suited for ILC physics (ISR, beamstrahlung); used for ILC reference event files
- **New release this summer: Whizard 2.0/O'Mega 1.0** (LHC approaching!!)
  - ▶ Fancier support for full color flows
  - ▶ LHAPDF support
  - ▶ new BSM models: extMSSM, ext.Dim., Little Higgs, NCSM
  - ▶ new syntax for arbitrary cut functions
- Virtual (SUSY) Corrections (all  $2 \rightarrow 2$  processes for ILC)
- Future features: Parton Shower/Matrix Element matching

# Tests and Checks of MSSM implementation



JR et al., 2005; Hagiwara/Kilian/Krauss/Ohl/Plehn/Rainwater/JR/Schumann, 2006

- MSSM: doubled spectrum, 100 parameters, 5000 vertices
- Unitarity checks:  $\sigma(2 \rightarrow 2, s), \sigma(2 \rightarrow 3, s) \sim \text{const}$  or  $1/s$  ✓
- Gauge invariance: Ward- and Slavnov-Taylor identities ✓
- Supersymmetry: Ward-/Slavnov-Taylor identities ✓ JR, 2002; Ohl/JR, 2002
- Comparison of codes ( $\mathcal{O}(600)$  processes): JR et al., 2005; K. Hagiwara/.../JR/..., 2006

Reference: [http://james.physik.uni-freiburg.de/~reuter/susy\\_comparison.html](http://james.physik.uni-freiburg.de/~reuter/susy_comparison.html)

		$\tau^+\tau^- \rightarrow X$					
Process	status	Madgraph/Helas		Whizard/O'Mega		Sherpa/A'Megic	
		0.5 TeV	2 TeV	0.5 TeV	2 TeV	0.5 TeV	2 TeV
$\tilde{\tau}_1 \tilde{\tau}_1^*$	●	257.57(7)	79.63(4)	257.32(1)	79.636(4)	257.30(1)	79.638(4)
$\tilde{\tau}_2 \tilde{\tau}_2^*$	●	46.55(1)	66.86(2)	46.368(2)	66.862(3)	46.372(2)	66.862(3)
$\tilde{\tau}_1 \tilde{\tau}_2^*$	●	95.50(3)	19.00(1)	94.637(3)	19.0015(8)	94.645(5)	19.000(1)
$\tilde{\nu}_\tau \tilde{\nu}_\tau^*$	●	502.26(7)	272.01(8)	502.27(2)	272.01(1)	502.30(3)	272.01(1)
$\tilde{\chi}_1^0 \tilde{\chi}_1^0$	●	249.94(2)	26.431(1)	249.954(9)	26.431(1)	249.96(1)	26.431(1)
$\tilde{\chi}_1^0 \tilde{\chi}_2^0$	●	69.967(3)	9.8940(3)	69.969(2)	9.8940(4)	69.968(3)	9.8937(5)
$\tilde{\chi}_1^0 \tilde{\chi}_3^0$	●	17.0387(3)	0.7913(1)	17.0394(1)	0.79136(2)	17.040(1)	0.79137(5)
$\tilde{\chi}_1^+ \tilde{\chi}_1^-$	●	185.09(3)	45.15(1)	185.093(6)	45.147(2)	185.10(1)	45.151(2)
$h^0 h^0$	●	0.3533827(3)	0.0001242(2)	0.35339(2)	0.00012422(3)	0.35340(2)	0.000124218(6)
$A^0 A^0$	●	—	0.07975(3)	—	0.079758(6)	—	0.079744(4)
$Z h^0$	●	59.591(3)	3.1803(8)	59.589(3)	3.1802(1)	59.602(3)	3.1829(2)
$Z A^0$	●	2.9915(4)	4.682(5)	2.99162(9)	4.6821(3)	2.9917(2)	4.6817(2)

## Parameter point under consideration

Following discussions do not depend on the special parameter point  
 SUGRA-inspired point, non-universal right-handed scalar masses  
 $\tan \beta = 20$

Particle	$M$ [GeV]	$\Gamma$ [GeV]	Particle	$M$ [GeV]	$\Gamma$ [GeV]
$h$	114.45	0.0050	$\tilde{\chi}_1^0$	46.84	—
$H$	300.15	2.2924	$\tilde{\chi}_2^0$	112.41	0.00005
$A$	300.00	2.7750	$\tilde{\chi}_3^0$	148.09	0.01162
$H^\pm$	310.96		$\tilde{\chi}_4^0$	236.77	1.0947
$\tilde{b}_1$	295.36	0.5395	$\tilde{\chi}_{1,2}^\pm$	106.60	
$\tilde{b}_2$	399.92	3.4956	$\tilde{\chi}_2^\pm$	237.25	
$\tilde{e}_L$	205.02		$\tilde{t}_1$	413.84	
$\tilde{e}_R$	205.65		$\tilde{t}_2$	978.88	

- ▶ (Very) light Higgs, directly above LEP limit
- ▶  $h \sim 47\%$  invisible decays to LSP
- ▶  $m_{\tilde{q}} \sim 430$  GeV
- ▶ Light sbottoms accessible at the ILC
- ▶ Low-energy data-compatible:  $b \rightarrow s\gamma$ ,  $B_s \rightarrow \mu^+\mu^-$ ,  $\Delta\rho$ ,  $g_\mu - 2$ , CDM
- ▶ Focus on  $\text{BR}(\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0) = 43.2\%$

# Sbottom production at the ILC

Cross sections for  $\sqrt{s} = 800$  GeV

- ▶ In contrast to the LHC: Electroweak production
- ▶ More channels contribute to  $e^+e^- \rightarrow b\bar{b}\tilde{\chi}_1^0\tilde{\chi}_1^0$ :  
 $e^+e^- \rightarrow Zh, ZH, Ah, HA, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_1^0\tilde{\chi}_3^0, \tilde{\chi}_1^0\tilde{\chi}_4^0, \tilde{b}_1\tilde{b}_1^*, \tilde{b}_1\tilde{b}_2^*$  (412 diagrams)
- ▶ Irreducible SM background:  $e^+e^- \rightarrow b\bar{b}\nu_i\bar{\nu}_i$  ( $WW$  fusion,  $Zh, ZZ$ ) (47 diagrams)

Channel	$\sigma_{2\rightarrow 2}$ [fb]	$\sigma \times \text{BR}$ [fb]	$\sigma_{\text{BW}}$ [fb]
$Zh$	20.574	1.342	1.335
$ZH$	0.003	0.000	0.000
$hA$	0.002	0.001	0.000
$HA$	5.653	0.320	0.314
$\tilde{\chi}_1^0\tilde{\chi}_2^0$	69.109	13.078	13.954
$\tilde{\chi}_1^0\tilde{\chi}_3^0$	24.268	3.675	4.828
$\tilde{\chi}_1^0\tilde{\chi}_4^0$	19.337	0.061	0.938
$\tilde{b}_1\tilde{b}_1^*$	4.209	0.759	0.757
$\tilde{b}_1\tilde{b}_2^*$	0.057	0.002	0.002
Sum		19.238	22.129
Exact w/ISR			19.624 22.552

Channel	$\sigma_{2\rightarrow 2/3}$ [fb]	$\sigma \times \text{BR}$ [fb]	$\sigma_{\text{BW}}$ [fb]
$ZZ$	202.2	12.6	13.1
$Zh$	20.6	1.9	1.9
$ZH$	0.0	0.0	0.0
$Z\bar{\nu}\nu$	626.1	109.9	111.4
$h\bar{\nu}\nu$	170.5	76.5	76.4
$H\bar{\nu}\nu$	0.0	0.0	0.0
Sum		186.5	187.7
Exact w/ISR			190.1 174.2

- ▶ Use widths to the same order as your process



# Sbottom production at the ILC

Cross sections for  $\sqrt{s} = 800$  GeV

- ▶ In contrast to the LHC: Electroweak production
- ▶ More channels contribute to  $e^+e^- \rightarrow b\bar{b}\tilde{\chi}_1^0\tilde{\chi}_1^0$ :  
 $e^+e^- \rightarrow Zh, ZH, Ah, HA, \tilde{\chi}_1^0\tilde{\chi}_2^0, \tilde{\chi}_1^0\tilde{\chi}_3^0, \tilde{\chi}_1^0\tilde{\chi}_4^0, \tilde{b}_1\tilde{b}_1^*, \tilde{b}_1\tilde{b}_2^*$  (412 diagrams)
- ▶ Irreducible SM background:  $e^+e^- \rightarrow b\bar{b}\nu_i\bar{\nu}_i$  ( $WW$  fusion,  $Zh, ZZ$ ) (47 diagrams)



Channel	$\sigma_{2\rightarrow 2}$ [fb]	$\sigma \times \text{BR}$ [fb]	$\sigma_{\text{BW}}$ [fb]
$Zh$	20.574	1.342	1.335
$ZH$	0.003	0.000	0.000
$hA$	0.002	0.001	0.000
$HA$	5.653	0.320	0.314
$\tilde{\chi}_1^0\tilde{\chi}_2^0$	69.109	13.078	13.954
$\tilde{\chi}_1^0\tilde{\chi}_3^0$	24.268	3.675	4.828
$\tilde{\chi}_1^0\tilde{\chi}_4^0$	19.337	0.061	0.938
$\tilde{b}_1\tilde{b}_1^*$	4.209	0.759	0.757
$\tilde{b}_1\tilde{b}_2^*$	0.057	0.002	0.002
Sum		19.238	22.129
Exact w/ISR			19.624
			22.552

Channel	$\sigma_{2\rightarrow 2/3}$ [fb]	$\sigma \times \text{BR}$ [fb]	$\sigma_{\text{BW}}$ [fb]
$ZZ$	202.2	12.6	13.1
$Zh$	20.6	1.9	1.9
$ZH$	0.0	0.0	0.0
$Z\bar{\nu}\nu$	626.1	109.9	111.4
$h\bar{\nu}\nu$	170.5	76.5	76.4
$H\bar{\nu}\nu$	0.0	0.0	0.0
Sum		186.5	187.7
Exact w/ISR			190.1
			174.2

- ▶ Use widths to the same order as your process


# Results

Off-shell decay  $\tilde{\chi}_3^0 \rightarrow (\tilde{b}_1)_{off} \bar{b} \rightarrow b\bar{b}\tilde{\chi}_1^0$  gives broad continuum



ISR/beamstrahlung: corrections of same order (effects all  $p_{miss}$  observables)

$b\bar{b}$  invariant mass with SM background:





Cut out the resonances

$$M_{b\bar{b}} < 150 \text{ GeV}$$

$$250 \text{ GeV} < M_{b\bar{b}} < 350 \text{ GeV}$$


# Results

Off-shell decay  $\tilde{\chi}_3^0 \rightarrow (\tilde{b}_1)_{of} f \bar{b} \rightarrow b \bar{b} \tilde{\chi}_1^0$  gives broad continuum



ISR/beamstrahlung: corrections of same order (effects all  $p_{miss}$  observables)

$b\bar{b}$  invariant mass with SM background:



Cut out the resonances

$$M_{b\bar{b}} < 150 \text{ GeV}$$


$$250 \text{ GeV} < M_{b\bar{b}} < 350 \text{ GeV}$$

# Results: Isolation of the Signal

Channel	$\sigma_{\text{BW}}$ [fb]	$\sigma_{\text{BW}}^{\text{cut}}$ [fb]
$Z\bar{\nu}\nu$	111.4	2.114
$h\bar{\nu}\nu$	76.4	0.002
$H\bar{\nu}\nu$	0.0	0.000
Sum	187.7	2.117
Exact	190.1	1.765
w/ISR	174.2	1.609

Channel	$\sigma_{\text{BW}}$ [fb]	$\sigma_{\text{BW}}^{\text{cut}}$ [fb]
$Zh$	1.335	0.009
$HA$	0.314	0.003
$\tilde{\chi}_1^0\tilde{\chi}_2^0$	13.954	0.458
$\tilde{\chi}_1^0\tilde{\chi}_3^0$	4.828	0.454
$\tilde{\chi}_1^0\tilde{\chi}_4^0$	0.938	0.937
$\tilde{b}_1\tilde{b}_1$	0.757	0.451
$\tilde{b}_1\tilde{b}_2$	0.002	0.001
Sum	22.129	<b>2.314</b>
Exact	19.624	<b>0.487</b>
w/ISR	22.552	<b>0.375</b>

$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$  decay kinematics affected





# Results: Isolation of the Signal

Channel	$\sigma_{\text{BW}}$ [fb]	$\sigma_{\text{BW}}^{\text{cut}}$ [fb]
$Z\bar{\nu}\nu$	111.4	2.114
$h\bar{\nu}\nu$	76.4	0.002
$H\bar{\nu}\nu$	0.0	0.000
Sum	187.7	2.117
Exact	190.1	1.765
w/ISR	174.2	1.609

Channel	$\sigma_{\text{BW}}$ [fb]	$\sigma_{\text{BW}}^{\text{cut}}$ [fb]
$Zh$	1.335	0.009
$HA$	0.314	0.003
$\tilde{\chi}_1^0\tilde{\chi}_2^0$	13.954	0.458
$\tilde{\chi}_1^0\tilde{\chi}_3^0$	4.828	0.454
$\tilde{\chi}_1^0\tilde{\chi}_4^0$	0.938	0.937
$\tilde{b}_1\tilde{b}_1$	0.757	0.451
$\tilde{b}_1\tilde{b}_2$	0.002	0.001
Sum	22.129	<b>2.314</b>
Exact	19.624	<b>0.487</b>
w/ISR	22.552	<b>0.375</b>

$\tilde{b}_1 \rightarrow b\tilde{\chi}_1^0$  decay kinematics affected



# Summary & Outlook

Precision predictions for SUSY pheno are important

- ▶ Higher orders: virtual corrections
- ▶ Higher orders: real corrections

Factorization in  $2 \rightarrow 2$  production and decay insufficient/wrong

Off-shell effects and interferences affect results (especially with cuts)

Use full matrix elements

Tools are available for ILC/LHC: Whizard/O'Mega

`http://james.physik.uni-freiburg.de/~reuter`

Reconsider all edge structures [LHC]: Alwall/Plehn/Rainwater/JR/Schumann

# Summary & Outlook

Precision predictions for SUSY pheno are important

- ▶ Higher orders: virtual corrections
- ▶ Higher orders: real corrections

Factorization in  $2 \rightarrow 2$  production and decay insufficient/wrong

Off-shell effects and interferences affect results (especially with cuts)

Use full matrix elements

Tools are available for ILC/LHC: Whizard/O'Mega

<http://james.physik.uni-freiburg.de/~reuter>

Reconsider all edge structures [LHC]: Alwall/Plehn/Rainwater/JR/Schumann

