

# Search for Extra Dimensions and Leptoquarks in Early LHC Data

Greg Landsberg



ILC/LHC Workshop

April 13, 2007


# Outline



- History and General Comments
- LHC Schedule
- Challenges
- Sensitivities
- Detector-Specific Challenges
- Word of Caution
- Conclusions

# History


- Drell-Yan dilepton production **historically has been a fruitful channel for discoveries** ( $J/\Psi$ ,  $Y$ ,  $Z$ )
  - So maybe the LHC case



UA1  $Z(ee)$   
candidate

# History, cont'd



- Well-understood SM cross section (calculated to NNLO level)
- Reasonably small dependence on PDF's and other systematics (except at very low x's)



$y$	0	2	4	0	2	4	0	2	4
	$M = 91.2 \text{ GeV}/c^2$			$M = 200 \text{ GeV}/c^2$			$M = 1000 \text{ GeV}/c^2$		
$x_1$	0.0065	0.0481	0.3557	0.0143	0.1056	0.7800	0.0714	0.5278	-
$x_2$	0.0065	0.0009	0.0001	0.0143	0.0019	0.0003	0.0714	0.0097	-



# History, cont'd

- Tevatron experience shows that these **analyses can be performed fast**, even with poorly understood detector:
  - Have a “**standard candle**” – the Z, which can be used for in situ measurement of the efficiencies and luminosity
  - Have been **among the first analyses shown by the Tevatron collaborations in Run II and still yield world's best sensitivity to date to a number of new phenomena**



# History, cont'd


- Beautifully clean events!



# General Comments

- **Lepton Resonances:**
  - Very clean channel with easy triggering and possibility of a cross check ( $ee/\mu\mu/\tau\tau$ )
  - In situ calibration with the Z, possibly  $J/\Psi$  and  $Y$
  - (Largely) insensitive to precise tracking alignment, jet energy scale, precise calibration
- **Leptoquarks and other gauge bosons:**
  - Once dilepton channels are understood, one can start adding other objects, e.g. jets
  - More complicated final states, but still relatively clean
  - Resonant states among the final state particles
- **Challenges:**
  - Commissioning detectors at the same time as trying to get the first results out
  - Changing beam conditions
  - Changing trigger definition and event reconstruction

# LHC Schedule, 2007



## Calibration run in 2007



Installation Hardware Commissioning	Hardware Commissioning 450GeV	Engineering Run 450GeV	Shutdown
--	-------------------------------------	------------------------------	----------

Machine checkout 450GeV	Beam commissioning 450GeV	Calibration run 450GeV
----------------------------	------------------------------	---------------------------


$k_b$	43	43	156	156
$I_b$ ( $10^{10}$ )	2	4	4	10
$\beta^*$ (m)	18	4	4	4
luminosity	$10^{28}$	$2 \cdot 10^{29}$	$7 \cdot 10^{29}$	$4 \cdot 10^{30}$
Event rate (kHz) <sup>1</sup>	0.4	8	28	180
W rate <sup>2</sup> (per 24h)	1	17	62	386
Z rate <sup>3</sup> (per 24h)	0.1	1.7	6.2	39

- |    |  |       |
|----|--|-------|
| 1. | Assuming 450GeV inelastic cross section          | 40mb  |
| 2. | Assuming 450GeV cross section $W \rightarrow lv$ | 1nb   |
| 3. | Assuming 450GeV cross section $Z \rightarrow ll$ | 100pb |



# LHC Schedule

- First 14 TeV Collisions: June 2008
- **Effective ATLAS/CMS running time/year:**  $\sim 1000$  hours  $\sim 4 \times 10^6$  s  $\sim 4 \times 10^{39}$  cm<sup>-2</sup> =  $4 \times 10^{15}$  b<sup>-1</sup> =  $4$  fb<sup>-1</sup> @  $10^{33}$  cm<sup>-2</sup>s<sup>-1</sup>
- Expected luminosity:  $\sim 1$  fb<sup>-1</sup> on 2008;  $\sim 5$  fb<sup>-1</sup> in 2009



# 'Early Phase' Assumptions




- Still large uncertainties on the delivered luminosity in 2008-2009
- Define 'early physics' as **1-10 fb<sup>-1</sup> of good quality data**
- This is the dataset we could expect by the time of the Tevatron shutdown
- This may vary from channel to channel, but gives an overall guidance on what to expect from early LHC run
- Focus on dielectron and dimuon reach; will comment on di-taus as well
- Assuming typical acceptance of 50% and 80% detection efficiency expect **~2,000 events produced for a process with 1pb cross section**
- Assuming at least ~5 events necessary for discovery, **can probe cross sections up to ~2 fb**
- Disclaimer: will use slightly more CMS than ATLAS studies, as CMS has brand new Physics TDR
  - Gustaaf will hopefully compensate this slight misbalance

# ED Models: The Stage

## ADD Model:

- Winding modes with energy spacing  $\sim 1/r$ , i.e. 1 meV – 100 MeV
- Can't resolve these modes – they appear as continuous spectrum


Gravitational coupling per mode; many modes



## TeV<sup>-1</sup> Scenario:

- Winding modes with nearly equal energy spacing  $\sim 1/r$ , i.e.  $\sim \text{TeV}$
- Can excite individual modes at colliders or look for indirect effects


$$M_i = \sqrt{M_0^2 + i^2/r^2}$$



## RS Model:

- "Particle in a box" with a special metric
- Energy eigenvalues are given by zeroes of Bessel function  $J_1$
- Light modes might be accessible at colliders

$$M_0 = 0; M_i = M_1 x_i / x_1 \approx M_1, 1.83M_1, 2.66M_1, 3.48M_1, 4.30M_1, \dots$$






# Dielectrons: Discovery Channel

- Excellent resolution 5-10%/sqrt(E, GeV) (calorimeter based) and detection efficiency
- Low background above ~1 TeV

	KK Z		$G, c = 0.01$	$G, c = 0.1$	SSM $Z'$	
$M$	4.0	6.0	1.5	3.5	1.0	5.0
$M_w$	3.5-4.5	5.0-6.7	1.47-1.52	3.30-3.65	0.92-1.07	4.18-5.81
$N_s$	50.6	1.05	18.8	7.30	72020	0.58
$N_b$	0.13	0.005	4.16	0.121	85.5	0.025
$S$	22.5	3.0	6.39	6.83	225	1.63



CMS, 30 fb<sup>-1</sup>

$Z_{KK}$  production





# Dimuons: Confirmation Channel?

- Generally worse rapidity coverage, detection efficiency





- Significantly worse momentum resolution than for electrons







- Nevertheless: generally lower instrumental background may make dimuons a discovery channel along with dielectrons

# What About Tau's?


- Less studied for the discovery reach by the LHC collaborations, but still can be accessible for early physics
- N.B. The first Tevatron Run II cross section paper was DØ  $Z(\tau\tau)$  measurement
- Very interesting reach for MSSM Higgs and other resonances; could also be tricky?



# Randall-Sundrum Graviton Reach







Coupling constant $c$	Estimator	$1 \text{ fb}^{-1}$	$10 \text{ fb}^{-1}$
0.01	$S_{cP}$	0.75	1.20
	$S_{cL}$	0.77	1.21
	$S_L$	0.78	1.23
0.02	$S_{cP}$	1.21	1.72
	$S_{cL}$	1.22	1.72
	$S_L$	1.22	1.74
0.05	$S_{cP}$	1.83	2.48
	$S_{cL}$	1.85	2.49
	$S_L$	1.85	2.51
0.1	$S_{cP}$	2.34	3.11
	$S_{cL}$	2.36	3.13
	$S_L$	2.36	3.16








# Angular Distributions?

- Not in the early running!
  - “One event – discovery; two events – cross section measurement; three events – angular distributions”







# KK Excitations of the Z Boson




- Dramatic reach even with  $\sim 1 \text{ fb}^{-1}$




# Leptoquarks

- Once II channel is understood, adding extra objects is easy, even if they are as messy as jets!
- Focus on the  $lljj$  channel
  - $evjj$  is a possibility, but no existing studies
  - $vvjj$  will take long time




$M_{LQ}$ (TeV)	$\sigma$ (fb)	Signal	Background	$S/\sqrt{B}$
1.0	4.96	98.5	2.84	58
1.2	1.33	22.0	2.43	14
1.3	0.713	12.8	1.44	11
1.5	0.223	3.62	0.376	5.9




# Leptoquarks: Reach

- Reach plot available for  $300 \text{ fb}^{-1}$ 
  - Scale significance as  $\sqrt{L}$
  - $S/\sqrt{B}$  of 50 correspond to 5 sigma at  $3 \text{ fb}^{-1}$



# Right-handed W



- $W_R$  production, with  $W_R \rightarrow l + N \rightarrow l + ljj$
- Interesting reach even with  $\sim 1 \text{ fb}^{-1}$



# Challenges: General



- Broad resonance are possible at high masses; signal start looking compositeness (or instrumental effect!) like
- Reduces the reach; requires different optimization of the search

Example: bulk  $Z_{KK}$  in RS model






# Challenges: CMS

- ECAL saturation: a single crystal saturates at  $\sim 1.7$  TeV; start seeing effect for  $>4$  TeV  $Z'$
- Correct energy at a slight resolution loss using “charge-sharing” technique
- Triggering with saturation could present another challenge!



# Challenges: ATLAS



- Electron efficiency drops fast with mass when “standard” isolation cut is used
  - Loosely confirmed by full simulation
- New set of isolation cuts is being developed to recover efficiency at high masses






# Last-Bin Effect

- Note the last-bin effects: saturation can easily cause a peak
  - Unlikely that confirmation could come from the dimuon (resolution) or ditau (ID, trigger,  $Z(\tau\tau)$  first!) channel at the time of discovery
  - Thus many cross checks will be required



# Memento Pentaquarks...

- Involved about a **dozen (!) of groups** all over the globe!
- Generated about **400 references** in the literature
- Net result: **every single claim of 2003 has been disputed** by at least one other group
- IMHO: **quite a shame for the field**, which is over 100 years old and so proud of the widely-accepted  $5\sigma$  discovery standard
- Unfortunately **follows a long trail of “miscoveries”** from split  $A_2$  to the Heidelberg-Moscow  $0\nu\beta\beta$  decay claim and DAMA story



Experiment	Reaction	State	Mode
LEPS(1)	$\gamma C_{12} \rightarrow K^+ K^- X$	$\theta^+$	$K^+ n$
LEPS(2)	$\gamma d \rightarrow K^+ K^- X$	$\theta^+$	$K^+ n$
CLAS(d)	$\gamma d \rightarrow K^+ K^- (n)p$	$\theta^+$	$K^+ n$
CLAS(p)	$\gamma p \rightarrow K^+ K^- \pi^+ (n)$	$\theta^+$	$K^+ n$
SAPHIR	$\gamma p \rightarrow K_S^0 K^+ (n)$	$\theta^+$	$K^+ n$
COSY	$pp \rightarrow \Sigma^+ K_S^0 p$	$\theta^+$	$K_S^0 p$
JINR	$p(C_3H_8) \rightarrow K_S^0 p X$	$\theta^+$	$K_S^0 p$
SVD	$pA \rightarrow K_S^0 p X$	$\theta^+$	$K_S^0 p$
DIANA	$K^+ X e \rightarrow K_S^0 p (X e)'$	$\theta^+$	$K_S^0 p$
$\nu BC$	$\nu A \rightarrow K_S^0 p X$	$\theta^+$	$K_S^0 p$
NOMAD	$\nu A \rightarrow K_S^0 p X$	$\theta^+$	$K_S^0 p$
HERMES	quasi-real photoproduction	$\theta^+$	$K_S^0 p$
ZEUS	$ep \rightarrow K_S^0 p X$	$\theta^+$	$K_S^0 p$
NA49	$pp \rightarrow \Xi \pi X$	$\Xi_3$	$\Xi \pi$
H1	$ep \rightarrow (D^+ p) X$	$\theta_c$	$D^+ p$

[Dzierba et al, hep-ex/0412077]

# Peaks Are Not Easy to Fake?




Lee Ann Womack – “Now You See Me, Now You Don't”

Better take a good look before I disappear  
Because I'm just about to be your used-to-be  
You might catch a glimpse of my taillights in the dust  
And if you notice something missin', well it's me  
'Cause I tried and you lied

...

Now you see me, now you don't  
First you do but then you won't  
Watch me vanish right before your eyes  
You might think you see me there  
In a cafe on a street somewhere  
Yeah, that might be me but I'll be gone  
Now you see me, now you don't

...



# Conclusions



- **Early LHC Physics** certainly hold **many surprises**: both pleasant and unpleasant ones
- The **machine and detector will come online** and will be commissioned **in parallel**, which adds to the challenges
- Nevertheless, **dilepton channel was historically very discovery-rich**, and if history is any guide, we should be able to discover multi-TeV resonances in these channels relatively fast
- This is a **brand new territory**, where Tevatron doesn't have any reach, so surprises may come very early
- Both ATLAS and CMS collaborations are well aware of challenges and have formed dilepton working groups looking at realistic scenarios and developing discovery strategy
  - **While a discovery can't be planned, it can and should be prepared for and accelerated!**

# The LHC Surprise ?

## •LHC vs. Tevatron:

- Nearly an order of magnitude higher c.o.m. energy (SSC would have been even better!)
- Significantly reduced relative rate of important SM backgrounds due to the pp initial state
- Consequently, surprises are possible literally on Day One of operations
- Nevertheless, how likely are they?

## •LHC event yield per 10 fb<sup>-1</sup>:

channel	recorded events per experiment for 10 fb <sup>-1</sup>
$W \rightarrow \mu\nu$	$7 \times 10^7$
$Z \rightarrow \mu\mu$	$1.1 \times 10^7$
$t\bar{t} \rightarrow \mu + X$	$0.08 \times 10^7$
QCD jets $p_T > 150$ GeV	$\sim 10^7$ (assuming 10% of trigger bandwidth)
minimum bias	$\sim 10^7$ (assuming 10% of trigger bandwidth)
$\tilde{g}\tilde{g}, m(\tilde{g}) \sim 1$ TeV	$10^3 - 10^4$

[Gianotti & Mangano, hep-ph/0504221]

## •What are the Tevatron lessons in some of these channels?

[Gianotti & Mangano, hep-ph/0504221]

