


1 SOLENOID HOMEWORK

1. Consider a 200 MeV/c particle starting on the axis with a transverse momentum of 20 MeV/c in an axial solenoidal field of 3.33 T.
 - (a) What is its motion in the lab frame and out to what transverse distance from the axis does it get.
 - (b) What is the distance along the axis before it first returns to that axis?
 - (c) What is the wavelength λ in the Larmor frame?
 - (d) what is the lattice parameter β_{\perp} for that particle

2. Consider again a $200 \text{ MeV}/c$ particle starting on the axis with a transverse momentum of $20 \text{ MeV}/c$ in an axial solenoidal field of 3.33 T . After a distance
- A) corresponding to $1/2$ a helix rotation, or
 - B) corresponding to a full helix rotation,
- the field abruptly doubles to 6.66 T .

In the two cases determine:

- (a) The shape of the motion projected onto the x, y plane
- (b) The following length in z for one helix rotation (λ_{helix})



2 COOLING HOMEWORK

1. In a linear cooling channel, assume $\beta_{\perp} = 0.4$ m, $C(mat, E) = 38 \cdot 10^{-4}$, $\beta_v = 0.85$.


a) What is the expected equilibrium transverse emittance?

2. A Guggenheim cooling channel, with emittance exchange in wedges cools all 6 dimensions. Assume $\beta_{\perp} = 0.4$ m, dispersion at the hydrogen wedge $D = 7$ cm, the length of the wedge on axis $\ell = 28.6$ cm, and the height from the axis to the apex of the wedge $h = \frac{\ell}{2 \tan(100^\circ/2)} = 12$ cm. Assume that the sum of partition functions $\sum J_i \approx 2.0$, $C(mat, E) = 38 \cdot 10^{-4}$, and assume good mixing between x and y . As before, $\beta_v = 0.85$.

a) What are the three partition functions in this case?

b) What is the expected equilibrium transverse emittance?

3. Consider a solenoid with $B_z = 3.33$ T and a muon with starting on axis with $p_t = 20$ MeV/c and $p_z = 200$ MeV/c. Imagine an ideal transverse cooling system with continuous energy loss and re-acceleration so that all transverse momenta are reduced to near zero, then the above particle will settle at half its maximum distance from the axis $r_2 = r_1/2$ and pass straight down the field lines at p_z .



- (a) What now is its motion in the Larmor frame?
 (b) If now the field B_z suddenly stops, what is the further motion of the muon?