Fast Beam-ion Instabilities

Vacuum Specifications in CLIC Main Linac

Adrian Oeftiger

Supervisor: G. Rumolo

CERN, BE/ABP–CC3

October 20, 2010

Outline

1. Introduction
2. Scan of Ion Species
   - Fourier Transformed Modes
   - Summarized Results
3. Main Components of Residual Gas
   - Results for H_2O
   - Results for CO
   - Results for CO_2
4. Summary and Perspectives

Fast beam-ion instability I

Fast beam-ion instability:

- electron bunches create ions out of residual gases by
  - scattering ionization
  - field ionization – after threshold (beam transversely small enough): currently $\sim$4km for a final energy of 1.5TeV
- electromagnetic interaction between electrons and ions
  - ions created bunch by bunch and kicked by passing bunches
  - electron bunches feel effect of ion field
- resulting in
  1. extra phase advance shift over bunch train
  2. possible excitation of unstable ion-electron coupled motion (ions do not necessarily need to be trapped)

Fast beam-ion instability II

- two-stream instability:
  - accumulation of ions around beam for long enough times $\Longrightarrow$ coupled oscillations between electrons and ions triggered (resonance, grows by its own)



source: http://sps-impedance.web.cern.ch/sps-impedance/USPAS/TwoStream.pdf

FASTION code

FASTION code produces several output files, e.g.

- bunch by bunch offsets x, x', y, y' over whole train for all lattice points



Figure: Bunch offsets – ion species: H_2O, pressure: 20nTorr

FASTION code

FASTION code produces several output files, e.g.

- bunch by bunch offsets x, x', y, y' over whole train for all lattice points



Figure: Bunch offsets – ion species: H_2O, pressure: 20nTorr

FASTION code

FASTION code produces several output files, e.g.

- bunch by bunch offsets x, x', y, y' over whole train for all lattice points
- sample ion trajectories recorded uniformly distributed



Figure: Ion trajectories – ion species: H_2O, pressure: 20nTorr

FASTION code

FASTION code produces several output files, e.g.

- bunch by bunch offsets x, x', y, y' over whole train for all lattice points
- sample ion trajectories recorded uniformly distributed



Figure: Ion trajectories – ion species: H_2O, pressure: 20nTorr

Specifications

Parameters used in the study of CLIC Main Linac:

Energy	p_0 [GeV]	9 to 1500
Norm. transv. emitt.	$\epsilon_{x,y}$ [nm]	660, 10
Bunch population	N	4×10^9
Number of bunches	N_b	312
Bunch spacing	ΔT_b [ns]	0.5
Bunch length	σ_z [ps]	0.15
Linac length	L [km]	20.5
Gas pressure	P [nTorr]	scanned
Ioniz. cross sect.	σ_{ions} [MBarn]	2
Thresh. electr. field	E_{th} [GV/m]	15

Beam Transverse Size Shrinking

What makes Main Linac special? (compared to transfer line)





- field ionization can start (small transverse beam size)
- trapping condition changing along Main Linac
- excited frequencies different along Main Linac

Approach for Scan

- scanned mass numbers: 2 (H_2), 4 (He), 5, 6, 7, 10, 12, 14, 16, 18 (H_2O), 20 (Ne), 28 (CO), 32 (O_2), 44 (CO_2), 70 ($^{35}Cl_2$), 84 (Kr), 132 (Xe)
- pressure set to 20nTorr
- analyzing bunch train motion and its fourier transformed modes

Why scan?

- evaluation of frequency dependencies on ion mass number

Fourier Transformed Modes for $A = 18$ (H_2O)



Fourier Transformed Modes for $A = 18$ (H_2O)



Fourier Transformed Modes for $A = 18$ (H_2O)



Fourier Transformed Modes for $A = 84$ (Kr)



Fourier Transformed Modes for $A = 84$ (Kr)



Fourier Transformed Modes for $A = 84$ (Kr)



Results

- upcoming unstable resonance frequencies at $A = 5$, clear beam instability from $A = 7$ on (20nTorr)
- frequency mean values and standard deviation

x-plane



y-plane



Composition of Vacuum

Three different possibilities for vacuum composition:

	Unbaked Vacuum	Baked Vacuum	BakedNEG Vacuum
H_2	40%	80%	90%
H_2O	40%	10%	4%
CO	10%	5%	3%
CO_2	10%	5%	3%

Thanks to S. Calatroni for providing the compositions

Results

- H_2 stable at least until 200nTorr, 2000nTorr is unstable (negligibly high)

Evolution of peak frequency amplitude for H_2O:





- beam instabilities in both x- and y-plane occur from 4nTorr on

Results

Evolution of peak frequency amplitude for CO:





- beam instabilities in both x- and y-plane occur from 3nTorr on

Results

Evolution of peak frequency amplitude for CO_2:





- beam instabilities occur from 2nTorr (x-plane) resp. from 3nTorr (y-plane) on

Summary

- for considerable pressures, beam instabilities occur from $A \geq 5$ on
- ion frequencies range between 0.1 and 0.5 GHz in x- and y-plane
- H_2O becomes unstable from 4nTorr on
- CO becomes unstable from 3nTorr on
- CO_2 becomes unstable from 2nTorr (x-plane) and 3nTorr (y-plane) on

Perspectives

- for combinations of gas species, thresholds could be more relaxed under constant total pressure
- simulations with several ion species (real gas compositions) can be tricky:
  - e.g. for too many H_2 macro-particles, we do not see instabilities at expected pressures compared to simulations without H_2 (due to a numerical problem)
- however, H_2 has no effect on beam $\Rightarrow$ leave it aside (fix total pressure and consider only rest parts of realistic gas composition)
- pressure threshold for energy scaled Linac, 500 GeV
- parameter scans