

Status of Higgs self-coupling analysis

Junping Tian (KEK)

Taikan Suehara (Tokyo U')

Tomohiko Tanabe (Tokyo U')

Keisuke Fujii (KEK)

Nov. 14 @ ILD Analysis Meeting

outline

- analysis strategy
- main difference from LoI analysis (flavor tagging and isolated-lepton selection)
- status of DBD analysis: ZHH @ 500 GeV
- samples request for $\nu\nu$ HH (fusion) @ 1TeV

analysis strategy and status

$$e^+ + e^- \rightarrow ZHH @ 500 \text{ GeV}$$

main backgrounds in each mode:

- ♦ **llHH:** llbb (ZZ, γ Z, bbZ), lvbbqq (tt-bar), llbbbb (ZZZ / ZZH) preliminary
- ♦ **vvHH:** bbbb (ZZ, γ Z, bbZ), τ vbbqq (tt-bar), vvbbbb (ZZZ / ZZH) ongoing
- ♦ **qqHH:** bbbb (ZZ, γ Z, bbZ), bbqqqq (tt-bar), qqbbbb (ZZZ / ZZH) preliminary


after isolated-lepton selection (or rejection) and jet-clustering, a neural-net is trained for each dominant background process (in total 9)

to make the result stable, high statistics is necessary

main improvements to the LoI analysis:



- ♦ better flavor tagging (tracking, PFA, LCFIPlus, B-baryon fixed)
- ♦ better lepton selection (muon detector, vertex constrained, bremsstrahlung and FSR recovered)

flavor tagging performance in qqHH mode



Isolated lepton selection (llHH)

$$(E_{tot} = E_{ecal} + E_{hcal})$$



electron ID

muon ID

- ◆ $E_{ecal} / E_{tot} > 0.9$ $E_{yoke} > 1.2$
- ◆ $0.5 < E_{tot} / P < 1.3$ $E_{tot} / P < 0.3$
- ◆ from primary vertex from primary vertex
- ◆ $P > 12.2 + 0.87E_{cone}$ $P > 12.6 + 4.62E_{cone}$

BS and FSR recovery adapted from ZFinder

efficiency of two isolated lepton selection
(much better for DBD)

Eff (%)	eeHH	$\mu\mu$ HH	bbbb	evbbqq	$\mu\nu$ bbqq
DBD	85.7	88.4	0.028	1.44	0.10
LoI	81.9	85.4	0.43	2.71	1.94

$$e^+ + e^- \rightarrow ZHH \rightarrow (q\bar{q})(b\bar{b})(b\bar{b}) \rightarrow q\bar{q} + 4 \text{ bjets}$$

full simulation @ 500GeV

pre-selection:

- isolated-charged-leptons rejected
- 6-jets clustering (LCFIPlus, Durham)
- combine the six jets by minimizing, and require the b tagging

$$\chi^2 = \frac{(M(b, \bar{b}) - M_H)^2}{\sigma_{H_1}^2} + \frac{(M(b, \bar{b}) - M_H)^2}{\sigma_{H_2}^2} + \frac{(M(q, \bar{q}) - M_Z)^2}{\sigma_Z^2}$$

requirement implied in the pre-selection:


- b-tagged four jets from two Higgs (b-likeness > 0.16)

final selection:


- separate to two categories: bbHH dominant and light qqHH dominant
- train the neural-nets, each event is also reconstructed as from ZZ, tt-bar, ZZZ and ZZH, and various variables are input to NN
- optimize the cuts on NN-output and tighter b-tagging

some distributions (qqHH)


b-likeness of the two jets from Z




TMVA response for classifier: MLP



TMVA response for classifier: MLP



TMVA response for classifier: MLP



preliminary

$P(e^-, e^+) = (-0.8, +0.3)$

reduction table

$E_{\text{cm}} = 500\text{GeV}, M_H = 120\text{GeV}$

$(\text{probZ1} + \text{probZ2} > 0.56)$

$\int L dt = 2\text{ab}^{-1}$

normalized	expected	MC	pre-selection	probZ1+probZ2>0.56	MissPt < 60	MLP_bbbb>0.7 4	MLP_bbqqq> 0.34	MLP_qbbbbb> 0.0	Bmax3>0.82 Bmax4>0.21
qqhh(qqbbbb)	310(129)	3.73×10^5	111(85.3)	26.7(23.0)	25.9(22.8)	20.6(18.8)	20.1(18.4)	20.0(18.3)	12.4(11.8)
bbbb	4.02×10^4	7.19×10^5	22889	2289	2253	9.04	8.06	7.94	3.32
lvbbqq	7.40×10^5	3.56×10^6	17240	357	172	8.47	6.69	6.69	0.03
qqbbbb	140	3.03×10^4	82.3	13.6	13.5	7.43	6.96	3.94	2.36
bbuddu	1.56×10^5	8.87×10^5	565	11.2	11.2	8.82	6.73	6.73	0.73
bbcdu	3.12×10^5	1.26×10^6	6109	86.8	86.4	61.6	44.6	44.1	2.41
bbcsc	1.56×10^5	1.17×10^6	12456	256	254	177	126	125	4.71
qqqqH(ZZH)	381	not available yet							
ttqq	2169	not available yet							
BG			59342	3013	2790	273	199	197	11.0

bbHH dominant:

$n_S = 12.4, n_B = 11.0 \sim 2.7\sigma$

samples of ZZH and ttqq are already available, to be added soon

preliminary

$P(e^-,e^+) = (-0.8, +0.3)$

reduction table

$E_{cm} = 500\text{GeV}, M_H = 120\text{GeV}$

$(\text{probZ1} + \text{probZ2} < 0.56)$

$\int L dt = 2\text{ab}^{-1}$

normalized	expected	MC	pre-selection	probZ1+probZ2<0.56	MissPt < 60	MLP_bbbb>0.6 3	MLP_bbqqq> 0.55	MLP_qqbbbb> 0.15	Bmax3>0.85 Bmax4>0.43
qqhh(qqbbbb)	310(129)	3.73×10^5	111(85.3)	84.3(62.3)	80.9(61.8)	66.9(53.5)	45.9(37.7)	44.5(36.6)	21.4(18.6)
bbbb	4.02×10^4	7.19×10^5	22889	20600	20282	152	62.9	53.5	25.6
lvbbqq	7.40×10^5	3.56×10^6	17240	16884	7937	536	115	105	1.36
qqbbbb	140	3.03×10^4	82.3	68.7	68.3	42.5	20.7	14.9	7.03
bbuddu	1.56×10^5	8.87×10^5	565	554	550	434	105	99.2	11.3
bbcsdu	3.12×10^5	1.26×10^6	6109	6022	5987	4559	977	917	25.4
bbcsc	1.56×10^5	1.17×10^6	12456	12200	12115	9181	1655	1556	19.2
qqqqH(ZZH)	381	not available yet							
ttqq	2169	not available yet							
BG			59342	56329	46939	14906	2936	2745	89.9

light qqHH dominant:

$n_S = 21.4, n_B = 89.9 \sim 2.0\sigma$

$$e^+ + e^- \rightarrow ZHH \rightarrow (l\bar{l})(b\bar{b})(b\bar{b}) \rightarrow l\bar{l} + 4 \text{ bjets}$$

full simulation @ 500GeV

pre-selection:

- two isolated-charged-leptons
- 4-jets clustering (LCFIPlus, Durham)
- combine the six jets by minimizing

$$\chi^2 = \frac{(M(b, \bar{b}) - M_H)^2}{\sigma_{H_1}^2} + \frac{(M(b, \bar{b}) - M_H)^2}{\sigma_{H_2}^2} + \frac{(M(l, \bar{l}) - M_Z)^2}{\sigma_Z^2}$$




requirement implied in the pre-selection:

- $|M(l\bar{l}) - M(Z)| < 40 \text{ GeV}$

final selection:

- separate to two categories: electron-type and muon-type
- train the neural-nets, each event is also reconstructed as from ZZ, tt-bar, ZZZ and ZZH, and various variables are input to NN
- optimize cuts on NN-output and b-tagging

neural-net output (1lHH)



preliminary

$P(e^-,e^+) = (-0.8, +0.3)$

reduction table

$E_{cm} = 500\text{GeV}, M_H = 120\text{GeV}$

(muon-type)

$\int L dt = 2\text{ab}^{-1}$

normalized	expected	MC	pre-selection	ltype = 13	$E_{cm12} + 4E_{cm12} < 60$ PLep1+PLep2 > 80 (Mll) > M(Z) - 27	MLP_llbb > 0.53	MLP_lvbbqq > 0.2	Bmax3 > 0.16	MLP_llbbbb > 0.52
llhh(llbbbb)	46.5(19.3)	3.88×10^5	26.5(11.0)	13.3(5.53)	13.0(5.38)	10.6(5.24)	10.4(5.23)	5.76(4.79)	4.47(3.76)
eebb	2.84×10^5	4.18×10^6	3950	0	0	0	0	0	0
$\mu\mu$ bb	4.96×10^4	1.00×10^6	1944	1943	1750	73.3	72.8	7.28	2.33
evbbqq	2.48×10^5	1.51×10^6	2437	0	0	0	0	0	0
$\mu\nu$ bbqq	2.46×10^5	1.48×10^6	239	215	95.7	65.7	33.3	2.78	0
$\tau\nu$ bbqq	2.46×10^5	1.35×10^6	156	7.76	2.62	1.82	0.80	0	0
bbqqqq	6.24×10^5	3.90×10^6	107	1.09	0	0	0	0	0
bbbb	4.02×10^4	1.02×10^6	5.84	0.08	0	0	0	0	0
llbbbb(ZZZ)	69.5	1.06×10^5	15.0	7.57	7.10	5.92	5.90	5.38	1.29
llqqh(ZZH)	157	6.30×10^4	138	69.7	68.4	54.3	54.0	12.8	2.36
BG	1.74×10^6	1.46×10^7	8992	2244	1924	201	167	28.2	5.97

muon-type:

$n_S = 4.5, n_B = 6.0 \sim 1.2\sigma$

preliminary

$P(e^-,e^+) = (-0.8, +0.3)$

reduction table

$E_{cm} = 500\text{GeV}, M_H = 120\text{GeV}$

(electron-type)

$\int Ldt = 2\text{ab}^{-1}$

normalized	expected	MC	pre-selection	ltype = 11	$E_{con12} + 4E_{con12-90}$ (M11)M(Z)1-32	MLP_l1bb>0.56	MLP_l1bbqq> 0.81	Bmax3>0.19	MLP_l1bbbb >0.5
llhh(llbbbb)	46.5(19.3)	3.88×10^5	26.5(11.0)	13.1(5.50)	12.3(5.18)	10.1(5.02)	8.60(4.57)	4.64(4.08)	3.73(3.30)
eebb	2.84×10^5	4.18×10^6	3950	3950	2762	75.4	57.8	3.88	0.81
$\mu\mu$ bb	4.96×10^4	1.00×10^6	1944	0.74	0.10	0	0	0	0
evbbqq	2.48×10^5	1.51×10^6	2437	2437	928	675	25.7	1.93	0.46
$\mu\nu$ bbqq	2.46×10^5	1.48×10^6	239	24.5	0.52	0.36	0	0	0
$\tau\nu$ bbqq	2.46×10^5	1.35×10^6	156	148	38.6	30.3	1.50	0.25	0
bbqqqq	6.24×10^5	3.90×10^6	107	106	3.93	3.93	1.04	0.16	0.16
bbbb	4.02×10^4	1.02×10^6	5.84	5.76	0.10	0	0	0	0
llbbbb(ZZZ)	69.5	1.06×10^5	15.0	7.42	6.69	5.44	4.68	4.18	0.97
llqqh(ZZH)	157	6.30×10^4	138	68.1	65.0	51.1	46.9	9.92	1.93
BG	1.74×10^6	1.46×10^7	8992	6748	3806	842	138	20.3	4.32

electron-type:

$n_S = 3.7, n_B = 4.3 \sim 1.1\sigma$

Status of DBD analysis

preliminary

$$P(e^-, e^+) = (-0.8, 0.3)$$

$$e^+ + e^- \rightarrow ZHH$$

$$M(H) = 120 \text{ GeV} \quad \int L dt = 2 \text{ ab}^{-1}$$

Energy (GeV)	Modes	signal	background	significance	
				excess (I)	measurement (II)
500	$ZHH \rightarrow (l\bar{l})(b\bar{b})(b\bar{b})$	3.7	4.3	1.5 σ	1.1 σ
		4.5	6.0	1.5 σ	1.2 σ
500	$ZHH \rightarrow (\nu\bar{\nu})(b\bar{b})(b\bar{b})$	-	-	-	-
500	$ZHH \rightarrow (q\bar{q})(b\bar{b})(b\bar{b})$	12.4	11.0	3.1 σ	2.7 σ
		21.4	89.9	2.2 σ	2.0 σ

- ◆ qqHH mode only, significance is already as same as using all modes in LoI
- ◆ llHH mode into two categories is helpful
- ◆ vvHH still ongoing
- ◆ samples @ 500 GeV are more or less ready

samples request for $\nu\nu\text{HH}$ @ 1TeV

(http://www-jlc.kek.jp/jlc/sites/default/files/users/tianjp/vvhh_1TeV_new.list)

ID	Process	Polarization (e-,e+)	X-Section / fb	Expected Events	Generated Events
I106801	nlnlhh	(-1,+1)	0.274	295.92	100000
I106802	nlnlhh	(+1,-1)	0.014	1.12	20000
I106803	n2n2hh	(-1,+1)	0.0218	23.544	100000
I106804	n2n2hh	(+1,-1)	0.014	1.12	10000
I106805	n3n3hh	(-1,+1)	0.0218	23.544	100000
I106806	n3n3hh	(+1,-1)	0.014	1.12	10000
I106807	nlnlbbbb	(-1,+1)	0.572	617.76	100000
I106808	nlnlbbbb	(+1,-1)	0.00389	0.3112	10000
I106809	n2n2bbbb	(-1,+1)	0.0147	15.876	100000
I106810	n2n2bbbb	(+1,-1)	0.00387	0.3096	10000
I106811	n3n3bbbb	(-1,+1)	0.0147	15.876	100000
I106812	n3n3bbbb	(+1,-1)	0.00387	0.3096	10000
I106813	bbnle1du	(-1,+1)	52.8	57024	300000
I106814	bbnle1du	(+1,-1)	17.3	1384	30000
I106815	bbnle1sc	(-1,+1)	52.4	56592	300000
I106816	bbnle1sc	(+1,-1)	17.26	1380.8	30000
I106817	bbn2e2du	(-1,+1)	48.8	52704	300000
I106818	bbn2e2du	(+1,-1)	17.3	1384	30000
I106819	bbn2e2sc	(-1,+1)	48.8	52704	300000
I106820	bbn2e2sc	(+1,-1)	17.28	1382.4	30000
I106821	bbn3e3du	(-1,+1)	48.8	52704	300000
I106822	bbn3e3du	(+1,-1)	17.28	1382.4	30000
I106823	bbn3e3sc	(-1,+1)	48.8	52704	300000
I106824	bbn3e3sc	(+1,-1)	17.26	1380.8	30000
I106825	bbuddu	(-1,+1)	66.1	71388	200000
I106826	bbuddu	(+1,-1)	23.6	1888	20000
I106827	bbcsdu	(-1,+1)	131.8	142344	400000
I106828	bbcsdu	(+1,-1)	47	3760	40000
I106829	bbcssc	(-1,+1)	65.9	71172	200000
I106830	bbcssc	(+1,-1)	23.5	1880	40000
I106831	nlnlccbb	(-1,+1)	0.932	1006.56	50000
I106832	nlnlccbb	(+1,-1)	0.00762	0.6096	10000
I106833	n2n2ccbb	(-1,+1)	0.0283	30.564	50000
I106834	n2n2ccbb	(+1,-1)	0.00758	0.6064	10000
I106835	n3n3ccbb	(-1,+1)	0.0283	30.564	50000
I106836	n3n3ccbb	(+1,-1)	0.00758	0.6064	10000
I106837	nlnlbbh	(-1,+1)	2.894	3125.52	50000
I106838	tth	(-1,+1)	6.23	6728.4	100000
I106839	tth	(+1,-1)	2.8	224	10000

~5M

Frank and Jan will help generate the samples (many thanks)

(now I'm preparing the generators to adapt DESY-site)

backup

result of LoI analysis

@ALCPG11

- ♦ focus on the ZHH @ 500 GeV, $M(H) = 120$ GeV.
- ♦ three decay modes of ZHH ($Z \rightarrow ll, \nu\nu, qq, H \rightarrow bb$) are investigated, based on ILD full simulation.
- ♦ neural-net methods are used to improve the background suppression.
- ♦ effects of different beam polarizations are checked.

$P(e^-, e^+) = (-0.8, 0.3)$

$$e^+ + e^- \rightarrow ZHH$$

$$M(H) = 120 \text{ GeV} \quad \int L dt = 2 \text{ ab}^{-1}$$

Energy (GeV)	Modes	signal	background	significance	
				excess (I)	measurement (II)
500	$ZHH \rightarrow (l\bar{l})(b\bar{b})(b\bar{b})$	6.4	6.7	2.1σ	1.7σ
500	$ZHH \rightarrow (\nu\bar{\nu})(b\bar{b})(b\bar{b})$	5.2	7.0	1.7σ	1.4σ
500	$ZHH \rightarrow (q\bar{q})(b\bar{b})(b\bar{b})$	8.5	11.7	2.2σ	1.9σ
		16.6	129	1.4σ	1.3σ

$$\sigma_{ZHH} = 0.22 \pm 0.07 \text{ fb}$$

precision of cross section: **32%**
 precision of Higgs self-coupling: **57%**