

Physics at ILC

Keisuke Fujii (KEK)

Electroweak Symmetry Breaking

Mystery of something in the vacuum

- Success of the SM = success of gauge principle

W_T and Z_T = gauge fields of the EW gauge symmetry

- Gauge symmetry forbids explicit mass terms for W and Z

→ it must be broken by something condensed in the vacuum: $\langle 0 | I_3, Y | 0 \rangle \neq 0$ $\langle 0 | I_3 + Y | 0 \rangle = 0$

- This “something” supplies 3 longitudinal modes of W and Z:

$$W_L^+, W_L^-, Z_L \longleftarrow \chi^+, \chi^-, \chi_3 : \text{Goldstone modes}$$

- Left- (f_L) and right-handed (f_R) matter fermions carry different EW charges.

Their explicit mass terms also forbidden by the EW gauge symmetry

They must be generated through their Yukawa interactions with some weak-charged vacuum


- In the SM, the same “something” mixes f_L and f_R → generating masses and inducing flavor-mixings
- In order to form the Yukawa interaction terms, we need a complex doublet scalar field, which has four real components. The SM identifies three of them with the Goldstone modes.
- We need one more to form a complex doublet, which is the physical Higgs boson.
- This SM symmetry breaking sector is the simplest and the most economical, but there is no reason for it. The symmetry breaking sector might be more complex.
- We don't know whether the “something” is elementary or composite.
- We knew it's there in the vacuum with a vev of 246 GeV. But other than that we didn't know almost anything about the “something” until July 4, 2012.

Since the July 4th, the world has changed!

The discovery of the ~ 125 GeV boson at LHC could be called a quantum jump.



- $X(125) \rightarrow \gamma\gamma$ means X is a neutral boson and $J \neq 1$ (Landau-Yang theorem). Recent LHC results prefer $J^P=0^+$.
- $X(125) \rightarrow ZZ^*, WW^* \Rightarrow \exists XVV$ couplings: ($V=W/Z$: gauge bosons)
- There is no gauge coupling like XVV , only $XXVV$ or XXV
 - $\Rightarrow XVV$ probably from $XXVV$ with one X replaced by $\langle X \rangle \neq 0$, namely $\langle X \rangle XVV$
 - \Rightarrow There must be $\langle X \rangle \langle X \rangle VV$, a mass term for V .
 - \Rightarrow X is at least part of the origin of the masses of $V=W/Z$.
 - \Rightarrow This is a great step forward but we need to know whether $\langle X \rangle$ saturates the SM vev = 246 GeV.

- $X \rightarrow ZZ^*$ means, X can be produced via $e^+e^- \rightarrow Z^* \rightarrow ZX$.




- By the same token, $X \rightarrow WW^*$ means, X can be produced via W fusion: $e^+e^- \rightarrow \nu\nu X$.

- So we now know that the major Higgs production mechanisms in e^+e^- collisions are indeed available at the ILC \Rightarrow No lose theorem for the ILC.
- ~ 125 GeV is the best place for the ILC, where variety of decay modes are accessible.
- We need to check this ~ 125 GeV boson in detail to see if it has indeed all the required properties of the something in the vacuum.



What Properties to Measure?

The Key is the Mass-Coupling Relation



- Properties to measure are
 - mass, width, J^{PC}
 - Gauge quantum numbers (multiplet structure)
 - Yukawa couplings
 - Self-coupling
- The key is to measure the mass-coupling relation

If the 125GeV boson is the one to give masses to all the SM particles, coupling should be proportional to mass.

Any deviation from the straight line signals BSM!

The Higgs is a window to BSM physics!

Our Mission = Bottom-up Model-Independent Reconstruction of the EWSB Sector through Precision Higgs Measurements

- **Multiplet structure :**
 - Additional singlet?
 - Additional doublet?
 - Additional triplet?
- **Underlying dynamics :**
 - Weakly interacting or strongly interacting? = elementary or composite ?
- Relations to other questions of HEP :
 - DM
 - EW baryogenesis
 - neutrino mass
 - inflation?

There are many possibilities!

Different models predict different deviation patterns --> **Fingerprinting!**

Model	μ	τ	b	c	t	g_V
Singlet mixing	↓	↓	↓	↓	↓	↓
2HDM-I	↓	↓	↓	↓	↓	↓
2HDM-II (SUSY)	↑	↑	↑	↓	↓	↓
2HDM-X (Lepton-specific)	↑	↑	↓	↓	↓	↓
2HDM-Y (Flipped)	↓	↓	↑	↓	↓	↓

Mixing with singlet

$$\frac{g_{hVV}}{g_{SMVV}} = \frac{g_{hff}}{g_{SMff}} = \cos\theta \simeq 1 - \frac{\delta^2}{2}$$

Composite Higgs

$$\frac{g_{hVV}}{g_{SMVV}} \simeq 1 - 3\%(1 \text{ TeV}/f)^2$$


$$\frac{g_{hff}}{g_{SMff}} \simeq \begin{cases} 1 - 3\%(1 \text{ TeV}/f)^2 & \text{(MCHM4)} \\ 1 - 9\%(1 \text{ TeV}/f)^2 & \text{(MCHM5)} \end{cases}$$

SUSY


$$\frac{g_{hbb}}{g_{SMbb}} = \frac{g_{h\tau\tau}}{g_{SM\tau\tau}} \simeq 1 + 1.7\% \left(\frac{1 \text{ TeV}}{m_A} \right)^2$$

Expected deviations are small --> **Precision!**

ILD



SiD



For the precision we need a 500GeV LC and high precision detectors

Why 250-500 GeV?

Three well known thresholds

ZH @ 250 GeV ($\sim M_Z + M_H + 20 \text{ GeV}$) :

- Higgs mass, width, J^{PC}
- Gauge quantum numbers
- Absolute measurement of HZZ coupling (recoil mass) \rightarrow couplings to H (other than top)
- $\text{BR}(h \rightarrow VV, qq, ll, \text{invisible})$: $V=W/Z$ (direct), g, γ (loop)

ttbar @ 340-350 GeV ($\sim 2m_t$) : ZH meas. Is also possible

- Threshold scan \rightarrow **theoretically clean m_t measurement**: $\Delta m_t(\overline{MS}) \simeq 100 \text{ MeV}$
 \rightarrow test stability of the SM vacuum
 \rightarrow **indirect meas. of top Yukawa coupling**
- A_{FB} , Top momentum measurements
- Form factor measurements

$\gamma\gamma \rightarrow \text{HH}$ @ 350 GeV possibility

vvH @ 350 - 500 GeV :

- HWW coupling \rightarrow **total width** \rightarrow absolute normalization of Higgs couplings






ZHH @ 500 GeV ($\sim M_Z + 2M_H + 170 \text{ GeV}$) :

- Prod. cross section attains its maximum at around 500 GeV \rightarrow **Higgs self-coupling**

ttbarH @ 500 GeV ($\sim 2m_t + M_H + 30 \text{ GeV}$) :

- Prod. cross section becomes maximum at around 800 GeV.
- QCD threshold correction enhances the cross section \rightarrow **top Yukawa** measurable at 500 GeV concurrently with the self-coupling



We can complete the mass-coupling plot at $\sim 500 \text{ GeV}$!



Main Production Processes

Single Higgs Production

Production cross section



ZH dominates at 250 GeV
(~80k ev: 250 fb⁻¹)

vvH takes over at 500 GeV
(~125k ev: 500 fb⁻¹)



Possible to rediscover the Higgs in one day!

ILC 250

Recoil Mass Measurement

The flagship measurement of ILC 250

Recoil Mass



$$M_X^2 = (p_{CM} - (p_{\mu^+} + p_{\mu^-}))^2$$

Invisible decay detectable!

$$250 \text{ fb}^{-1} @ 250 \text{ GeV} \quad m_H = 125 \text{ GeV}$$

$$\Delta\sigma_H / \sigma_H = 2.6\%$$

$$\Delta m_H = 30 \text{ MeV}$$

$$BR(\text{invisible}) < 1\% @ 95\% \text{ C.L.}$$

scaled from $m_H = 120 \text{ GeV}$


Model-independent absolute measurement of σ_{ZH} (the HZZ coupling)

Branching Ratio Measurements

for $b, c, g, \tau, WW^*, \dots$

DBD Physics Chap.

$250 \text{ fb}^{-1} @ 250 \text{ GeV}$
 $m_H = 125 \text{ GeV}$
 scaled from $m_H = 120 \text{ GeV}$



	@250GeV
process	ZH
Int. Lumi. [fb^{-1}]	250
$\Delta\sigma/\sigma$	2.6%
decay mode	$\Delta\sigma\text{Br}/\sigma\text{Br}$
$H \rightarrow b\bar{b}$	1.2%
$H \rightarrow c\bar{c}$	8.3%
$H \rightarrow gg$	7.0%
$H \rightarrow WW^*$	6.4%
$H \rightarrow \tau\tau$	4.2%
$H \rightarrow ZZ^*$	19%
$H \rightarrow \gamma\gamma$	29-38%

What we measure is not BR itself but σBR .

To extract BR from σBR , we need σ from the recoil mass measurement.

--> $\Delta\sigma/\sigma=2.6\%$ eventually limits the BR measurements.

--> If we want to improve this situation, we need more data at 250GeV.

We need to seriously think about luminosity upgrade scenario.

preliminarily

Total Width and Coupling Extraction

One of the major advantages of the LC

To extract couplings from BRs, we need the total width:

$$g_{HAA}^2 \propto \Gamma(H \rightarrow AA) = \Gamma_H \cdot BR(H \rightarrow AA)$$

To determine the total width, we need at least one partial width and corresponding BR:

$$\Gamma_H = \Gamma(H \rightarrow AA) / BR(H \rightarrow AA)$$

In principle, we can use $A=Z$, or W for which we can measure both the BRs and the couplings:

$BR(H \rightarrow ZZ^*)$

$\Gamma(H \rightarrow ZZ^*)$

BR=O(1%): precision limited by low stat.
for $H \rightarrow ZZ^*$ events

$250 \text{ fb}^{-1} @ 250 \text{ GeV}$
 $\Delta\Gamma_H / \Gamma_H \simeq 20\%$

$\Gamma(H \rightarrow WW^*)$

$BR(H \rightarrow WW^*)$

More advantageous but not easy at low E

$250 \text{ fb}^{-1} @ 250 \text{ GeV}$
 $\Delta\Gamma_H / \Gamma_H \simeq 11\%$

C.F.Durig, Helmholtz Alliance
6th WS, Dec. 2012

ILC 500


Width and BR Measurements at 500 GeV

Addition of 500GeV data to 250GeV data

E_{cm} [GeV]	independent measurements	relative error
250	σ_{ZH}	2.6%
	$\sigma_{ZH} \cdot Br(H \rightarrow b\bar{b})$	1.2%
	$\sigma_{ZH} \cdot Br(H \rightarrow c\bar{c})$	8.3%
	$\sigma_{ZH} \cdot Br(H \rightarrow gg)$	7.0%
	$\sigma_{ZH} \cdot Br(H \rightarrow WW^*)$	6.4%
	$\sigma_{ZH} \cdot Br(H \rightarrow \tau^+\tau^-)$	4.2%
	$\sigma_{\nu\bar{\nu}H} \cdot Br(H \rightarrow b\bar{b})$	10.5%
500	$\sigma_{ZH} \cdot Br(H \rightarrow b\bar{b})$	1.8%
	$\sigma_{ZH} \cdot Br(H \rightarrow c\bar{c})$	13%
	$\sigma_{ZH} \cdot Br(H \rightarrow gg)$	11%
	$\sigma_{ZH} \cdot Br(H \rightarrow WW^*)$	9.2%
	$\sigma_{ZH} \cdot Br(H \rightarrow \tau^+\tau^-)$	5.4%
	$\sigma_{\nu\bar{\nu}H} \cdot Br(H \rightarrow b\bar{b})$	0.66%
	$\sigma_{\nu\bar{\nu}H} \cdot Br(H \rightarrow c\bar{c})$	6.2%
	$\sigma_{\nu\bar{\nu}H} \cdot Br(H \rightarrow gg)$	4.1%
	$\sigma_{\nu\bar{\nu}H} \cdot Br(H \rightarrow WW^*)$	2.4%

250 fb⁻¹ @250 GeV
 +500 fb⁻¹ @500 GeV
 $m_H = 125$ GeV

ILD DBD Full Simulation Study



comes in as a powerful tool!




$$\Delta\Gamma_H/\Gamma_H \simeq 6\%$$

Mode	$\Delta\text{BR}/\text{BR}$
bb	2.8 (2.9)%
cc	5.3 (8.7)%
gg	4.3 (7.5)%
WW*	3.6 (6.9)%
$\tau\tau$	4.1 (4.9)%

The numbers in the parentheses are as of 250 fb⁻¹ @250 GeV

Top Yukawa Coupling

The largest among matter fermions, but not yet directly observed



A factor of 2 enhancement from QCD bound-state effects

Cross section maximum at around $E_{cm} = 800 \text{ GeV}$

Philipp Roloff, LCWS12

Tony Price, LCWS12

DBD Full Simulation

$$1 \text{ ab}^{-1} @ 500 \text{ GeV} \quad m_H = 125 \text{ GeV}$$

$$\Delta g_Y(t) / g_Y(t) = 9.9\%$$

Tony Price, LCWS12


scaled from $m_H = 120 \text{ GeV}$

Notice $\sigma(500+20 \text{ GeV}) / \sigma(500 \text{ GeV}) \sim 2$
Moving up a little bit helps significantly!





Higgs Self-coupling

What force makes the Higgs condense in the vacuum?

We need to **measure the Higgs self-coupling**




= We need to **measure the shape of the Higgs potential**






The measurement is very difficult even at ILC.

The Problem : BG diagrams dilute self-coupling contribution


Signal diagram






Irreducible BG diagrams



Signal diagram





Irreducible BG diagrams



$$\sigma = \lambda^2 S + \lambda I + B$$

$$\frac{\Delta\lambda}{\lambda} = F \cdot \frac{\Delta\sigma}{\sigma}$$

F=0.5 if no BG diagrams




Junping Tian LC-REP-2013-003

Higgs self-coupling @ 500 GeV (combined)

$P(e^-, e^+) = (-0.8, +0.3)$
 $e^+ + e^- \rightarrow ZHH$
 $M(H) = 120 \text{ GeV}$
 $\int L dt = 2 \text{ ab}^{-1}$

Energy (GeV)	Modes	signal	background (tt, ZZ, ZZH/ ZZZ)	significance	
				excess (I)	measurement (II)
500	$ZHH \rightarrow (l\bar{l})(b\bar{b})(b\bar{b})$	3.7	4.3	1.5 σ	1.1 σ
		4.5	6.0	1.5 σ	1.2 σ
500	$ZHH \rightarrow (\nu\bar{\nu})(b\bar{b})(b\bar{b})$	8.5	7.9	2.5 σ	2.1 σ
500	$ZHH \rightarrow (q\bar{q})(b\bar{b})(b\bar{b})$	13.6	30.7	2.2 σ	2.0 σ
		18.8	90.6	1.9 σ	1.8 σ

Hypothesis test





$\sigma_{ZHH} = 0.22 \pm 0.06 \text{ fb}$

$\frac{\delta\sigma}{\sigma} = 27\%$

$\frac{\delta\lambda}{\lambda} = 44\%$

(cf. 80% for qqbbbb at the LoI time)

χ^2 as a function of cross section



2013

ILC 1000

Higgs Physics at Higher Energy

Self-coupling with WBF, top Yukawa at xsection max., other higgses, ...

vvH @ at >1TeV : $> 1 \text{ ab}^{-1}$ (pol e^+, e^-)=(+0.2,-0.8)





- allows us to measure rare decays such as $H \rightarrow \mu^+ \mu^-$, ...
- further improvements of coupling measurements

vvHH @ 1TeV or higher : 2 ab^{-1} (pol e^+, e^-)=(+0.2,-0.8)

- cross section increases with E_{cm} , which compensates the dominance of the background diagrams at higher energies, thereby giving a better precision for the self-coupling.
- If possible, we want to see the running of the self-coupling (very very challenging).

t \bar{t} H @ 1TeV : 1 ab^{-1}

- Prod. cross section becomes maximum at around 800GeV.
- CP mixing of Higgs can be unambiguously studied.



Obvious but most important advantage of higher energies in terms of Higgs physics is, however, its **higher mass reach to other Higgs bosons** expected in extended Higgs sectors and **higher sensitivity to $W_L W_L$ scattering** to decide whether the Higgs sector is strongly interacting or not.

In any case we can improve the mass-coupling plot by including the data at 1TeV!

Independent Higgs Measurements at ILC

Canonical ILC program


250 GeV: 250 fb⁻¹
 500 GeV: 500 fb⁻¹
 1 TeV: 1000 fb⁻¹

(M_H = 125 GeV)

Ecm	250 GeV		500 GeV		1 TeV
luminosity [fb ⁻¹]	250		500		1000
polarization (e ⁻ , e ⁺)	(-0.8, +0.3)		(-0.8, +0.3)		(-0.8, +0.2)
process	ZH	vvH(fusion)	ZH	vvH(fusion)	vvH(fusion)
cross section	2.6%	-	-		
	σ·Br	σ·Br	σ·Br	σ·Br	σ·Br
H→bb	1.2%	10.5%	1.8%	0.66%	0.32%
H→cc	8.3%		13%	6.2%	3.1%
H→gg	7.0%		11%	4.1%	2.3%
H→WW*	6.4%		9.2%	2.4%	1.6%
H→ττ	4.2%		5.4%	9.0%	3.1%
H→ZZ*	19%		25%	8.2%	4.1%
H→γγ	29-38%		29-38%	20-26%	7-10%
H→μμ	-	-	-	-	31%

Top Yukawa Coupling at 1TeV


The largest among matter fermions, but not yet observed



Cross section maximum at around $E_{cm} = 800\text{GeV}$

Tony Price & Tomohiko Tanabe: ILD DBD Study
Philipp Roloff & Jan Strube: SiD DBD Study

DBD Full Simulation



Similar significance in both modes

8-jet mode: 7.9σ (TMVA)

L+6-jet mode: 8.4σ (TMVA)

Tony Price & Tomohiko Tanabe: ILD DBD Study

$$1 \text{ ab}^{-1} @ 500 \text{ GeV} \quad m_H = 125 \text{ GeV}$$

$$\Delta g_Y(t) / g_Y(t) = 9.9\%$$

Tony Price, LCWS12

scaled from $m_H=120 \text{ GeV}$



$$1 \text{ ab}^{-1} @ 1 \text{ TeV} \quad m_H = 125 \text{ GeV}$$

$$\Delta g_Y(t) / g_Y(t) = 3.0\%$$

ILD / SiD DBD Studies

Higgs self-coupling @ 1 TeV

$P(e^-,e^+) = (-0.8, +0.2)$
 $e^+ + e^- \rightarrow \nu\bar{\nu}HH$
 $M(H) = 120\text{GeV}$
 $\int Ldt = 2\text{ab}^{-1}$


	Expected	After Cut
$\nu\nu hh$ (WW F)	272	35.7
$\nu\nu hh$ (ZHH)	74.0	3.88
BG (tt/ $\nu\nu$ ZH)	7.86×10^5	33.7
significance	0.30	4.29

- better sensitive factor
- benefit more from beam polarization
- BG tt x-section smaller
- more boosted b-jets

$\frac{\Delta\sigma}{\sigma} \approx 23\%$
 $\frac{\Delta\lambda}{\lambda} \approx 18\%$

Double Higgs excess significance: $> 7\sigma$

Higgs self-coupling significance: $> 5\sigma$



HHH Prospects

Scenario A: $HH \rightarrow bbbb$, full simulation done

Scenario B: by adding $HH \rightarrow bbWW^*$, full simulation ongoing,
expect $\sim 20\%$ relative improvement

Scenario C: color-singlet clustering, future improvement,
expected $\sim 20\%$ relative improvement (conservative)

HHH	500 GeV			500 GeV + 1 TeV		
Scenario	A	B	C	A	B	C
Canonical	104%	83%	66%	26%	21%	17%
LumiUP	58%	46%	37%	16%	13%	10%

ILC 250+500+1000

Model-independent Global Fit for Couplings

Canonical ILC program

($M_H = 125 \text{ GeV}$)

250 GeV: 250 fb⁻¹

500 GeV: 500 fb⁻¹

1 TeV: 1000 fb⁻¹

$P(e^-,e^+) = (-0.8, +0.3) @ 250, 500 \text{ GeV}$


$P(e^-,e^+) = (-0.8, +0.2) @ 1 \text{ TeV}$

coupling	250 GeV	250 GeV + 500 GeV		250 GeV + 500 GeV + 1 TeV	
HZZ	1.3%	1.3%		1.3%	
HWW	4.8%	1.4%		1.4%	
Hbb	5.3%	1.8%		1.5%	
Hcc	6.8%	2.9%		2.0%	
Hgg	6.4%	2.4%		1.8%	
H $\tau\tau$	5.7%	2.4%		1.9%	
H $\gamma\gamma$	18%	8.4%		4.1%	
H $\mu\mu$	-	-		16%	
Γ_0	11%	5.9%		5.6%	
Htt	-	14%		3.2%	
HHH	-	104%	66%(*)	26%	17%(*)

) With H \rightarrow WW (preliminary) and expected improvements in jet clustering

Mass Coupling Relation

After Canonical ILC Program



LHC + ILC

Higgs couplings


LHC including Moriond/Aspen data [SFitter: Klute, Lafaye, TP, Rauch, Zerwas]

- focus SM-like [secondary solutions possible]
- six couplings and ratios from data
 - g_b from width
 - g_g vs g_t not yet possible
- [similar: Ellis etal, Djouadi etal, Strumia etal, Grojean etal]
- poor man's analyses: $\Delta_H, \Delta_V, \Delta_f$
- Tevatron $H \rightarrow b\bar{b}$ with little impact

Future dinosaurs

- LHC extrapolations unclear [SFitter version 0]
- theory extrapolations tricky [SFitter version 0]
- ILC case obvious [500 GeV for now]
- interplay in loop-induced couplings
- $t\bar{t}H$ important at LHC and ILC


LHC+LC500 Synergy!



Expected Precision and Deviation

Combined Fit with LHC data

$g(hAA)/g(hAA)|_{SM}^{-1}$ LHC/ILC1/ILC/ILCTeV



Assumed Luminosities

LHC = LHC14TeV: 300fb⁻¹

HLC = ILC250: 250fb⁻¹

ILC = ILC500: 500fb⁻¹

ILCTeV = ILC1000: 1000fb⁻¹

Maximum deviation when nothing but the 125 GeV object would be found at LHC

	ΔhVV	$\Delta h\bar{t}t$	$\Delta h\bar{b}b$
Mixed-in Singlet	6%	6%	6%
Composite Higgs	8%	tens of %	tens of %
Minimal Supersymmetry	< 1%	3%	10% ^a , 100% ^b
LHC 14 TeV, 3 ab ⁻¹	8%	10%	15%

R.S.Gupta, H.Rzehak, J.D.Wells

arXiv: 1206.3560v1

Mixing with singlet

$$\frac{g_{hVV}}{g_{SMVV}} = \frac{g_{hff}}{g_{SMff}} = \cos\theta \simeq 1 - \frac{\delta^2}{2}$$

Composite Higgs

$$\frac{g_{hVV}}{g_{SMVV}} \simeq 1 - 3\%(1 \text{ TeV}/f)^2$$

$$\frac{g_{hff}}{g_{SMff}} \simeq \begin{cases} 1 - 3\%(1 \text{ TeV}/f)^2 & \text{(MCHM4)} \\ 1 - 9\%(1 \text{ TeV}/f)^2 & \text{(MCHM5)} \end{cases}$$

SUSY

$$\frac{g_{hbb}}{g_{SMbb}} = \frac{g_{h\tau\tau}}{g_{SM\tau\tau}} \simeq 1 + 1.7\% \left(\frac{1 \text{ TeV}}{m_A} \right)^2$$

Figure 2: Comparison of the capabilities of LHC and ILC for model-independent measurements of Higgs boson couplings. The plot shows (from left to right in each set of error bars) 1 σ confidence intervals for LHC at 14 TeV with 300 fb⁻¹, for ILC at 250 GeV and 250 fb⁻¹ ('ILC1'), for the full ILC program up to 500 GeV with 500 fb⁻¹ ('ILC'), and for a program with 1000 fb⁻¹ for an upgraded ILC at 1 TeV ('ILCTeV'). More details of the presentation are given in the caption of Fig. 1. The marked horizontal band represents a 5% deviation from the Standard Model prediction for the coupling.

Fingerprinting is possible or we will get lower bounds on the BSM scale!

HL-ILC ?

High Luminosity ILC

The current ILC design is rather conservative!



ILC Luminosity Upgrade

- Concept: increase n_b from 1312 → 2625
 - Reduce linac bunch spacing 554 ns → 336 ns
 - Increase pulse current 5.8 → 8.8 mA
 - Increase number of klystrons by ~50%
- Doubles beam power → $\times 2$ L ($3.6 \times 10^{34} \text{cm}^{-2}\text{s}^{-1}$)
- Damping ring:
 - Electron ring doubles current (389mA → 778mA)
 - Positron ring: possible 2nd (stacked) ring (e-cloud limit)
- AC power: 161 MW → 204 MW (est.)
 - AC power increased by $\times 1.5$
 - shorter fill time and longer beam pulse results in higher RF-beam efficiency (44% → 61%)

14 March, 2013

Marc Ross, SLAC


11



ILC at low/high Ecm

- Low E_{cm} operation of upgraded ILC:
 - $L_{250} \sim 3e34$; Wall plug 200 MW
 - Higgs Factory Option
- High $E_{\text{cm}} \sim 1.5$ TeV
 - $L_{1500} \sim 6e34$; Wall plug 340 MW

Assumes 2x improved efficiency; 2450 bunches



14 March, 2013

Marc Ross, SLAC

13

**x4 upgrade
@250GeV**

Snowmass e^+e^- Collider Luminosity (fb^{-1})

based on 3×10^7 s running time for ILC & LEP3/TLEP

E_{cm} (GeV)	ILC	ILC Lum Upgrade	LEP3	TLEP
250	250	900	300	1500
350	300	950		200
500	500	1100		
1000	1500	1500		

Independent Higgs Measurements

Hypothetical HL-ILC

($M_H = 125 \text{ GeV}$)

250 GeV: 250 fb⁻¹
500 GeV: 500 fb⁻¹
1 TeV: 1000 fb⁻¹



250 GeV: 1150 fb⁻¹
500 GeV: 1600 fb⁻¹
1 TeV: 2500 fb⁻¹

Ecm	250 GeV		500 GeV		1 TeV
luminosity · fb	250		500		1000
polarization (e-,e+)	(-0.8, +0.3)		(-0.8, +0.3)		(-0.8, +0.2)
process	ZH	vvH(fusion)	ZH	vvH(fusion)	vvH(fusion)
cross section	1.2%	-	-		
	$\sigma \cdot \text{Br}$	$\sigma \cdot \text{Br}$	$\sigma \cdot \text{Br}$	$\sigma \cdot \text{Br}$	$\sigma \cdot \text{Br}$
H-->bb	0.56%	4.9%	1.0%	0.37%	0.20%
H-->cc	3.9%		7.2%	3.5%	2.0%
H-->gg	3.3%		6.0%	2.3%	1.4%
H-->WW*	3.0%		5.1%	1.3%	1.0%
H--> $\tau\tau$	2.0%		3.0%	5.0%	2.0%
H-->ZZ*	8.8%		14%	4.6%	2.6%
H--> $\gamma\gamma$	16%		19%	13%	5.4%
H--> $\mu\mu$	-	-	-	-	20%

Coupling Measurements

Hypothetical HL-ILC

($M_H = 125 \text{ GeV}$)

250 GeV: 1150 fb⁻¹
 500 GeV: 1600 fb⁻¹
 1 TeV: 2500 fb⁻¹

$P(e^-,e^+) = (-0.8, +0.3) @ 250, 500 \text{ GeV}$

$P(e^-,e^+) = (-0.8, +0.2) @ 1 \text{ TeV}$

coupling	250 GeV	250 GeV + 500 GeV	250 GeV + 500 GeV + 1 TeV
HZZ	0.61%	0.61%	0.61%
HWW	2.3%	0.67%	0.65%
Hbb	2.5%	0.90%	0.74%
Hcc	3.2%	1.5%	1.1%
Hgg	3.0%	1.3%	0.93%
H $\tau\tau$	2.7%	1.2%	0.99%
H $\gamma\gamma$	8.2%	4.5%	2.4%
H $\mu\mu$	-	-	10%
Γ_0	5.4%	2.8%	2.7%
Htt	-	7.8%	2.0%

HHH	-	58%	37%(*)	16%	10%(*)
-----	---	-----	--------	-----	--------

) With H->WW (preliminary) and expected improvements in jet clustering

Conclusions

- The primary goal for the next decades is to uncover the secret of the EW symmetry breaking. This will open up **a window to BSM** and **set the energy scale for the E-frontier machine that will follow LHC and ILC 500.**
- **Probably LHC will hit systematic limits at O(5-10%) for most of $\sigma \times \text{Br}$ measurements, being not enough to see the BSM effects if we are in the decoupling regime.**
To achieve the primary goal we hence need a 500 GeV LC for self-contained precision Higgs studies **to complete the mass-coupling plot**
 - starting from $e^+e^- \rightarrow ZH$ at $E_{\text{cm}} = 250\text{GeV}$,
 - then $t\bar{t}$ at around 350GeV,
 - and then ZHH and $t\bar{t}H$ at 500GeV.
- **The ILC to cover up to 500 GeV is an ideal machine to carry out this mission** (regardless of BSM scenarios) and we can do this with staging starting from 250GeV. We may need more data depending on the size of the deviation. **Lumi-upgrade possibility should be always kept in our scope.**
- If we are lucky, some extra Higgs boson or some other new particle might be within reach already at ILC 500. Let's hope that the upgraded LHC will make another great discovery in the next run.
- If not, we will most probably need **the energy scale information from the precision Higgs studies.** Guided by the energy scale information, we will go hunt direct BSM signals with a new machine, if necessary.

Last but Not Least

- In this talk I have been focusing on the case where $X(125\text{GeV})$ alone would be the probe for BSM physics, but there is a good chance for the higher energy run of LHC to bring us more.
- It is also very important to stress that ILC, too, is an energy frontier machine. It will access the energy region never explored with any lepton collider. There can be a zoo of new uncolored particles or new phenomena that are difficult to find at LHC but can be discovered and studied in detail at ILC.
- For instance
 - Natural SUSY : naturalness prefers μ not far above 100GeV
 - > light chargino/neutralinos will be higgsino-dominant and nearly degenerate
 - > typically Δm of a few GeV or less (very difficult for LHC)
 - > Δm as small as 50MeV possible with ISR tagging at ILC
 - > If $\Delta m=800\text{MeV}$ --> possible to measure m to 1.5GeV and Δm to 20MeV
 - > ILC will also be a Higgsino factory!
 - Possible anomalies in precision studies of properties of top, W/Z, and two-fermion processes

Higgsinos in Natural SUSY ($\Delta M < \text{a few GeV}$)


Hale Sert
ECFA LCWS 2013, DESY

$$e^+e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \gamma$$

$$e^+e^- \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_1^0 \gamma$$

ISR Tagging

Ref: C.-H. Chen et al. hep-ph:9512230



Only very soft particles in the final states → Require a hard ISR to kill huge two-photon BG!



Hale Sert

- Light Higgsinos are well motivated by naturalness
- It is a challenging scenario for LHC
- Separation of Higgsinos at the reconstructed level is possible at the ILC
- Assumed

- ▶ $\sqrt{s} = 500 \text{ GeV}$
- ▶ $\int \mathcal{L} dt = 500 \text{ fb}^{-1}$ with $P(e^+, e^-) = (+30\%, -80\%)$ and $P(e^+, e^-) = (-30\%, +80\%)$ each



- Statistical uncertainties for $P(e^+, e^-) = (+30\%, -80\%)$

$m_h = 124 \text{ GeV}$

- ▶ $\delta(\sigma \times BR) \approx 3\%$ $\delta M_{\tilde{\chi}_1^\pm}(M_{\tilde{\chi}_2^0}) \approx 2.1(3.7) \text{ GeV}$ $\delta \Delta M(\tilde{\chi}_1^\pm, \tilde{\chi}_1^0) = 70 \text{ MeV}$


$m_h = 127 \text{ GeV}$

- ▶ $\delta(\sigma \times BR) \approx 1.5\%$ $\delta M_{\tilde{\chi}_1^\pm}(M_{\tilde{\chi}_2^0}) \approx 1.5(1.6) \text{ GeV}$ $\delta \Delta M(\tilde{\chi}_1^\pm, \tilde{\chi}_1^0) = 20 \text{ MeV}$



Top Quark

Anomalous Couplings in Open Top Production at 500 GeV



LAL 11-222

Figure 34: Predictions of various groups [40,42–44] on deviations from Standard Model couplings of the t quark within Randall-Sundrum Models. The cartoon is taken from [47].

coupling	LHC, 300 fb ⁻¹	e ⁺ e ⁻ [23]	coupling	LHC, 300 fb ⁻¹	e ⁺ e ⁻ [23]
$\Delta\tilde{F}_{1V}^\gamma$	+0.043 -0.041	+0.047, 200 fb ⁻¹ -0.047, 200 fb ⁻¹	$\Delta\tilde{F}_{1V}^Z$	+0.24 -0.62	+0.012, 200 fb ⁻¹ -0.012, 200 fb ⁻¹
$\Delta\tilde{F}_{1A}^\gamma$	+0.051 -0.048	+0.011, 100 fb ⁻¹ -0.011, 100 fb ⁻¹	$\Delta\tilde{F}_{1A}^Z$	+0.052 -0.060	+0.013, 100 fb ⁻¹ -0.013, 100 fb ⁻¹
$\Delta\tilde{F}_{2V}^\gamma$	+0.038 -0.035	+0.038, 200 fb ⁻¹ -0.038, 200 fb ⁻¹	$\Delta\tilde{F}_{2V}^Z$	+0.27 -0.19	+0.009, 200 fb ⁻¹ -0.009, 200 fb ⁻¹
$\Delta\tilde{F}_{2A}^\gamma$	+0.16 -0.17	+0.014, 100 fb ⁻¹ -0.014, 100 fb ⁻¹	$\Delta\tilde{F}_{2A}^Z$	+0.28 -0.27	+0.052, 100 fb ⁻¹ -0.052, 100 fb ⁻¹


Table 12: Sensitivities achievable at 68.3% CL for the anomalous ttV ($V = \gamma, Z$) couplings $\tilde{F}_{1V,A}^V$ and $\tilde{F}_{2V,A}^V$ of Eq. (59) at the LHC for integrated luminosities of 300 fb⁻¹, and the ILC with $\sqrt{s} = 500$ GeV (taken from Ref. [23]). Only one coupling at a time is allowed to deviate from its SM value. Table and caption have been copied from [16].

arXiv:hep-ph/0601112v2


Top Quark

Threshold Region

M.Stahlhofen Top Phys WS 2012



Theory improving!





Expected accuracies

$$\Delta m_t = 34 \text{ MeV}$$


$$\Delta \alpha_s(m_Z) = 0.0023$$

$$\Delta \Gamma_t = 42 \text{ MeV}$$

Threshold scan alone



F.Simon Top Phys WS 2012



+ A_{FB} & Top Momentum

$$\Delta m_t = 19 \text{ MeV}$$

$$\Delta \alpha_s(m_Z) = 0.0012$$

$$\Delta \Gamma_t = 32 \text{ MeV}$$

arXiv:hep-ph/0601112v2

$$\Delta m_t(\overline{MS}) \simeq 100 \text{ MeV}$$



Vacuum Stability of the SM

With the discovered 126 GeV Higgs boson, λ becomes negative below Planck Scale

Cut off $\Lambda = 10^7 - 10^{15}$ GeV
large uncertainty comes
from large Δm_t

At ILC, $\Delta m_t \approx 30$ MeV is expected
Cutoff Λ can be better determined

At Planck Scale, $\lambda(M_{pl}) < 0$, but the theory satisfies the condition of
the meta-stable vacuum



arXiv:1205.6497, Degraasi et al


Last but Not Least

- In this talk I have been focusing on the case where $X(125\text{GeV})$ alone would be the probe for BSM physics, but there is a good chance for the higher energy run of LHC to bring us more.
- It is also very important to stress that ILC, too, is an energy frontier machine. It will access the energy region never explored with any lepton collider. There can be a zoo of new uncolored particles or new phenomena that are difficult to find at LHC but can be discovered and studied in detail at ILC.
- Examples
 - Natural SUSY : naturalness prefers μ not far above 100GeV
 - > light chargino/neutralinos will be higgsino-dominant and nearly degenerate
 - > typically Δm of a few GeV or less (very difficult at LHC)
 - > Δm as small as 50MeV possible with ISR tagging at ILC
 - > If $\Delta m=800\text{MeV}$ --> possible to measure m to 1.5GeV and Δm to 20MeV
 - > ILC will also be a Higgsino factory!
 - Possible anomalies in precision studies of properties of top, W/Z, and two-fermion processes
- Whatever new physics is awaiting for us, clean environment, polarized beams, and excellent jet energy resolution to reconstruct W/Z/t/H in their hadronic decays will enable us to uncover the nature of the new physics through model-independent precision measurements.

Backup


Coupling Precisions

Running Scenarios



Self-coupling Measurement

Weighting Method to Enhance the Sensitivity to λ



$$\frac{d\sigma}{dx} = B(x) + \lambda I(x) + \lambda^2 S(x)$$

irreducible
interference
self-coupling

Observable: weighted cross-section

$$\sigma_w = \int \frac{d\sigma}{dx} w(x) dx$$

Equation for the optimal $w(x)$ (variational principle):


$$\sigma(x)w_0(x) \int (I(x) + 2S(x))w_0(x)dx = (I(x) + 2S(x)) \int \sigma(x)w_0^2(x)dx$$

General solution:



$$w_0(x) = c \cdot \frac{I(x) + 2S(x)}{\sigma(x)}$$

c: arbitrary normalization factor

optimal weighing function



Expected Coupling Precision as a Function of Ecm




ILD DBD Study (Junping Tian)

Sensitivity Factor

$$\frac{\Delta\lambda}{\lambda} = F \cdot \frac{\Delta\sigma}{\sigma}$$

$F=0.5$ if no BG diagrams there

BG diagrams dominate at high E_{cm}



$\Rightarrow F$ grows quickly with E_{cm} !

Coupling Precision

ZHH : optimal $E_{cm} \sim 500$ GeV

though the cross section maximum is at around $E_{cm} = 600$ GeV

$\nu\nu HH$:

Precision slowly improves with E_{cm}

Expected Coupling Precision as a Function of Ecm

