


# Higgs recoil mass analysis: status and prospects

Taikan Suehara (Kyushu)  
A. Miyamoto (KEK), T. Tomita (Kyushu)

# Recoil in ILC

Direct measurement of  $g_{ZZH}$  with high precision  
NOT the “ratio” of Higgs couplings  
NOT suffered from QCD uncertainty




# Recoil mass

$$m_{\text{recoil}}^2 = (\sqrt{s} - E_{\ell\ell})^2 - |\vec{p}_{\ell\ell}|^2$$

- Using 4-momentum conservation (not possible in LHC)
- Excellent mass resolution (lepton only: no jet ambiguity)  $\sim 30$  MeV in ILC (1<sup>st</sup> phase only)
- Usable as a cut to separate the main background ZZ (recoil mass peaked at 91.2 GeV)

only looking 2f from Z



# global fit of higgs couplings

Snowmass energy frontier report

Model independent

Facility		ILC		ILC(LumiUp)
$\sqrt{s}$ (GeV)	250	500	1000	250/500/1000
$\int \mathcal{L} dt$ (fb $^{-1}$ )	250	+500	+1000	1150+1600+2500
$P(e^-, e^+)$	(-0.8, +0.3)	(-0.8, +0.3)	(-0.8, +0.2)	(same)
$\Gamma_H$	11%	5.9%	5.6%	2.7%
$BR_{inv}$	< 0.69%	< 0.69%	< 0.69%	< 0.32%
$\kappa_\gamma$	18%	8.4%	4.1%	2.4%
$\kappa_g$	6.4%	2.4%	1.8%	0.93%
$\kappa_W$	4.8%	1.4%	1.4%	0.65%
$\kappa_Z$	1.3%	1.3%	1.3%	0.61%
$\kappa_\mu$	—	—	16%	10%
$\kappa_\tau$	5.7%	2.4%	1.9%	0.99%
$\kappa_c$	6.8%	2.9%	2.0%	1.1%
$\kappa_b$	5.3%	1.8%	1.5%	0.74%
$\kappa_t$	—	14%	3.2%	2.0%

Taikan Suehara, Geneva **limited by  $\sigma_{ZH}$  @ 250 GeV**

# Further improvements

- $llh$  250  $\rightarrow$  3.0% (model independent)  
/ 2.6% (model dependent)
- Combining other modes
  - Higher energies: ILC500
  - Jet recoil  $qqh$ 
    - 350/500 GeV
    - 250 GeV

# lh at 500 GeV

- Cross section ( $\mu\mu h$ ) eL(0.8) pR(0.3)
  - 250 GeV: 10 fb
  - 500 GeV: 3.3 fb
  - Combining two may improve the resolution
- Recoil mass
  - Smearred in 500 GeV
- Background
  - Large t-channel diagram in 500 GeV
- Analysis
  - Almost the same as 250 GeV

# Cuts

Almost same as Lol

- Lepton ID
- Z mass (81.2 to 101.2 GeV)
- di-lepton  $p_T > 20$  GeV
- recoil mass (115-250  $\mu\mu$ , 100-250 ee)
- acoplanarity ( $\pi \pm 0.1$  rad vetoed)
- $p_T$  balance  
|di-lepton  $p_T - p_T$  of the most energetic neutral particle|  $> 10$  GeV


# Likelihood

Same as Lol

- di-lepton  $p_T$
- $\cos\theta$  of di-lepton
- acolinearity
- Z mass

TMVA used

- BDT & likelihood gives similar results
- Likelihood adopted







# mumu channel: cut table

cut	eeh	mmh	ooh	2f	4f	6f	o
m>=2	128	1691	3490	1.03e+6	966334	56719	361231
z mass	7.2	1499	141	225706	111908	4203	2765
ptdl>20	6.2	1492	124	64342	95766	3861	1573
recoil	1.6	1211	11	15198	30594	232	68
acop	1.3	1208	11	14598	28447	227	56
ptbal	1.2	1206	10	4544	27618	217	35
like>0.8	0	997	6.5	1632	3345	63	1.3

# Recoil fit


- BG fit (Gaus \* 2<sup>nd</sup> pol for mm, linear for ee)
- Fit will all parameters free
  - GPET with 5 parameters
    - Gaus (left side)
    - Gaus + expo (right side)
  - background distribution from fit function
    - to avoid large fluctuation due to the small stat
- Toy-MC – 10000 times
  - Poisson from data (sig), func (bg)
  - Fit with fixing shape parameters (mean and amplitude free)

# mumu: recoil fit




# ee: brems recovery

all neutral particle at  $\cos\theta > 0.99$  are added







# ee: cut table

cut	eeh	mmh	ooh	2f	4f	6f	o
e>=2	4582	76	6241	500994	1.6e+4	94156	413309
z mass	1654	2.3	559	18976	151032	14827	15562
ptdl>20	1627	2.0	459	12987	115384	12983	11496
recoil	1126	0.6	14	478	31324	443	1278
acop	1118	0.6	14	426	30123	429	1193
ptbal	1113	0.6	14	388	28417	417	1154
l>0.9	757	0.6	1.4	26	2763	66	84



# ee: recoil fit



# Summary of Ilh 500

- mmh: 6.49%
- eeh: 7.10%
- combined: 4.8%


# qqh recoil

- llh: only 6% of Z ( $e+\mu$ )
- qqh: 70% → hopeful!
- $\nu\nu h$ ,  $\tau\tau h$ : impossible to get recoil
  
- Jet clustering
  - efficiency can be different among higgs decay modes (6-jet, 4-jet, 2-jet)
- Jet energy resolution
  - Worse than leptons
  - Wider peak of recoil mass

# qqh 350: W/Z separation

A. Miyamoto

sales point of  
PFA calorimeter




# 350 GeV result

A. Miyamoto

- $k_t$  clustering w/  $R=1.2$  (may not be optimal)
- Basic cuts of Z mass etc.

## Selection statistics


	eR80.pR30 (300fb <sup>-1</sup> )	eL80.pR30 (300fb <sup>-1</sup> )
qqh	7581	11263
2f_h	13049	33326
4f_h	15726	109011
4f_sl	10767	65971
4f_l	597	1755
Higgs	1313	1975
aa/ae/ea	3971	7320
tt	4124	8441
all-bkg	49546	227800
S/sqrt(S+B)	31.72	23.04
sqrt(S+B)/S	0.03153	0.04341
S/N	0.153	0.049



150 fb<sup>-1</sup> each to  
-80/+30 & +80/-30%

$\Delta\sigma/\sigma \sim 3.6\%$

# 500 GeV



Event statistics for 500 fb<sup>-1</sup>,  
eL80.pR30 at 500 GeV

	Process	Nevents	Fraction(%)
S	qqh	11113	
	ffh	338	0.19
B	2f	47377	27.01
	4f	121086	69.02
	6f	6357	3.62
	aa_2f	91	0.05
	ae/ea	186	0.11

Hard to see recoil mass peak,  
but excess due to qqh could be seen

$$S/N=0.0633$$

$$\sqrt{S+N}/S=0.0389$$

$$\Delta\sigma/\sigma \sim 3.9\%$$

# Summary


---

- A possibility to measure  $\sigma_{ZH}$  using using  $Z \rightarrow qq$  mode was investigated.
- At 350 GeV, Higgs peak in jet recoil mass can be seen. Combining eL80/eR30  $150\text{fb}^{-1}$  and eR80/eL30  $150\text{fb}^{-1}$ ,  $\Delta\sigma/\sigma \sim 3.6\%$  is expected.
- At 500 GeV, hard to see Higgs peak in jet recoil mass distribution. But from event excess in qqh like events,  $\Delta\sigma/\sigma \sim 3.9\%$  is expected for  $500\text{fb}^{-1}$ , eL80.pR30 beam polarization.
- Further improvement may be possible by more sophisticated analysis.
- A possibility to use  $Z \rightarrow qq$  at 250 GeV should be investigated.


# qqh 250

- 250 GeV analysis ( $\sigma_{ZH} \sim \text{maximum}$ )
  - Durham y-fix clustering (should be optimized)
  - or dedicated  $Z \rightarrow qq$  finder from PFOs
- Difference of efficiency in Higgs decay modes should be investigated (as systematic error study)
- Also used for detector optimization
  - performance with various jet energy resolution will be investigated

## qqh 250: some plots



$y_{\text{cut}} = 0.001$




$y_{\text{cut}} = 0.005$

black: all  
 red:  $H \rightarrow bb$   
 blue:  $H \rightarrow WW^*$   
 black:  $H \rightarrow \text{others}$


Z mass

# qqh 250: some plots



$y_{\text{cut}} = 0.001$


The combination with mass nearest to Z is selected



$y_{\text{cut}} = 0.005$

black: all  
red:  $H \rightarrow bb$   
blue:  $H \rightarrow WW^*$   
black:  $H \rightarrow \text{others}$


# recoil mass qqh 250: some plots



$y_{\text{cut}} = 0.001$

The combination with  
mass nearest to  $Z$  is selected

cut on  $81.2 < m_Z < 101.2$



$y_{\text{cut}} = 0.005$

black: all

red:  $H \rightarrow b\bar{b}$

blue:  $H \rightarrow W W^*$

black:  $H \rightarrow \text{others}$

# qqh 250 summary

- Recoil mass peak observed
- will see WW/ZZ events (before LCWS)
- Optimization with ycut
- Kinematic fit
- Preliminary number? (not sure)
- Dedicated  $Z \rightarrow qq$  finder (after LCWS)