


CLIC Higgs capabilities and prospects for BSM physics

Philipp Roloff (CERN)
on behalf of the CLICdp collaboration



International Workshop on Future Linear Colliders (LCWS14)
Belgrade, Serbia, 06/10/2014

Overview:

- Introduction
 - CLIC Higgs capabilities
- Prospects for BSM physics
- Summary and conclusions

CLIC energy stages

CLIC would be implemented in stages:

- Optimised running conditions over a wide energy range
- **The energy stages are defined by physics** (with additional technical considerations)
→ The strategy can be adapted to discoveries at the LHC at 13/14 TeV

Example scenario assumed for this talk:


- **Stage 1:** 350 / 375 GeV, 500 fb⁻¹
(under discussion, [see CLICdp parallel meeting](#))
SM Higgs physics, $t\bar{t}$ threshold scan

- **Stage 2:** 1.4 TeV, 1.5 ab⁻¹
Targeted at BSM physics,
rare Higgs processes and decays

- **Stage 3:** 3 TeV, 2 ab⁻¹
Targeted at BSM physics,
rare Higgs processes and decays
(each stage corresponds to 4-5 years)

New CLIC staging baseline

→ [see next talk by Steinar Stapnes](#)





Detector benchmark studies


- Studies in this talk obtained using the CLIC_ILD and CLIC_SiD detector concepts

- New CLIC detector concept in preparation


More details → [see talks by Sophie Redford, Rosa Simoniello](#)



- Pile-up from $\gamma\gamma \rightarrow \text{hadrons}$ interactions overlaid to the physics events




- 1.3(3.2) events per BX at 1.4(3) TeV
- Suppressed using **timing capabilities** of the detectors and **hadron-collider type jet algorithms**




CLIC Higgs capabilities:

- Single Higgs production
- Processes at high energy
 - Combined analysis


Single Higgs production at CLIC



**Higgs-
strahlung**




Single Higgs production at CLIC



Some numbers

	350 GeV	1.4 TeV	3 TeV
L_{int}	500 fb^{-1}	1.5 ab^{-1}	2 ab^{-1}
# ZH events	68 000	20 000	11 000
# $H\nu_e\bar{\nu}_e$ events	17 000	370 000	830 000
# He^+e^- events	3 700	37 000	84 000





- Large samples of Higgs bosons produced at CLIC
- Measurements at high energy benefit from good detectors in the forward region

- Benchmark studies assume unpolarised beams

Polarization $P(e^-) : P(e^+)$	Enhancement factor	
	$e^+e^- \rightarrow ZH$	$e^+e^- \rightarrow H\nu_e\bar{\nu}_e$
unpolarized	1.00	1.00
-80% : 0%	1.18	1.80

Higgsstrahlung at 350 GeV (1)

Higgsstrahlung process





HZ events can be identified from Z recoil mass
→ **model independent** measurements of the g_{HZZ} coupling

$$\Delta(\sigma_{HZ}) / \sigma_{HZ} \approx 4\% \rightarrow \Delta(g_{HZZ}) / g_{HZZ} \approx 2\% \quad \text{from } Z \rightarrow \mu^+\mu^- \text{ and } Z \rightarrow e^+e^-$$

Higgsstrahlung at 350 GeV (2)


Higgsstrahlung process



- Substantial improvement using hadronic Z decays
- Challenge: $Z \rightarrow q\bar{q}$ reconstruction may depend on Higgs decay mode
- Even extreme variations of the SM Higgs BRs lead to bias $\leq \frac{1}{2}$ stat. error

$$\Delta(\sigma_{HZ}) / \sigma_{HZ} \approx 1.8\% \rightarrow \Delta(g_{HZZ}) / g_{HZZ} \approx 0.9\% \quad \text{from hadronic Z decays}$$

$\sigma \times \text{BR}$ measurements at 350 GeV



Higgs parallel session:

$H \rightarrow b\bar{b}/c\bar{c}/gg$: [Marco Szalay](#)

$H \rightarrow WW^*$: [Mila Pandurovic](#)

$H \rightarrow \tau^+\tau^-$: [Ph.R.](#)

Measurement	Observable	Stat. precision
$\sigma(HZ) \times \text{BR}(H \rightarrow \tau^+\tau^-)$	$g_{HZZ}^2 g_{H\tau\tau}^2 / \Gamma_H$	6.2%
$\sigma(HZ) \times \text{BR}(H \rightarrow b\bar{b})$	$g_{HZZ}^2 g_{Hbb}^2 / \Gamma_H$	1% (estimated)
$\sigma(HZ) \times \text{BR}(H \rightarrow c\bar{c})$	$g_{HZZ}^2 g_{Hcc}^2 / \Gamma_H$	5% (estimated)
$\sigma(HZ) \times \text{BR}(H \rightarrow gg)$		6% (estimated)
$\sigma(HZ) \times \text{BR}(H \rightarrow WW^*)$	$g_{HZZ}^2 g_{HWW}^2 / \Gamma_H$	2% (estimated)
$\sigma(H\nu_e \bar{\nu}_e) \times \text{BR}(H \rightarrow b\bar{b})$	$g_{HWW}^2 g_{Hbb}^2 / \Gamma_H$	3% (estimated)


Assuming
unpolarised beams

In addition: $\text{BR}(H \rightarrow \text{inv.}) < 0.97\%$ at 90% C.L.

Measurements using $H\nu_e\bar{\nu}_e$ events

Large Higgs samples produced in WW fusion at high energy:

- Precision measurements of $\sigma \times \text{BR}$
- Access to rarer decay modes



Measurement	Observable	Stat. precision (1.4 TeV)	Stat. precision (3 TeV)
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow \tau^+\tau^-)$	$g_{HWW}^2 g_{H\tau\tau}^2 / \Gamma_H$	4.2%	tbd
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow b\bar{b})$	$g_{HWW}^2 g_{Hbb}^2 / \Gamma_H$	0.3%	0.2%
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow c\bar{c})$	$g_{HWW}^2 g_{Hcc}^2 / \Gamma_H$	2.9%	2.7%
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow gg)$		1.8%	1.8%
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow \mu^+\mu^-)$	$g_{HWW}^2 g_{H\mu\mu}^2 / \Gamma_H$	38%	16%
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow \gamma\gamma)$		15%	tbd
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow Z\gamma)$		42%	tbd
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow ZZ^*)$	$g_{HWW}^2 g_{HZZ}^2 / \Gamma_H$	3% (estimated)	2% (estimated)
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow WW^*)$	g_{HWW}^4 / Γ_H	1.4%	0.9% (estimated)

Assuming unpolarised beams




Higgs parallel session: $H \rightarrow ZZ^*$: Gordana Milutinovic-Dumbelovic,
 $H \rightarrow WW^*$: Mila Pandurovic, $H \rightarrow \tau^+\tau^-$ P.R.

Precision measurements

$H \rightarrow b\bar{b}/c\bar{c}/g\bar{g}$:

- Separation of the different hadronic final states using precise flavour tagging
- $H \rightarrow c\bar{c}$ and gg impossible at hadron colliders
- In addition, the Higgs mass can be extracted from the $H \rightarrow b\bar{b}$ invariant mass distribution ($\pm 40\text{MeV}$ at 1.4 TeV, $\pm 33\text{MeV}$ at 3 TeV)


Measurement	1.4 TeV	3 TeV
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow b\bar{b})$	$\pm 0.3\%$	$\pm 0.2\%$
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow c\bar{c})$	$\pm 2.9\%$	$\pm 2.7\%$
$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow g\bar{g})$	$\pm 1.8\%$	$\pm 1.8\%$



Rare decays



$$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow \mu^+\mu^-):$$

- Very small BR ($\approx 0.022\%$)
- Requires precision tracking
- $\Delta(\sigma \times \text{BR}) = 38\%(16\%)$ at 1.4(3) TeV



$$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow \gamma\gamma):$$


- $\text{BR}(H \rightarrow \gamma\gamma) \approx 0.23\%$
- $\Delta(\sigma \times \text{BR}) = 15\%$ at 1.4 TeV




$$\sigma(H\nu_e\bar{\nu}_e) \times \text{BR}(H \rightarrow Z\gamma):$$

- $\text{BR}(H \rightarrow Z\gamma) \approx 0.16\%$
- Hadronic Z decays usable (in contrast to hadron colliders)
- $\Delta(\sigma \times \text{BR}) = 42\%$ at 1.4 TeV


Other processes at higher energy




$t\bar{t}H$ production:
maximum at
around 800 GeV





Other processes at higher energy




$t\bar{t}H$ production:
maximum at
around 800 GeV



Double Higgs production:
requires high energy



The $t\bar{t}H$ final state at 1.4 TeV



→ The $t\bar{t}H$ cross section is **directly sensitive to the top Yukawa coupling $g_{t\bar{t}H}$**

Investigated final states:

“6 jets”: $t(\rightarrow qq\bar{b})\bar{t}(\rightarrow l\nu\bar{b})H(\rightarrow b\bar{b})$

“8 jets”: $t(\rightarrow qq\bar{b})\bar{t}(\rightarrow qq\bar{b})H(\rightarrow b\bar{b})$

→ **Four b-quarks in the final state**



Combination of both final states:

$$\Delta\sigma(t\bar{t}H) / \sigma(t\bar{t}H) = 8.4\%$$


$$\rightarrow \Delta g_{t\bar{t}H} / g_{t\bar{t}H} = 4.5\%$$

More details


→ [see talk by Sophie Redford in the Higgs session](#)



Double Higgs production at high energy



- The $HH\nu_e\bar{\nu}_e$ cross section is sensitive to the Higgs self coupling, λ , and the quartic $HHWW$ coupling
- Only 225 (1200) $e^+e^- \rightarrow HH\nu_e\bar{\nu}_e$ events at 1.4 (3) TeV
 \rightarrow high energy and luminosity crucial




Measurement	1.4 TeV	3 TeV
$\Delta(g_{HHWW})$	7% (preliminary)	3% (preliminary)
$\Delta(\lambda)$	32%	16%
$\Delta(\lambda)$ for $P(e^-) = -80\%$	24%	12%

CLIC Higgs studies

Channel	Measurement	Observable	Statistical precision		
			350 GeV 500 fb ⁻¹	1.4 TeV 1.5 ab ⁻¹	3.0 TeV 2.0 ab ⁻¹
ZH	Recoil mass distribution	m_H	120 MeV	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow \text{invisible})$	Γ_{inv}	0.6%	—	—
ZH	$H \rightarrow b\bar{b}$ mass distribution	m_H	tbd	—	—
Hv _e $\bar{\nu}_e$	$H \rightarrow b\bar{b}$ mass distribution	m_H	—	40 MeV*	33 MeV*
ZH	$\sigma(HZ) \times BR(Z \rightarrow \ell^+ \ell^-)$	g_{HZZ}^2	4.2%	—	—
ZH	$\sigma(HZ) \times BR(Z \rightarrow q\bar{q})$	g_{HZZ}^2	1.8%	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow b\bar{b})$	$g_{HZZ}^2 g_{Hbb}^2 / \Gamma_H$	1% [†]	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow c\bar{c})$	$g_{HZZ}^2 g_{Hcc}^2 / \Gamma_H$	5% [†]	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow gg)$	—	6% [†]	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow \tau^+ \tau^-)$	$g_{HZZ}^2 g_{H\tau\tau}^2 / \Gamma_H$	6.2%	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow WW^*)$	$g_{HZZ}^2 g_{HWW}^2 / \Gamma_H$	2% [†]	—	—
ZH	$\sigma(HZ) \times BR(H \rightarrow ZZ^*)$	$g_{HZZ}^2 g_{HZZ}^2 / \Gamma_H$	tbd	—	—
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow b\bar{b})$	$g_{HWW}^2 g_{Hbb}^2 / \Gamma_H$	3% [†]	0.3%	0.2%
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow c\bar{c})$	$g_{HWW}^2 g_{Hcc}^2 / \Gamma_H$	—	2.9%	2.7%
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow gg)$	—	—	1.8%	1.8%
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow \tau^+ \tau^-)$	$g_{HWW}^2 g_{H\tau\tau}^2 / \Gamma_H$	—	4.2%	tbd
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow \mu^+ \mu^-)$	$g_{HWW}^2 g_{H\mu\mu}^2 / \Gamma_H$	—	38%	16%
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow \gamma\gamma)$	—	—	15%	tbd
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow Z\gamma)$	—	—	42%	tbd
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow WW^*)$	g_{HWW}^4 / Γ_H	tbd	1.4%	0.9% [†]
Hv _e $\bar{\nu}_e$	$\sigma(Hv_e \bar{\nu}_e) \times BR(H \rightarrow ZZ^*)$	$g_{HWW}^2 g_{HZZ}^2 / \Gamma_H$	—	3% [†]	2% [†]
He ⁺ e ⁻	$\sigma(He^+ e^-) \times BR(H \rightarrow b\bar{b})$	$g_{HZZ}^2 g_{Hbb}^2 / \Gamma_H$	—	1% [†]	0.7% [†]
t \bar{t} H	$\sigma(t\bar{t}H) \times BR(H \rightarrow b\bar{b})$	$g_{Htt}^2 g_{Hbb}^2 / \Gamma_H$	—	8%	tbd
HHv _e $\bar{\nu}_e$	$\sigma(HHv_e \bar{\nu}_e)$	g_{HHWW}	—	7%*	3%*
HHv _e $\bar{\nu}_e$	$\sigma(HHv_e \bar{\nu}_e)$	λ	—	32%	16%
HHv _e $\bar{\nu}_e$	with -80% e ⁻ polarization	λ	—	24%	12%

*: preliminary
†: estimated

Putting it all together




Parameter	Measurement precision		
	350 GeV 500 fb ⁻¹	+ 1.4 TeV +1.5 ab ⁻¹	+3.0 TeV +2.0 ab ⁻¹
g_{HZZ}	0.8 %	0.8 %	0.8 %
g_{HWW}	1.8 %	0.9 %	0.9 %
g_{Hbb}	2.0 %	1.0 %	0.9 %
g_{Hcc}	3.2 %	1.4 %	1.1 %
$g_{H\tau\tau}$	3.7 %	1.7 %	1.5 %
$g_{H\mu\mu}$	—	14.1 %	5.6 %
g_{Htt}	—	4.1 %	≤ 4.1 %
g_{Hgg}^{\dagger}	3.6 %	1.2 %	1.0 %
$g_{H\gamma\gamma}^{\dagger}$	—	5.7 %	< 5.7 %
Γ_H	5.0 %	3.6 %	3.4 %

- Fully model-independent, **only possible at a lepton collider**
- All results limited by 0.8% from $\sigma(HZ)$ measurement
- The Higgs width is extracted with 5 – 3.5% precision

More details → [talk by Frank Simon in the Higgs/EW session](#)

Analysis similar to LHC experiments



Parameter	Measurement precision		
	350 GeV 500 fb ⁻¹	+ 1.4 TeV +1.5 ab ⁻¹	+3.0 TeV +2.0 ab ⁻¹
κ_{HZZ}	0.44 %	0.31 %	0.23 %
κ_{HWW}	1.5 %	0.17 %	0.11 %
κ_{Hbb}	1.7 %	0.37 %	0.22 %
κ_{Hcc}	3.1 %	1.1 %	0.75 %
$\kappa_{H\tau\tau}$	3.7 %	1.5 %	1.2 %
$\kappa_{H\mu\mu}$	—	14.1 %	5.5 %
κ_{Htt}	—	4.0 %	≤ 4.0 %
κ_{Hgg}	3.6 %	0.79 %	0.55 %
$\kappa_{H\gamma\gamma}$	—	5.6 %	< 5.6 %
$\Gamma_{H,md,derived}$	1.6 %	0.32 %	0.22 %

$$\kappa_i^2 = \frac{\Gamma_i}{\Gamma_i^{SM}}$$

No invisible decays:

$$\Gamma_{H,model} = \sum_i \kappa_i^2 \cdot BR_i^{SM}$$


Sub-percent precisions
at high energy

→ Results strongly dependent
on fit assumptions

-80% electron polarisation at 1.4 and 3 TeV

What's next?

- **Single Higgs production:** addressing a few channels not covered so far ($e^+e^- \rightarrow H\nu_e\bar{\nu}_e \rightarrow WW^*\nu_e\bar{\nu}_e$ at 350 GeV, $H \rightarrow \gamma\gamma$ at 3 TeV, ZZ fusion at 3 TeV)
- **Reanalysis of double Higgs production:** add the $HH \rightarrow b\bar{b}WW^*$ final state (40% more events compared to $HH \rightarrow b\bar{b}b\bar{b}$ alone)
- **Looking at differential distributions:**
example: **CP properties of the Higgs boson**
 - using $t\bar{t}H$ events:
extension of top Yukawa coupling study
 - using WW and ZZ fusion events:
large statistics at CLIC promising



More details

→ [see the CLICdp parallel session](#)

Snowmass Higgs WG report, [arXiv:1310.8361](#)


Prospects for BSM physics

- **Two approaches:**

1.) Pair production of new particles if $M \leq \sqrt{s} / 2$

→ CLIC especially attractive for electroweak states

→ Precision measurement of new particle masses and couplings




Many examples of SUSY particle production studied for CLIC CDR

2.) Indirect searches through precision observables

→ possibility to reach much higher mass scales


One of the priorities for future benchmarking studies

Investigated SUSY models




CDR Model I, 3 TeV:

- Squarks
- Heavy Higgs




CDR Model II, 3 TeV:

- Smuons, selectrons
- Gauginos



CDR Model III, 1.4 TeV:


- Smuons, selectrons
- Staus
- Gauginos



Wider applicability than only SUSY: Reconstructed particles can be classified simply as **states of given mass, spin and quantum numbers**


Reconstruction of SUSY particles

Endpoints of energy spectra:



$$\begin{aligned} m(\tilde{\mu}_R) &: \pm 5.6 \text{ GeV} \\ m(\tilde{e}_R) &: \pm 2.8 \text{ GeV} \\ m(\tilde{\nu}_e) &: \pm 3.9 \text{ GeV} \\ m(\tilde{\chi}_1^0) &: \pm 3.0 \text{ GeV} \\ m(\tilde{\chi}_1^\pm) &: \pm 3.7 \text{ GeV} \end{aligned}$$

$$e^+e^- \rightarrow \tilde{\mu}_R^+ \tilde{\mu}_R^- \rightarrow \mu^+ \mu^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$$



Jet reconstruction

Precision on the measured gaugino masses
(few hundred GeV):
1 - 1.5%

$$e^+e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 W^+ W^-$$

$$e^+e^- \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_2^0 \rightarrow hh \tilde{\chi}_1^0 \tilde{\chi}_1^0$$



$$e^+e^- \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_2^0 \rightarrow Zh \tilde{\chi}_1^0 \tilde{\chi}_1^0$$

Complex final states:

$$e^+e^- \rightarrow HA \rightarrow b\bar{b}b\bar{b}$$

$$e^+e^- \rightarrow H^+H^- \rightarrow t\bar{b}b\bar{t}$$

≈0.3% precision on
heavy Higgs masses



Summary of the SUSY studies

\sqrt{s} (TeV)	Process	Decay mode	SUSY model	Measured quantity	Generator value (GeV)	Stat. uncertainty
3.0	Sleptons	$\tilde{\mu}_R^+ \tilde{\mu}_R^- \rightarrow \mu^+ \mu^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$	II	$\tilde{\ell}$ mass	1010.8	0.6%
				$\tilde{\chi}_1^0$ mass	340.3	1.9%
		$\tilde{e}_R^+ \tilde{e}_R^- \rightarrow e^+ e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$		$\tilde{\ell}$ mass	1010.8	0.3%
				$\tilde{\chi}_1^0$ mass	340.3	1.0%
		$\tilde{\nu}_e \tilde{\nu}_e \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 e^+ e^- W^+ W^-$		$\tilde{\ell}$ mass	1097.2	0.4%
				$\tilde{\chi}_1^\pm$ mass	643.2	0.6%
3.0	Chargino Neutralino	$\tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 W^+ W^-$	II	$\tilde{\chi}_1^\pm$ mass	643.2	1.1%
		$\tilde{\chi}_2^0 \tilde{\chi}_2^0 \rightarrow h/Z^0 h/Z^0 \tilde{\chi}_1^0 \tilde{\chi}_1^0$		$\tilde{\chi}_2^0$ mass	643.1	1.5%
3.0	Squarks	$\tilde{q}_R \tilde{q}_R \rightarrow q \bar{q} \tilde{\chi}_1^0 \tilde{\chi}_1^0$	I	\tilde{q}_R mass	1123.7	0.52%
3.0	Heavy Higgs	$H^0 A^0 \rightarrow b \bar{b} b \bar{b}$	I	H^0/A^0 mass	902.4/902.6	0.3%
		$H^+ H^- \rightarrow t \bar{b} b \bar{t}$		H^\pm mass	906.3	0.3%
1.4	Sleptons	$\tilde{\mu}_R^+ \tilde{\mu}_R^- \rightarrow \mu^+ \mu^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$	III	$\tilde{\ell}$ mass	560.8	0.1%
				$\tilde{\chi}_1^0$ mass	357.8	0.1%
		$\tilde{e}_R^+ \tilde{e}_R^- \rightarrow e^+ e^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$		$\tilde{\ell}$ mass	558.1	0.1%
				$\tilde{\chi}_1^0$ mass	357.1	0.1%
		$\tilde{\nu}_e \tilde{\nu}_e \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 e^+ e^- W^+ W^-$		$\tilde{\ell}$ mass	644.3	2.5%
				$\tilde{\chi}_1^\pm$ mass	487.6	2.7%
1.4	Stau	$\tilde{\tau}_1^+ \tilde{\tau}_1^- \rightarrow \tau^+ \tau^- \tilde{\chi}_1^0 \tilde{\chi}_1^0$	III	$\tilde{\tau}_1$ mass	517	2.0%
1.4	Chargino Neutralino	$\tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 W^+ W^-$	III	$\tilde{\chi}_1^\pm$ mass	487	0.2%
		$\tilde{\chi}_2^0 \tilde{\chi}_2^0 \rightarrow h/Z^0 h/Z^0 \tilde{\chi}_1^0 \tilde{\chi}_1^0$		$\tilde{\chi}_2^0$ mass	487	0.1%

Precision studies of $e^+e^- \rightarrow \mu^+\mu^-$

Minimal anomaly-free Z' model:

Charge of the SM fermions
under $U(1)'$ symmetry:

$$Q_f = g_Y'(Y_f) + g_{BL}'(B-L)_f$$

Observables:



- total $e^+e^- \rightarrow \mu^+\mu^-$ cross section
- forward-backward-asymmetry
- left-right asymmetry
($\pm 80\%$ e^- polarisation)

If LHC discovers Z'
(e.g. for $M = 5$ TeV):

Precise measurement of the
effective couplings


Otherwise:

Discovery reach up to tens of TeV
(depending on the couplings)



Composite Higgs bosons

- Higgs as **composite bound state of fermions**
- m_ρ : mass of the vector resonance of the composite theory
- $\xi = (v / f)^2$ measures the strengths of the Higgs interactions



CLIC provides an indirect probe of a Higgs composite scale of 70 TeV

What's next?

- **Interesting SUSY signatures not yet studied for CLIC:**

- 1.) Gauginos/Higgsinos with small mass splittings

→ Main signal: γ + missing energy + soft particles
(challenging in the presence of beam-induced backgrounds)

- 2.) Top squark production

e.g. $\tilde{t}_1 \rightarrow t + \tilde{\chi}_1^0 \rightarrow$ boosted top quarks

- **Model-independent searches for Dark Matter**

using the γ + missing energy final state

- **Higher-dimensional effective operators**

- **More on compositeness, weakly interacting exotica, ...**

Crucial: need to be ready to respond to
theoretical interpretation of new LHC data

More details
→ see CLICdp
parallel session

Precision measurements: top

$t\bar{t}$ events:

- So far focussed on top mass at lower energies (350 GeV and 500 GeV)
- Explore **potential of $t\bar{t}$ events to probe for new physics**, examples:

- A_{FB}^t (and A_{FB}^b)
- $\sin^2\theta_W$
- top quark couplings to γ , W and Z

→ At high energy and possibly for the first stage

V_{tb} from $e^-\gamma \rightarrow \bar{t}b\nu_e$ at high energy:

200000 $e\gamma \rightarrow tb\nu$ events expected at 3 TeV

(no $t\bar{t}$ contribution in contrast
to $e^+e^- \rightarrow t\bar{b}\nu_e$ or $\gamma\gamma \rightarrow t\bar{b}\nu_e$)

More details


→ [see CLICdp parallel session](#)

Precision measurements: EW

Triple and quartic gauge couplings using $e^+e^- \rightarrow W^+W^- (\nu\bar{\nu}/e^+e^-)$:
Important to choose parametrisation comparable to other studies/experiments!

W boson mass determination at high energy:

- Large samples of single W events produced at high-energy CLIC
- Potential for competitive measurement of M_W using $W^\pm \rightarrow q\bar{q}$
- Need full simulation study to understand the impact of systematic effects



More details
→ [see CLICdp parallel session](#)

Summary and conclusions

Summary and conclusions: Higgs


- The first stage of a CLIC collider at 350 GeV provides **precise determinations of the absolute values of many Higgs boson couplings**
- Subsequent **high-energy running**, here assumed at 1.4 and 3 TeV, improves the precision of many observables significantly and gives access to **rare Higgs decays**
- High-energy CLIC operation provides the potential to **measure the trilinear Higgs self-coupling at the 10% level**
- **Combined fits** to all measurements at 350 GeV, 350 GeV + 1.4 TeV and 350 GeV + 1.4 TeV + 3 TeV were performed to extract the Higgs couplings and width simultaneously
- **A comprehensive paper on (SM-)Higgs physics at CLIC is being completed**

Summary and conclusions: BSM

- CLIC operated at high-energy (**1.4 and 3 TeV**) provides significant discovery potential for BSM phenomena
- Measurement of the gaugino, slepton and heavy Higgs masses with $O(1\%)$ precision up to the kinematic limit ($M \approx 1.5$ TeV)
- In addition to studying new particles directly: sensitivity to New Physics at large scales (tens of TeV) through **precision measurements** (examples: Z' and composite models)
- Many more studies started / will start soon: also on BSM sensitivity through precision top / SM observables


Backup slides

If you want to know more...



CLIC Conceptual Design Report (CDR) Vol. 2: Physics and Detectors (mostly at 3 TeV)

[arXiv:1202.5940](https://arxiv.org/abs/1202.5940)



CLIC CDR Vol. 3: Staged construction, SUSY at 1.4 TeV, Z'

[arXiv:1209.2543](https://arxiv.org/abs/1209.2543)



Snowmass white paper: Most of the Higgs studies

[arXiv:1307.5288](https://arxiv.org/abs/1307.5288)
(last update: 01/10/2013)


2-beam acceleration scheme

Drive beam supplies RF power:

- 12 GHz bunch structure
- Low energy:
2.4 GeV – 240 MeV
- High current: **100 A**

Main beam for physics:

- High energy: **9 GeV – 1.5 TeV**
- Current: 1.2 A




Selected CLIC parameters

CLIC at 3 TeV	
L ($\text{cm}^{-2}\text{s}^{-1}$)	$5.9 \cdot 10^{34}$
Bunch separation	0.5 ns
#Bunches / train	312
Train duration	156 ns
Train rep. rate	50 Hz
Crossing angle	20 mrad
Particles / bunch	$3.72 \cdot 10^9$
σ_x / σ_y (nm)	$\approx 45 / 1$
σ_z (μm)	44


Drive timing requirements for CLIC detector

Very small beam profile at the interaction point




CLIC: trains at 50 Hz, 1 train = 312 bunches, 0.5 ns apart

Beam related backgrounds



- e⁺e⁻ pairs
- $\gamma\gamma \rightarrow \text{hadrons}$



Coherent e⁺e⁻ pairs:

7 · 10⁸ per BX, very forward


Incoherent e⁺e⁻ pairs:

3 · 10⁵ per BX, rather forward


→ **Detector design issue**
(high occupancies)

$\gamma\gamma \rightarrow \text{hadrons}$

- “Only” 3.2 events per BX at 3 TeV
 - Main background in calorimeters and trackers
- **Impact on physics**




CLIC detector concepts




Background suppression

Triggerless readout of full bunch train:



1.) Identify t_0 of physics event in offline event filter

- Define reconstruction window around t_0
- All hits and tracks in this window are passed to the reconstruction
→ **Physics objects with precise p_T and cluster time information**




2.) Apply cluster-based timing cuts

- Cuts depend on particle-type, p_T and detector region
→ **Protects physics objects at high p_T**


In addition: hadron-collider type jet algorithms (FastJet)

Impact of the timing cuts

$e^+e^- \rightarrow t\bar{t}$ at 3 TeV with background from $\gamma\gamma \rightarrow \text{hadrons}$ overlaid



1.2 TeV background
in the reconstruction
window



100 GeV background
after timing cuts

Physics studies are based on Geant4 simulations including pile-up from $\gamma\gamma \rightarrow \text{hadrons}$