



# TASK FOR THIS WORKSHOP & AFTER WORKSHOP

Masakazu Kurata

The University of Tokyo

HLReco Week, 07/06/2015–07/10/2015

1

# LCFIPLUS IMPROVEMENT

- Now, focusing on:
  - Vertex finding efficiency improvement itself
  - Flavor separation in the case of 0vtx jet
  - Vertex Mass Recovery using  $\pi^0$ s
- Other topics especially using PID:
  - Soft lepton tagger for flavor tag using PID
  - Muon ID during jet clustering
- We will incorporate above topics into LCFIPlus
  - Short term task
  - Middle term task

# WHICH CAN BE ACHIEVED DURING THIS WORKSHOP?

## ○ Soft lepton tagger using PID

- Can be easily include into LCFIPlus
- Will be during this workshop
- Directly check PID output and apply the lepton selection

SIMPLE:  $PDG==11$  for electron,  $PDG==13$  for muon

- Need to check performance


## ○ AVF algorithm

- Only AVF part, we will include during this workshop
- Create flag to switch AVF/nominal  $\chi^2$  algorithm
- But AVF case needs BNess tagger for fake rejection...

# BNESS TAGGER & VERTEX MASS – MIDDLE PLAN?

- Slightly complicated
- Need to check this ROC curve improvement

- Did I use really correct variable?
- Please give me some time to check



- Need CNess? → I have already constructed CNess(just constructed)
- My feeling is this can be achieved after workshop


- Vertex mass recovery:

- Need pi0 reco! →Graham's pi0 reco. techinque? or continue to use my pi0 reco.?
- Need to check performance with Graham's reco. & AVF  
→retrain pi0vertexfinder is necessary

✂it takes some time!

## IN SHORT... (MY OPINION)

- We will be able to obtain during this workshop:
  - Soft lepton tagger
  - AVF code include
- After workshop:
  - BNess tagger(& CNess tagger?) after some check
- Near future
  - Vertex mass recovery
- Finally, final flavor tagger
- Good improvement about flavor tagging will be after above them



## BACK UPS




# ADAPTIVE VERTEX FITTING

- To introduce the effect of multi-vertex fitting
  - Introduce weight function to estimate which vertex a track belongs to
  - Weight function definition: k-th track's weight on n-th vertex

$$w_{nk} = \frac{e^{-\chi_{nk}^2/2T}}{e^{-\chi_{\text{cut}}^2/2T} + \sum_{i=1}^N e^{-\chi_{ik}^2/2T}}$$

- Parameter: temperature T
  - If T small, decision is like  $\chi^2$  minimization
  - If T large, multi-vertex effect becomes large (suppress the weight function)
- Apply it to associate IP tracks to secondary vertices:
- Algorithm: Adaptive Vertex Fitting



- Tracks will belong to the vertices when  $w_{nk} > 0.5$  (k-th track belongs to n-th vertex)
  - Try to fit more tracks than nominal algorithm

# VERTEX FINDING OF BJETS

- Common parameters are set at same values for comparison
- Same event sample(qqHH sample@500GeV) 19889 events
- 6 jet clustering, jet matching with MCtruth is performed
- Num. of jets

method	bjet with 2vtx	bjet with 1+1vtx	bjet with 1vtx	total
Nominal Algorithm	10577	9159	12804	32504
AVF&BNess	13461	6502	14256	34219

- Jets with vtx: **~5%** increased
  - Jets with 2vtx: **~24%** increased →move from 1+1
  - Jets with 1vtx: **~11%** increased
- Fake track rate per vtx: how many are fake tracks contaminated to vertices?
  - Seems fake singletrk is increased→need opt. and more selection

method	bjet with 2vtx	bjet with 1+1vtx	bjet with 1vtx
Nominal Algorithm	$0.011 \pm 0.0007$	$0.007 \pm 0.0006$	$0.025 \pm 0.001$
AVF&BNess	$0.010 \pm 0.0006$	$0.011 \pm 0.0009$	$0.021 \pm 0.001$



## VERTEX FINDING OF C JETS

- Common parameters are set at same values for comparison
- Same event sample(nnH sample@500GeV) 99432 events
  - H→cc: 6461 events
- 2 jet clustering, jet matching with MCtruth is performed
- Num. of vertices

method	cjet with 2vtx	cjet with 1+1vtx	cjet with 1vtx	total
Nominal Algorithm	43	165	6537	6745
AVF&BNess	84	215	6960	7259

- Total:  $\sim 7\%$  increased
- Vertex mis-ID eff. is increased(but, 2vtx jet has pure vertices)
  - Though num. of vertices is small
  - need additional selection for singletrk? (e.g.)vertex mass?)
- Fake track rate per vtx:



method	cjet with 2vtx	cjet with 1+1vtx	cjet with 1vtx
Nominal Algorithm	$0.00 \pm 0.00$	$0.012 \pm 0.006$	$0.0014 \pm 0.004$
AVF&BNess	$0.00 \pm 0.00$	$0.018 \pm 0.007$	$0.0013 \pm 0.004$

# BNESS TAGGER

- Flavor separation of 0vtx jet is most difficult situation
  - Only impact parameter implies the existence of secondary vertices for flavor separation
- BNess tagger will be worth trying in this case!
  - Developed in CDF
  - Focus on individual tracks and evaluate jet flavor only using single track
  - Track's potential for coming from heavy flavor particle(D&B meson and baryons) should be evaluated(using MVA)
- Difficulty in ILC
  - In CDF, it is important to separate b and other flavor → c quark separation is not required
  - In ILC, separation among b, c and other is very important → bc separation is a key for flavor tagger
- How is bc separation using BNess tagger?



# BNESS OUTPUT

- Collect **Highest score BNess** track in 0vtx jets
- Final BNess is defined as BNess(bl)+ BNess(bc)
- Well separated between bjets and l jets
- Difference can be seen between bjets and cjets



# RESULTS OF BNESSTAGGER ON FLAVOR TAGGING

- Construct a “toy” flavor tagger
  - Convert nominal input variables to BNesstagger variables
  - Compare with ROC curve
- For bc separation, some improvement can be obtained
- For bl separation, becomes too good? under investigation
- b-l separation will be very good!
- Need optimization
- Especially, precise study of b-c-l flavor separation is necessary





# BNESS TAGGER FOR FAKE TRACK REJECTION

- Loosen the track selection to try to attach as many tracks as possible to vertices
  - Fake track rate will be increased
- To reject fakes, BNess tagger is used
  - So far, just use BNess(bl)
- So far, only BNess is checked



→ some bias for D meson tracks?

Example: looking for single track




# VERTEX MASS RECOVERY

- Using  $\pi^0$ s which escape from vertices
  - Need to choose good  $\pi^0$  candidates –construct  $\pi^0$  vertex finder
  - Key issue – $\pi^0$  kinematics, very collinear to vertex direction




- Particle ID is the other key to classify vertices
  - Different particle patterns have different vertex mass patterns
- Construct  $\pi^0$  Vertex finder using MVA
  - Identify which vertex  $\pi^0$ s are coming from



# VTX MASSES OF BJETS IN DOUBLE-HIGGS PROCESS

- Vtx mass distributions for each vertex pattern(ntrk)
  - These results are the outputs of LCFIPlus(unofficial ver.)!
  - Difference is limited by mis-pairing of gammas and mis-attachment of pi0s





# PI0 RECO USING NAÏVE BAYES FOR VERTEX MASS RECOVERY



- Good pairing eff. & mis-pairing eff.

	Correct pair	Wrong pair
eff. (%)	$46.0 \pm 0.3$	$54.0 \pm 0.4$

- Kin. plots of pi0 reco. results



MC truth  
Pi0 finder




Num. of pi0s to be reconstructed  
Num. of pi0s from pi0 finder

- Integrate pi0 reconstruction?



# JET CLUSTERING WITH BEAM BACKGROUND REJECTION

- Now in LCFIPlus, Valencia jet clustering is available!
- We also include Durham jet clustering with beam b.g. rejection
  - Assumed very large energy jet exists in beam direction
- Compare the performance between Durham, Kt and Valencia



w/o beam b.g. rejection

Kt

Valencia

Durham

$\nu \nu Z@500\text{GeV}$

2 jet clustering


Parameters are tuned  
for better result

## OTHER TOPICS

- Other change from DBD LCFIPlus:
  - Automatic creation of Joint Probability plots
    - We can calibrate for that variable
    - Performance check for new joint probability is ongoing
- Paper available:
  - arxiv: 1506:08371
  - NIM paper has been submitted

# SUMMARY AND PROSPECTS

- For flavor tagging improvement:
  - New vertexing algorithm(AVF) will provide better vertex finding efficiency
  - BNesstagger will give some improvement for 0vtx jet flavor separation
  - There seems hope for attaching  $\pi^0$ s to vertices to recover vertex mass
- So far, AVF will provide **4–7%** improvement of vertex finding in bjets
  - Need to check the bias of fake track rejection using BNesstagger
  - Vertex quality check is necessary
  - This study will lead to vertex charge assignment improvement
- 0vtx jet case will improve well not only b–c separation, but also b–l separation
  - More Precise study of b–c–l separation is necessary
- Vertex mass recovery is reasonable
  - Will provide better flavor tagger
  - Of course, many checks are necessary
- Finally, incorporate all the ideas and check the final flavor tagging effs.in LCFIPlus!




## BACK UPS2

20

# TRACK MVA(BNESS)

- To identify track which comes from heavy flavor particle  
→ using MVA
  - Signal: tracks which come from B mesons or B baryons
  - Background: tracks produced in hadronization process
- Most significant tracks with both plus and minus signed impact parameters in a jet are collected

- Significance:  $sig = \sqrt{\left(\frac{d_0}{\sigma}\right)^2 + \left(\frac{z_0}{\sigma}\right)^2}$



# VTX MASSES

- Vtx mass distributions for each vertex pattern(ntrk)
  - not so bad
  - Difference is coming from **mis-pairing of gammas** and mis-attachment of pi0s

