

Searching for new light scalars at the ILC

Yan Wang (DESY, IHEP), Jenny List (DESY), Mikael Berggren (DESY),
Shin-ichi Kawada (DESY)

ALCW 2018 — Fukuoka

May 28, 2018



HELMHOLTZ
RESEARCH FOR GRAND CHALLENGES



Motivation

The higgs boson found at 2012: the SM Higgs?




Many BSMs predict one or more extra scalars:

- ▶ 2HDM, NMSSM, Randall Sundrum model
- ▶ a scalar lighter than 125 GeV is well motivated.

JHEP 12 (2016) 068


survived after indirect + LEP + LHC constrains



- ▶ LEP/LHC constrains rely on the model details: CP, mass hierarchy, couplings, etc.
- ▶ want a better result?



ILC — The International Linear Collider



- ▶ ILC properties:
 - ▶ e^+e^- collider, with polarized beams (e^- : ± 0.8 , e^+ : ∓ 0.3).
- ▶ ILC running scenario for 22 years:
 - ▶ $\sqrt{s} = 250$ GeV and $\int Ldt = 2/ab$ for the first stage
- ▶ Energy-upgradable
 - ▶ $\sqrt{s} = 350$ GeV and $\int Ldt = 0.2/ab$
 - ▶ $\sqrt{s} = 500$ GeV and $\int Ldt = 4/ab$
 - ▶ if possible 1 TeV.

Comparing LEP/LHC and ILC

- ▶ comparing with LEP: ILC is sensitive to lighter scalars with smaller hZZ coupling.

	LEP	ILC	improvement
$\max \sqrt{s}$ (GeV)	189-209	250	
m_h region (GeV)	<115	<125	
luminosity	totally 2461 pb^{-1}	2000 fb^{-1}	recoil mass
polorization	×	✓	angle correlation
detector	ALEPH,DELPHI,OPAL,L3	ILD, SID	resolution
searching channel	2b2q,2b2 ν ,2b2l, $\tau\tau qq$	model independent	

Phys.: Conf. Ser. 110 042030


- ▶ comparing with LHC
 - ▶ LHC, complex initial states and backgrounds, $h \rightarrow \gamma\gamma/ZZ\dots$ channel, large uncertainties.
 - ▶ ILC, e^+e^- initial states, **clean environment, model-independent.**



The Recoil Method on SM Higgs at ILC


e^+e^- collider \rightarrow know the initial states behaviour \rightarrow recoil technique \rightarrow model independence

Higgsstrahlung process $e^+e^- \rightarrow Z + H/h$




- ▶ $M_{rec}^2 = (\sqrt{s} - E_{\mu\mu})^2 - |\vec{p}_{\mu\mu}|^2$
- ▶ $M_{\mu\mu} \sim M_Z, M_{rec} \sim M_{H/h}$

SM Higgs recoil mass distribution (ILD)



SM Higgs coupling for ILC and HL-LHC



the same method on light scalar searching, **SM $H \rightarrow$ a lighter h .**




The Recoil Method at LEP

LEP results (CERN-EP-2002-032):

- ▶ the OPAL detector
- ▶ Decay-mode independent searches for new scalar bosons
- ▶ energy & luminosity:
 - ▶ 91.2 GeV and 115.4 pb⁻¹ at LEP1
 - ▶ 161 to 202 GeV and **662.4 pb⁻¹** at LEP2.
- ▶ light higgs mass: 10 keV - 100 GeV

- ▶ $k = \frac{\sigma_{S^0 Z}}{\sigma_{H_{SM} Z}(m_{H_{SM}}=m_{S^0})}$



ILD (International Large Detector)


and full simulation of Signal and SM Background



- ▶ new trackers, calorimeters, 3.5T magnetic field, yoke for muon, forward system
- ▶ Requirements:
 - ▶ Impact parameter resolution:
 $\sigma_{r\phi} < 5 \oplus 10/(p \sin^{3/2}\theta)\mu\text{m}$
 - ▶ Momentum resolution:
 $\sigma_{1/p_T} < 2 * 10^{-5} \text{ GeV}^{-1}$
 - ▶ Energy resolution: $\sigma_E/E = 3 - 4\%$

The signal MC samples

- ▶ $M_{S^0} = 10, 15, 20, \dots, 120 \text{ GeV}$, every 5 GeV step.
- ▶ decay branch ratios are the same as the 125 GeV SM Higgs boson.



The background MC samples:

- ▶ 2-fermion ($2f^l, 2f^h$) leptonic/bhabha/hadronic
- ▶ 4-fermion ($4f^l, 4f^{sl}, 4f^h$) leptonic/semi-lepton/hadronic
- ▶ SM Higgs, $Higgs_{125}$
- ▶ $\gamma\gamma$ backgrounds



analysis flow

01

a muon pair

$$\chi^2(M_{\mu^+\mu^-}, M_{\text{rec}}) = \frac{(M_{\mu^+\mu^-} - M_Z)^2}{\sigma_{M_{\mu^+\mu^-}}^2} + \frac{(M_{\text{rec}} - M_h)^2}{\sigma_{M_{\text{rec}}}^2}.$$

02

$M_Z \in [73, 120]$ GeV

03


$P_T^Z \in [10, 128 - 4 \times \frac{M_h}{10}]$ GeV

04

$|\cos\theta_{\text{mis}}| < 0.98$ when $M_h > 50$ GeV


05

Multi-Variate Analysis : angles




recoil mass distribution

- ▶ recoil mass distribution for different M_{S^0} .
- ▶ $(-, +) / (+, -)$ 900 fb^{-1} ,
 $(-, -) / (+, +)$ 100 fb^{-1}
 polarization @ 250 GeV



mass region	main backgrounds	$M_{\text{rec}} \text{ (GeV/c}^2\text{)}$
$125 > M_{S^0} > M_Z$	$4f_{zz}^{sl}$, $4f_{zz/ww}$, SM Higgs	
$M_{S^0} \sim M_Z$	$4f_{zz}^l$, $4f_{zz}^{sl}$, $4f_{zz/ww}$, SM Higgs	
$M_Z > M_{S^0} > 40$	$2f_l$, $4f_{zz}$, $4f_{zz/ww}$	
$40 > M_{S^0}$	$2f_l$	


Comparing with MC results




- ▶ 95% CL upper bounds on scale factor of cross section with likelihood methods
- ▶ MC results with the same cuts.
- ▶ slightly different in the low mass region.




$2f_l$ process



$M_{S^0} = 15$ GeV process




$M_{S^0} = 50$ GeV process



- ▶ There is photon return effects in $2f_l$ process.
- ▶ identify ISR photon by
 - ▶ ISR photon in the central region ($\cos\theta < 0.95$): $E_{\text{central}} > 100$ GeV
 - ▶ ISR photon in the forward region ($0.95 < \cos\theta < 0.99$): $E_{\text{forward}} > 60$ GeV
 - ▶ ISR cone around photon axis: $\cos\alpha = 0.90$
 - ▶ Energy ratio inside the ISR photon cone: $\frac{E}{E_{\text{cone}}} = 0.95$

Comparing w/o photon veto results



- ▶ remove large amount of $2f_l$ background in small mass region.
- ▶ $S^{95} \in (0.0025 - 0.015) \rightarrow \kappa_{hZZ}^{95} \in (0.05-0.12)$.

OPAL's strategy


- ▶ at least two opposite charged leptons
- ▶ isolation of lepton tracks, $\alpha_{iso}^1 > 15^\circ$, $\alpha_{iso}^2 > 10^\circ$
- ▶ find two best leptons $m_{ll} \sim m_Z$
- ▶ invariant mass of the lepton pair, $M_{\mu\mu} \in [81.2, 101.2] \text{ GeV}$
- ▶ $p_{ll}^Z < 50 \text{ GeV}$
- ▶ polar angle of missing momentum, $|\theta_{mis}| < 0.95$ for $p_{mis} > 5 \text{ GeV}$
- ▶ acoplanarity
- ▶ ISR photon veto

my strategy

- ▶ at least two isolated muon, with IsolatedLeptonTagging Processor
- ▶ find two best leptons, $m_{ll} \sim m_Z$ and $m_{rec} \sim m_h$
- ▶ Recovery of bremsstrahlung and FSR photons
- ▶ Reconstruct Z boson mass $M_{\mu\mu} \in [73, 120] \text{ GeV}$.
- ▶ $70 \text{ GeV} > P_T^Z > 10 \text{ GeV}$
- ▶ the polar angle of the missing momentum, $|\theta_{mis}| < 0.98$
- ▶ MVA: $M_{\mu+\mu-}$, $\cos(\theta_Z)$, $\cos(\theta_{\mu+\mu-})$, $\cos(\theta_{\mu+})$, $\cos(\theta_{\mu-})$, acoplanarity
- ▶ ISR photon veto



Comparing with LEP results



- ▶ LEP recoil: LEP2 data from 161 GeV to 202 GeV, combined LEP1 data.
- ▶ LEP traditional: exclusive reconstruction of Z and h decay, mainly $h \rightarrow bb$, $h \rightarrow \tau\tau$.
- ▶ when $100 \geq M_h \geq 50$ GeV, trend are similar with LEP.
- ▶ generally, 1 order better than LEP recoil results

- ▶ A lighter higgs is favored in many BSM models
 - ▶ 2HDM, NMSSM, RS ...
- ▶ A model-independent analysis has been performed.
 - ▶ mass range [10, 120) GeV
 - ▶ 2000 fb^{-1} , when $\sqrt{s} = 250 \text{ GeV}$.
 - ▶ $(-+, +-, --, ++)$ = (45%, 45%, 5%, 5%) polarization scenario
- ▶ Exclusion limits for κ_{hZZ}^{95} coupling factor
 - ▶ $\kappa_{hZZ}^{95} \in (0.05-0.12)$.
 - ▶ 1 order better than LEP recoil results.







The higgs boson found at 2012: the SM Higgs?

Many BSMs predict one or more extra scalars:

- ▶ General Two Higgs Doublet Model (2HDM...)
 - with 2 scalars: h, H , 1 pseudoscalar A , 2 charged particles
- ▶ Next-to-Minimal Supersymmetric Standard Model (NMSSM)
 - with 3 scalars: h_1, h_2, h_3 , 2 pseudoscalars A_1, A_2 , 2 charged particles
- ▶ Randall Sundrum model
 - a radion

In these models, a scalar lighter than 125 GeV is well motivated.

LHC Higgs boson rather SM-like \rightarrow new higgs coupling to Z boson strongly suppressed.
Could we find it at the ILC?


Past Experiment Results parameters

LEP SM Higgs searches: constrain other extra scalars, whose properties, especially decay profile, are similar as SM higgs's.

LEP/LHC constrain rely on the model details: CP, mass hierarchy, couplings, etc.

JHEP 12 (2016) 068

survived after indirect + LEP + LHC constrains



2HDM, Type I:


$\tan\beta > 1.2$,

$m_A > 60$ GeV,

$m_{H^\pm} > 80$ GeV ..

Four regions.

mass region	main backgrounds
$125 > M_{S^0} > M_Z$	$4f_{zz}^{sl}$, $4f_{zz/ww}$, SM Higgs
$M_{S^0} \sim M_Z$	$4f_{zz}^{fl}$, $4f_{zz}^{sl}$, $4f_{zz/ww}$, SM Higgs
$M_Z > M_{S^0} > 40$	$2f_l$, $4f_{zz}$, $4f_{zz/ww}$
$40 > M_{S^0}$	$2f_l$



Cut efficiencies for different masses:

$\int L dt = 2000 fb^{-1}$	new higgs	$4f_l$	$4f_{sl}$	$2f_l$	total bk	cut efficiency	significance
$m_h = 115$ GeV	17419.6	61033.9	53869.4	13877.7	128781	0.67	45.56
$m_h = 90$ GeV	22198.2	63210.7	74563	18514.2	156288	0.59	52.54
$m_h = 70$ GeV	26841.3	51671.6	60357.7	37166.6	149196	0.57	63.97
$m_h = 50$ GeV	30493.5	46128.1	54372.8	80074.4	180575	0.54	66.37
$m_h = 30$ GeV	33843.7	51206.6	55743.3	213184	320134	0.49	56.88

$$\text{significance} = \frac{S}{\sqrt{S+B}}, \quad \text{and } S = \kappa_{gZZ}^2 \times \sigma_{h\mu\mu}^{m_h} \times \mathcal{L}, \quad \text{where } \kappa_{gZZ} = 1$$



- ▶ 2σ exclusion limits with a bin-by-bin comparison between the signal and backgrounds recoil mass histograms.
- ▶ the background-only hypothesis — no new higgs in the investigated mass range.
- ▶ the signal-plus-background hypothesis — the new higgs is assumed to be produced.
- ▶ a global test-statistic $X(m_h) = \mathcal{L}(s(m_h))/\mathcal{L}(0)$ is constructed to discriminate signal and background.
- ▶ the distributions of $X(m_h)$ are normalised to become probability density functions → integrated to be the confidence levels $CL_b(m_h)$ and $CL_{s+b}(m_h)$.
- ▶ the ratio $CL_s(m_h) = CL_{s+b}(m_h)/CL_b(m_h)$ is used to describe that the signal confidence one might have obtained in the absence of background.

