


Measuring the CP state of tau lepton pairs from Higgs decay at the ILC



arXiv:1804.01241
submitted to PRD

Daniel Jeans, KEK
ALCW18, Fukuoka



Motivation

Is the 125 GeV Higgs a CP eigenstate ?

$$h_{125} = \cos \psi_{CP} h^{CP\text{even}} + \sin \psi_{CP} A^{CP\text{odd}}$$

pure CP even: $\psi_{CP} = 0$ [Standard Model]

odd: $\psi_{CP} = \pi/2$ [excluded at LHC]

or a mixture?

Do Higgs couplings conserve CP ?

e.g. coupling to fermions: $\mathcal{L} \sim g \bar{f} (\cos \psi_{CP} + i \gamma^5 \sin \psi_{CP}) f H$

CP conserving coupling

$\psi_{CP} = 0$ [Standard Model]


maximally violating

$\psi_{CP} = \pi/2$


or partially violating ?

$$h_{125} = \cos \psi_{CP} h^{CP\text{even}} + \sin \psi_{CP} A^{CP\text{odd}}$$

$$g \bar{f} (\cos \psi'_{CP} + i \gamma^5 \sin \psi'_{CP}) f h_{125}$$





The **correlation** between spins of Higgs decay products
is sensitive to their **CP state** [in particular, the transverse correlation]



why use **tau leptons** to measure CP in Higgs sector?

- **fermion**: tree-level CP effects possible
($H \rightarrow WW, ZZ$ only via loops)
- **unstable** fermion:
distribution of tau decay products gives
access to tau spin direction
optimal estimator = “polarimeter vector”
easy to extract for $\tau^+ \rightarrow (\pi^+ \nu)$ and $\tau^+ \rightarrow (\pi^+ \pi^0 \nu)$ decay modes
- reasonable 6% **branching ratio**
- clean separation of the two fermion decays (no colour string as in $H \rightarrow b\bar{b}$)



distribution of $\Delta\phi$ is sensitive to CP mixing angle ψ_{CP}

amplitude of modulation in $\Delta\phi$ varies from event to event, depending on θ^\pm , according to contrast function:

$$c(\theta^+, \theta^-) \equiv \sin \theta^+ \sin \theta^- / (1 + \cos \theta^+ \cos \theta^-)$$




FIG. 3. Two-dimensional distribution of events in $\Delta\phi$ and $c(\theta^+, \theta^-)$ at MC truth level, for the case $\psi_{CP} = 0$.

In this analysis, we measure ψ_{CP} of the **tau pair** from **Higgs** decay in a model-independent way, by measuring the phase of the $\Delta\varphi$ distribution

we don't try to understand which mechanism creates the mixing:
explicitly CP violating coupling, mixed CP mass eigenstate, ...
→ requires combination with other measurements,
model assumptions, ...



Analysis uses full ILD simulation
signal and SM backgrounds
standard data reconstruction
(ilcsoft v01-16-02)

SIGNAL: $e^+ e^- \rightarrow Z H$

$Z \rightarrow$ electrons, muons, quarks

$H \rightarrow \tau^+ \tau^-$

$\tau^\pm \rightarrow (\pi^\pm \nu)$ or $(\pi^\pm \pi^0 \nu)$



SM backgrounds:

$e^+ e^- \rightarrow ff H, 4f, 2f$

assume 2 ab^{-1} of 250 GeV data: “H20-staged”

Full tau reconstruction

NIM A810 (2016) 51

[arXiv:1507.01700](https://arxiv.org/abs/1507.01700)

to reconstruct tau **polarimeter**, need
full reconstruction of tau decay products,
including the neutrino(s)

in hadronic tau decays (# neutrino = 1), if we know
the tau **production vertex**,
the **impact parameters** of charged tau decay products,
the p_T of the tau-tau system,

then the neutrino momenta can be reconstructed:

6 **unknowns**/event:

2 x neutrino 3-momenta

6 **constraints**/event:

2 x impact parameter defines plane of tau momentum

2 x tau invariant mass

2 from event p_T [p_x , p_y] → insensitive to ISR / beamstrahlung

[+ solve two-fold ambiguities from quadratic constraints using tau lifetime,
and, only if necessary, using reconstructed tau-tau mass]

vertex detector
tracking
photon reco.
Jet En. Res.

reconstruct $Z \rightarrow e e, \mu \mu, \text{jets} + 2 \times (1\text{-prong tau jets})$

simple preselection

some distributions after reconstruction and preselection:







TABLE II. Selection cuts [see text for details; (energies, momenta, and masses) in GeV/c^(0,1,2)], signal selection efficiencies ϵ (in %), and number of expected background events (BG) at various stages of the selection in the three selection channels e, μ, q . Event numbers are scaled to the 2 ab⁻¹ of 250 GeV data of the “H20-staged” running scenario.

event property	leptonic preselection			hadronic preselection				
	requirement	ϵ_e	ϵ_μ	BG _{lep}	requirement	ϵ_q	BG _{had}	
		100	100	142 M		100	142 M	
chg. PFOs	4 \rightarrow 7	91	93	10.1 M	≥ 8	98	95.7 M	
$Z \rightarrow ll$ candidate	≥ 1	88	90	1.03 M				
isolated prongs					≥ 2	91	45.8 M	
opp. chgd. prongs		84	87	903 k		84	33.5 M	
min. prong score					> 0.8	77	14.5 M	
impact par. error	$< 25\mu m$	76	79	491 k	$< 25\mu m$	74	13.2 M	
extra cone energy		72	75	438 k				
m_Z					60 \rightarrow 160	72	5.58 M	
m_{recoil}					50 \rightarrow 160	71	4.90 M	
τ decay mode		63	65	236 k		64	1.99 M	
full selection		$Z \rightarrow ee$		$Z \rightarrow \mu\mu$		$Z \rightarrow qq$		
event property	requirement	ϵ_e	BG _e	ϵ_μ	BG _{μ}	requirement	ϵ_q	BG _q
good $\tau^+\tau^-$ fit		57	112 k	59	99.5 k		58	1.64 M
$m_{\tau\tau}$	100 \rightarrow 140	46	618	52	366	100 \rightarrow 140	42	43.5 k
event p_T	< 5	43	309	50	268	< 20	42	31.6 k
m_{recoil}	> 120	42	252	50	162	> 100	41	23.5 k
m_Z	80 \rightarrow 105	41	186	49	136	80 \rightarrow 115	38	6.93 k
$ \cos\theta_Z $	< 0.96	40	168	47	124	< 0.96	37	6.22 k
event p_z	< 40	40	144	47	105	< 40	37	5.26 k
$ \cos\theta_P _{\text{min}}$	< 0.95	40	140	47	102	< 0.95	37	5.26 k
Sample purity (%)		19		26		11		

TABLE I. Migrations among τ -pair decay modes, for preselected and reconstructed signal events in which the Z boson decays to either muons or light quarks. All numbers are given in %.





Reco. decay	True decay		
	$(\pi\nu, \pi\nu)$	$(\pi\nu, \rho\nu)$	$(\rho\nu, \rho\nu)$
$Z \rightarrow \mu^+\mu^-$			
$(\pi\nu, \pi\nu)$	93	3	< 1
$(\pi\nu, \rho\nu)$	7	93	6
$(\rho\nu, \rho\nu)$	< 1	4	94
$Z \rightarrow qq(\text{uds})$			
$(\pi\nu, \pi\nu)$	89	6	< 1
$(\pi\nu, \rho\nu)$	11	89	12
$(\rho\nu, \rho\nu)$	< 1	5	87

reconstructing polarimeter vectors: in tau rest frame

$$\mathbf{h}(\tau^\pm \rightarrow \pi^\pm \nu) \propto \mathbf{p}_{\pi^\pm} \quad (6)$$



$$\begin{aligned} \mathbf{h}(\tau^\pm \rightarrow \pi^\pm \pi^0 \nu) \propto & m_\tau (E_{\pi^\pm} - E_{\pi^0}) (\mathbf{p}_{\pi^\pm} - \mathbf{p}_{\pi^0}) \\ & + 2(p_{\pi^\pm} + p_{\pi^0})^2 \mathbf{p}_\nu, \quad (7) \end{aligned}$$

some distributions after selection



group events according to expected sensitivity, based on:





- longitudinal component of polarimeters (contrast) → intrinsic sensitivity
- tau decay prongs' d0 measurement significance → reconstruction quality
- output of simple NN [6 inputs] (signal vs. main 4f bgs) → bg. contamination
- output of simple NN [4 inputs] (signal tau decays vs. others) → tau decay mis-identification



CP sensitive observable $\Delta\phi$ in different event sensitivity bins

signal background

error bars:
MC statistics







phase of **signal distributions** is sensitive to CP

estimating measurement sensitivity

unbinned maximum likelihood fit: simultaneously in all sensitivity bins and selection channels
fit a single parameter: the phase of $\Delta\phi$ distribution

perform series of toy pseudo-experiments using simulated distributions

results of 10k pseudo-exps




predicted sensitivity on ψ_{CP} under various conditions

TABLE IV. Estimated experimental precision $\delta\psi_{CP}$ on the CP phase in different scenarios.

$\int \mathcal{L}$ [ab ⁻¹]	beam pol.		notes	$\delta\psi_{CP}$ [mrad]
	e^-	e^+		
1.0	0	0	full analysis	116
1.0	0	0	only $Z \rightarrow ee$	450
1.0	0	0	only $Z \rightarrow \mu\mu$	412
1.0	0	0	only $Z \rightarrow qq$	122
1.0	0	0	only $(\pi\nu, \pi\nu)$	387
1.0	0	0	only $(\pi\nu, \rho\nu)$	198
1.0	0	0	only $(\rho\nu, \rho\nu)$	166
1.0	-1.0	+1.0	pure $e_L^- e_R^+$	97
1.0	+1.0	-1.0	pure $e_R^- e_L^+$	113
1.0	0	0	$\sigma_{ZH} + 20\%$	104
1.0	0	0	$\sigma_{ZH} - 20\%$	133
1.0	0	0	no bg.	76
1.0	0	0	perf. pol.	100
1.0	0	0	no bg., perf. pol./eff.	25
H20-staged: 250 GeV, 2 ab ⁻¹				
0.9	-0.8	+0.3	only $e_L^- e_R^+$	102
0.9	+0.8	-0.3	only $e_R^- e_L^+$	120
0.1	-0.8	-0.3	only $e_L^- e_L^+$	359
0.1	+0.8	+0.3	only $e_R^- e_R^+$	396
2.0	mixed		full analysis	75

sanity check: output = input phase



summary

demonstrated, using full detector simulation and backgrounds,
that CP mixing in tau-pair from Higgs decays
can be determined to 75 mrad \sim 4.3 deg
using 2 ab⁻¹ of ILC250 data
[10~15 years from ILC start]

potential for improved analysis methods
to significantly improve results [ultimately \rightarrow 25 mrad]

using only a few tau decay modes
 \rightarrow can probably increase sensitivity by also using other modes

backup

International Linear Collider



electron-positron collisions

initial collision energy 250 GeV → Higgs factory


linear design allows relatively simple energy upgrade in the future


longitudinally polarised beams

Technical Design Report published 2012

undergoing detailed review in MEXT

expect project decision from JP government
in next few months





ILC Candidate Location: Kitakami Area



Kaname
Kodashima


“Lots of
encounters
through the ILC”

Futago
Elementary
School,
Kitakami City

Comment: “I drew my
wish that there would
be a lot of new
encounters through the
ILC everyone is hoping
for.”




International Large Detector



One of two detector designs being studied for the ILC
dominated by groups from Europe and Japan

Design principles

- excellent vertexing: identification of b , c , τ
 - high precision and lightweight vertex detector
- highly efficient and precise charged particle tracking
 - large TPC in ~ 3.5 T field
- excellent jet energy resolution
 - make best use of dominant hadronic decays of W , Z , H
- highly granular calorimeters



ECAL

Time
Projection
Chamber

1.6 → 1.85m

silicon strips

FCAL

vertex detector
silicon pixels

HCAL







FIG. 7. Distributions of the two Neural Network outputs in the muon and hadronic selection channels. The structure in the output of NN2 is due to the three different combinations of τ lepton decay modes. Distributions are normalized to 0.9 ab^{-1} of data in the $e_L^- e_R^+$ beam polarization.