



We still believe in supersymmetry

You must be joking



SUSY Higgs Bosons: Reaching the LC Precision

Sven Heinemeyer, IFT/IFCA (CSIC, Madrid/Santander)

Sendai, 10/2019

1. The Quest for Precision
2. SUSY Higgs boson masses
3. MSSM Higgs production cross sections
4. (N)MSSM Higgs decays
5. Conclusions

1. The Quest for Precision



⇒ clear discovery at $\sim 125 \text{ GeV}$!

⇒ can be interpreted as the light(/heavy) CP -even MSSM Higgs

The Higgs mass accuracy: experiment vs. theory:

Experiment:

ATLAS: $M_h^{\text{exp}} = 125.36 \pm 0.37 \pm 0.18 \text{ GeV}$

CMS: $M_h^{\text{exp}} = 125.03 \pm 0.27 \pm 0.15 \text{ GeV}$

combined: $M_h^{\text{exp}} = 125.09 \pm 0.21 \pm 0.11 \text{ GeV}$

The Higgs mass accuracy: experiment vs. theory:

Experiment:

$$\text{ATLAS:} \quad M_h^{\text{exp}} = 125.36 \pm 0.37 \pm 0.18 \text{ GeV}$$

$$\text{CMS:} \quad M_h^{\text{exp}} = 125.03 \pm 0.27 \pm 0.15 \text{ GeV}$$

$$\text{combined:} \quad M_h^{\text{exp}} = 125.09 \pm 0.21 \pm 0.11 \text{ GeV}$$

MSSM theory:

LHCHSWG adopted **FeynHiggs** for the prediction of MSSM Higgs boson masses and mixings (considered to be the code containing the most complete implementation of higher-order corrections)

$$\text{FeynHiggs:} \quad \delta M_h^{\text{theo}} \sim 3 \text{ GeV (now 1 – 2 GeV?)}$$

→ rough estimate, FeynHiggs contains algorithm to evaluate uncertainty, depending on parameter point

Katharsis of Ultimate Theory Standards

11th meeting: 20-22 November 2019 (MPI Munich)

Precise Calculation of

(N)




Higgs Boson masses

Local organizers: T. Hahn, W. Hollik

Organized by:
M. Carena, H. Haber
R. Harlander, S. Heinemeyer
W. Hollik, P. Slavich, G. Weiglein

The MSSM:

⇒ Superpartners for Standard Model particles



Enlarged Higgs sector: Two Higgs doublets

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1 + i\chi_1)/\sqrt{2} \\ \phi_1^- \end{pmatrix}$$
$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2 + i\chi_2)/\sqrt{2} \end{pmatrix}$$

$$V = m_1^2 H_1 \bar{H}_1 + m_2^2 H_2 \bar{H}_2 - m_{12}^2 (\epsilon_{ab} H_1^a H_2^b + \text{h.c.})$$
$$+ \underbrace{\frac{g'^2 + g^2}{8}}_{\text{gauge couplings, in contrast to SM}} (H_1 \bar{H}_1 - H_2 \bar{H}_2)^2 + \underbrace{\frac{g^2}{2}}_{\text{gauge couplings, in contrast to SM}} |H_1 \bar{H}_2|^2$$

physical states: h^0, H^0, A^0, H^\pm

Goldstone bosons: G^0, G^\pm

Input parameters: (to be determined experimentally)

$$\tan \beta = \frac{v_2}{v_1}, \quad M_A^2 = -m_{12}^2 (\tan \beta + \cot \beta)$$

Enlarged Higgs sector: Two Higgs doublets with \mathcal{CP} violation

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1 + i\chi_1)/\sqrt{2} \\ \phi_1^- \end{pmatrix}$$

$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2 + i\chi_2)/\sqrt{2} \end{pmatrix} e^{i\xi}$$

$$V = m_1^2 H_1 \bar{H}_1 + m_2^2 H_2 \bar{H}_2 - m_{12}^2 (\epsilon_{ab} H_1^a H_2^b + \text{h.c.})$$

$$+ \underbrace{\frac{g'^2 + g^2}{8}}_{\text{gauge couplings, in contrast to SM}} (H_1 \bar{H}_1 - H_2 \bar{H}_2)^2 + \underbrace{\frac{g^2}{2}}_{\text{gauge couplings, in contrast to SM}} |H_1 \bar{H}_2|^2$$

physical states: h^0, H^0, A^0, H^\pm

2 \mathcal{CP} -violating phases: $\xi, \arg(m_{12}) \Rightarrow$ can be set/rotated to zero

Input parameters: (to be determined experimentally)

$$\tan \beta = \frac{v_2}{v_1}, \quad M_{H^\pm}^2$$

Z_3 invariant NMSSM

MSSM Higgs sector: Two Higgs doublets

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1 + i\chi_1)/\sqrt{2} \\ \phi_1^- \end{pmatrix}$$
$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2 + i\chi_2)/\sqrt{2} \end{pmatrix}$$

$$V = (\tilde{m}_1^2 + |\mu|^2)H_1\bar{H}_1 + (\tilde{m}_2^2 + |\mu|^2)H_2\bar{H}_2 - m_{12}^2(\epsilon_{ab}H_1^aH_2^b + \text{h.c.})$$
$$+ \frac{g'^2 + g^2}{8}(H_1\bar{H}_1 - H_2\bar{H}_2)^2 + \frac{g^2}{2}|H_1\bar{H}_2|^2$$

Z_3 invariant NMSSM

NMSSM Higgs sector: Two Higgs doublets + one Higgs singlet

$$H_1 = \begin{pmatrix} H_1^1 \\ H_1^2 \end{pmatrix} = \begin{pmatrix} v_1 + (\phi_1 + i\chi_1)/\sqrt{2} \\ \phi_1^- \end{pmatrix}$$

$$H_2 = \begin{pmatrix} H_2^1 \\ H_2^2 \end{pmatrix} = \begin{pmatrix} \phi_2^+ \\ v_2 + (\phi_2 + i\chi_2)/\sqrt{2} \end{pmatrix}$$

$$S = v_s + S_R + IS_I$$

$$\begin{aligned} V = & (\tilde{m}_1^2 + |\mu\lambda S|^2)H_1\bar{H}_1 + (\tilde{m}_2^2 + |\mu\lambda S|^2)H_2\bar{H}_2 - m_{12}^2(\epsilon_{ab}H_1^a H_2^b + \text{h.c.}) \\ & + \frac{g'^2 + g^2}{8}(H_1\bar{H}_1 - H_2\bar{H}_2)^2 + \frac{g^2}{2}|H_1\bar{H}_2|^2 \\ & + |\lambda(\epsilon_{ab}H_1^a H_2^b) + \kappa S^2|^2 + m_S^2|S|^2 + (\lambda A_\lambda(\epsilon_{ab}H_1^a H_2^b)S + \frac{\kappa}{3}A_\kappa S^3 + \text{h.c.}) \end{aligned}$$

Free parameters:

$$\lambda, \kappa, A_\kappa, M_{H^\pm}, \tan\beta, \mu_{\text{eff}} = \lambda v_s$$

Higgs spectrum:

\mathcal{CP} -even : h_1, h_2, h_3

\mathcal{CP} -odd : a_1, a_2

charged : H^+, H^-

Goldstones : G^0, G^+, G^-

Neutralinos:

$$\mu \rightarrow \mu_{\text{eff}}$$

compared to the MSSM: one singlino more

$$\rightarrow \tilde{\chi}_1^0, \tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0, \tilde{\chi}_5^0$$

Mass of the lightest \mathcal{CP} -even Higgs:

$$m_{h,\text{tree,NMSSM}}^2 = m_{h,\text{tree,MSSM}}^2 + M_Z^2 \frac{\lambda^2}{g^2} \sin^2 2\beta$$

Mass of the \mathcal{CP} -odd Higgs:

$$\text{MSSM} : M_A^2 = -m_{12}^2 (\tan \beta + \cot \beta) = \mu B (\tan \beta + \cot \beta)$$

$$\text{NMSSM} : "M_A^2" = \mu_{\text{eff}} B_{\text{eff}} (\tan \beta + \cot \beta)$$

$$\text{with } B_{\text{eff}} = A_\lambda + \kappa s, \mu_{\text{eff}} = \lambda s \quad \Rightarrow \text{one very light } a_1$$

Mass of the charged Higgs:

$$\text{MSSM} : M_{H^\pm}^2 = M_A^2 + M_W^2 = M_A^2 + \frac{1}{2} v^2 g^2$$

$$\text{NMSSM} : M_{H^\pm}^2 = M_A^2 + v^2 \left(\frac{g^2}{2} - \lambda^2 \right)$$

Mass of the lightest \mathcal{CP} -even Higgs:

$$m_{h,\text{tree,NMSSM}}^2 = m_{h,\text{tree,MSSM}}^2 + M_Z^2 \frac{\lambda^2}{g^2} \sin^2 2\beta$$

Mass of the \mathcal{CP} -odd Higgs:

$$\text{MSSM} : M_A^2 = -m_{12}^2 (\tan \beta + \cot \beta) = \mu B (\tan \beta + \cot \beta)$$

$$\text{NMSSM} : "M_A^2" = \mu_{\text{eff}} B_{\text{eff}} (\tan \beta + \cot \beta)$$

$$\text{with } B_{\text{eff}} = A_\lambda + \kappa s, \mu_{\text{eff}} = \lambda s \quad \Rightarrow \text{one very light } a_1$$

Mass of the charged Higgs:

$$\text{MSSM} : M_{H^\pm}^2 = M_A^2 + M_W^2 = M_A^2 + \frac{1}{2} v^2 g^2$$

$$\text{NMSSM} : M_{H^\pm}^2 = M_A^2 + v^2 \left(\frac{g^2}{2} - \lambda^2 \right)$$

$$\Rightarrow M_{h_1}^{\text{MSSM,tree}} \leq M_{h_1}^{\text{NMSSM,tree}}, \text{ one light } a_1, M_{H^\pm}^{\text{MSSM,tree}} \geq M_{H^\pm}^{\text{NMSSM,tree}}$$

Higgs coupling determination at e^+e^- collider:

recoil method: $e^+e^- \rightarrow ZH, Z \rightarrow e^+e^-, \mu^+\mu^-$

⇒ total measurement of Higgs production cross section

⇒ **NO** additional theoretical assumptions needed for absolute determination of partial widths

⇒ all observable channels can be measured with high accuracy

⇒ Cross section needed with high precision, better than $\sim 1\%$

Available: SM cross section predictions at the 1% accuracy level


⇒ improvements necessary ... full 2-loop calculations and more ... ?!

⇒ What about the MSSM cross sections?

HL-LHC vs. ILC in the most general κ framework:


[P. Bechtle, S.H., O. Stål, T. Stefaniak, G. Weiglein '14]

no theory assumptions, full fit



\Rightarrow strong improvement with the ILC

Future expectations for κ (kappa-3 framework)



⇒ very roughly similar results

⇒ FCC-hh/-he/-ee appears better

⇒ FCC-hh uses different theory assumptions, uncertainties $\lesssim 1\%$

⇒ also remember different time scales!

Needed for LHC/ILC/CLIC/... physics:

Precise and consistent prediction of

- Higgs boson masses
- Higgs boson mixings
- Higgs boson couplings
- Higgs boson production cross sections
- Higgs boson decay widths/branching ratios
- ...

⇒ (partially) provided by FeynHiggs

2. SUSY Higgs mass calculations

The “easy” case: MSSM:

Propagator/Mass matrix at tree-level:

$$\begin{pmatrix} q^2 - m_H^2 & 0 \\ 0 & q^2 - m_h^2 \end{pmatrix}$$

Propagator / mass matrix with higher-order corrections
(→ Feynman-diagrammatic approach):

$$M_{hH}^2(q^2) = \begin{pmatrix} q^2 - m_H^2 + \widehat{\Sigma}_{HH}(q^2) & \widehat{\Sigma}_{Hh}(q^2) \\ \widehat{\Sigma}_{hH}(q^2) & q^2 - m_h^2 + \widehat{\Sigma}_{hh}(q^2) \end{pmatrix}$$

$\widehat{\Sigma}_{ij}(q^2)$ ($i, j = h, H$) : renormalized Higgs self-energies

\mathcal{CP} -even fields can mix


⇒ complex roots of $\det(M_{hH}^2(q^2))$: $\mathcal{M}_{h_i}^2$ ($i = 1, 2$): $\mathcal{M}^2 = M^2 - iM\Gamma$

Calculation of renormalized Higgs boson self-energies:

$$\hat{\Sigma}(q^2) = \hat{\Sigma}^{(1)}(q^2) + \hat{\Sigma}^{(2)}(q^2) + \dots + \text{log resum}$$

Main contribution: t/\tilde{t} sector (\tilde{t} : scalar top, SUSY partner of the t)

Very leading 1-Loop:



Structure of higher-order corrections at one-loop:

$$\Delta M_h^2 \sim m_t^2 \alpha_t [L + L^0] , \quad L := \log \left(\frac{m_{\tilde{t}}}{m_t} \right)$$

Large $m_{\tilde{t}}$ \Rightarrow large L \Rightarrow resummation of logs necessary

Codes on the market:

1.) Fixed order codes: good for all scales low

- SuSpect
- SPheno/SARAH
- SoftSUSY/FlexibleSUSY
- H3m

2.) EFT codes (pure log resum): good for all scales high

- SusyHD
- MhEFT
- HSSUSY

3.) Hybrid codes: good always?!

- FeynHiggs
- FlexibleEFTHiggs
- SPheno/SARAH

Obviously: quality depends on the details implemented

Codes on the market:

1.) Fixed order codes: good for all scales low

- SuSpect
- SPheno/SARAH
- SoftSUSY/FlexibleSUSY
- H3m

2.) EFT codes (pure log resum): good for all scales high

- SusyHD
- MhEFT
- HSSUSY

3.) Hybrid codes: good always?!

- FeynHiggs \leftarrow our code / best code :-)
- FlexibleEFTHiggs
- SPheno/SARAH

Obviously: quality depends on the details implemented

Possible & necessary refinements of the EFT calculation:

- Inclusion of EWino mass scale in RGE's
- Inclusion of gluino mass scale in RGE's
- Inclusion of EW effects in RGE's
- Inclusion of 3-loop RGEs plus 2-loop thresholds etc.
- “Two Higgs Doublet Model” below M_S
- Splitting in the scalar top sector
- . . .


Possible & necessary refinements of the EFT calculation:

- Inclusion of EWino mass scale in RGE's
⇒ included into FeynHiggs
- Inclusion of gluino mass scale in RGE's
⇒ included into FeynHiggs
- Inclusion of EW effects in RGE's
⇒ included into FeynHiggs
- Inclusion of 3-loop RGEs plus 2-loop thresholds etc.
⇒ included into FeynHiggs
- “Two Higgs Doublet Model” below M_S
⇒ private version of FeynHiggs exists, other code: MhEFT
- Splitting in the scalar top sector
⇒ future work
- ...

Impact of precise M_h calculation (I):

Impact of non-degenerate $\mathcal{O}(\alpha_t^2)$ threshold corr. in EFT part:

One scale M_{SUSY} , but large stop sector splitting, $\tan\beta = 10$:




\Rightarrow important for large X_t (more in a moment)

Impact of precise M_h calculation (II):

Impact of pole mass determination improvements:
(ask me details over coffee!)


One scale M_{SUSY} , $\tan \beta = 10$:



⇒ calculation stabilized!

Improved uncertainty estimate:

[H. Bahl, S.H., W. Hollik, G. Weiglein '19 – PRELIMINARY]



⇒ reduced to 1-2 GeV (in simple single-scale scenario)

Going beyond the MSSM: the FeynHiggs Ansatz

(taken from talk by [P. Drechsel])

General idea: treat the MSSM part exactly as in the MSSM

- ▶ full inverse propagator in CP-even sector for mass determination

$$\Delta^{-1}(k^2) = i \left[k^2 \mathbb{1} - \underbrace{\mathcal{M}_{\phi\phi} + \hat{\Sigma}_{\phi\phi}^{(1L)}(k^2)}_{\text{NMSSM}} + \underbrace{\hat{\Sigma}_{\phi\phi}^{(2L)}(k^2 = 0)}_{\text{MSSM/FEYNHIGGS}} \right]$$


- ▶ included corrections from FEYNHIGGS at 2-loop order:
 - ▶ orders $\mathcal{O}(\alpha_s \alpha_t, \alpha_s \alpha_b, \alpha_t^2, \alpha_t \alpha_b)$
 - ▶ resummed large logarithms

⇒ any deviation from the MSSM can directly attributed to the extended model!


⇒ kind of obvious, but only FeynHiggs does it ...

Lightest Mass @ 1-Loop Order: Sample Scenario

Absolute difference between different mass predictions



$$\Delta M = \left| m_{h_1}^{(Y_t)} - m_{h_1}^{(Y_t, \lambda)} \right|$$



$$\Delta M = \left| m_{h_1}^{(Y_t, \lambda+H+G)} - m_{h_1}^{(1L)} \right|$$

include Higgs- & gauge-sector

⇒ influence of corrections beyond top/scalar top-sector is by far larger than those of the order $\mathcal{O}(Y_t \lambda, \lambda^2)$

⇒ we need two-loop calculations rather from the Higgs/gauge sector than from the genuine NMSSM t/\tilde{t} sector!

⇒ those are not available and more complicated

⇒ NMSSM has intrinsically larger uncertainties than the MSSM

3. MSSM Higgs production cross sections

Neutral Higgs production:

$$e^+e^- \rightarrow h_i Z, h_i \gamma, h_i h_j, h_i \nu \bar{\nu}, h_i e^+ e^-, h_i t \bar{t}, h_i b \bar{b}, \dots \quad (i, j = 1, 2, 3).$$

Now available in the **cMSSM** at the full one-loop level:

[S.H., C. Schappacher '15] [F. Arco, S.H., C. Schappacher '18]

$$\sigma(e^+e^- \rightarrow h_i h_j)$$

$$\sigma(e^+e^- \rightarrow h_i Z)$$

$$\sigma(e^+e^- \rightarrow h_i \gamma)$$

In the following:

few examples of each process, relevance of loop corrections

cMSSM parameters:

Table 2: MSSM default parameters for the numerical investigation; all parameters (except of t_β) are in GeV (calculated masses are rounded to 1 MeV). The values for the trilinear sfermion Higgs couplings, $A_{t,b,\tau}$ are chosen such that charge- and/or color-breaking minima are avoided [76], and $A_{b,\tau}$ are chosen to be real. It should be noted that for the first and second generation of sfermions we chose instead $A_f = 0$, $M_{\tilde{Q},\tilde{U},\tilde{D}} = 1500$ GeV and $M_{\tilde{L},\tilde{E}} = 500$ GeV.



Scen.	\sqrt{s}	t_β	μ	M_{H^\pm}	$M_{\tilde{Q},\tilde{U},\tilde{D}}$	$M_{\tilde{L},\tilde{E}}$	$ A_{t,b,\tau} $	M_1	M_2	M_3
S	1000	7	200	300	1000	500	$1500 + \mu/t_\beta$	100	200	1500

m_{h_1}	m_{h_2}	m_{h_3}
123.404	288.762	290.588


with \sqrt{s} , M_{H^\pm} , $\tan \beta$, ϕ_{A_t} varied

- Scenario chosen such that many processes are possible at the same time
- not chosen to maximize loop corrections

$$\underline{e^+e^- \rightarrow h_i h_j}$$




$e^+e^- \rightarrow h_1h_2$:




\Rightarrow loop corrections crucial!

$e^+e^- \rightarrow h_1h_2$:





\Rightarrow phase dependence more pronounced at loop-level

$e^+e^- \rightarrow h_1 h_1$ (purely loop induced):




\Rightarrow possibly observable!

$e^+e^- \rightarrow h_i Z$:




$e^+e^- \rightarrow h_1 Z$:




\Rightarrow loop corrections crucial

$e^+e^- \rightarrow h_3Z$:




\Rightarrow possibly observable, loop corrections crucial

$e^+e^- \rightarrow h_3Z$:




\Rightarrow pronounced phase dependence at the loop level


$e^+e^- \rightarrow h_i\gamma$: purely loop-induced!



$e^+e^- \rightarrow h_1\gamma$ (purely loop induced):




\Rightarrow possibly observable!



⇒ relevant variation with scalar top sector

$e^+e^- \rightarrow h_1\gamma$ (purely loop induced):

[F. Arco, S.H., C. Schappacher '18]



⇒ relevant variation with chargino sector

Charged MSSM Higgs Production at the LC

Charged Higgs production:

$$e^+e^- \rightarrow H^+H^-, H^\pm W^\mp, H^\pm e^\mp \nu, H^\pm tb, \dots$$

Now available in the **cMSSM** at the full one-loop level:

[S.H., C. Schappacher '17]

$$\sigma(e^+e^- \rightarrow H^+H^-)$$


$$\sigma(e^+e^- \rightarrow H^\pm W^\mp)$$

In the following:

few examples of each process, relevance of loop corrections

⇒ **BACKUP**

4. Higgs boson decays in the (N)MSSM



The FeynHiggs Ansatz for masses (taken from talk by [P. Drechsel])

General idea: treat the MSSM part exactly as in the MSSM

- ▶ full inverse propagator in CP-even sector for mass determination

$$\Delta^{-1}(k^2) = i \left[k^2 \mathbb{1} - \underbrace{\mathcal{M}_{\phi\phi} + \hat{\Sigma}_{\phi\phi}^{(1L)}(k^2)}_{\text{NMSSM}} + \underbrace{\hat{\Sigma}_{\phi\phi}^{(2L)}(k^2 = 0)}_{\text{MSSM/FEYNHIGGS}} \right]$$

- ▶ included corrections from FEYNHIGGS at 2-loop order:
 - ▶ orders $\mathcal{O}(\alpha_s \alpha_t, \alpha_s \alpha_b, \alpha_t^2, \alpha_t \alpha_b)$
 - ▶ resummed large logarithms

⇒ any deviation from the MSSM can directly attributed to the extended model!

⇒ kind of obvious, but only FeynHiggs does it ...

The FeynHiggs Ansatz for masses (taken from talk by [P. Drechsel])

General idea: treat the MSSM part exactly as in the MSSM

- ▶ full inverse propagator in CP-even sector for mass determination

$$\Delta^{-1}(k^2) = i \left[k^2 \mathbb{1} - \underbrace{\mathcal{M}_{\phi\phi} + \hat{\Sigma}_{\phi\phi}^{(1L)}(k^2)}_{\text{NMSSM}} + \underbrace{\hat{\Sigma}_{\phi\phi}^{(2L)}(k^2 = 0)}_{\text{MSSM/FEYNHIGGS}} \right]$$

- ▶ included corrections from FEYNHIGGS at 2-loop order:
 - ▶ orders $\mathcal{O}(\alpha_s \alpha_t, \alpha_s \alpha_b, \alpha_t^2, \alpha_t \alpha_b)$
 - ▶ resummed large logarithms

⇒ any deviation from the MSSM can directly attributed to the extended model!

⇒ kind of obvious, but only FeynHiggs does it ...

⇒ **same Ansatz for Higgs decays!**

What is included in FeynHiggs (so far):


Evaluation of all MSSM Higgs boson masses and mixing angles

- $M_{h_1}, M_{h_2}, M_{h_3}, M_{H^\pm}$, α_{eff} , \mathbf{Z}_{ij} , \mathbf{U}_{ij} , ... \Rightarrow precision disussed before

Evaluation of all neutral MSSM Higgs boson decay channels (so far)

- total decay width Γ_{tot}
- $\text{BR}(h_i \rightarrow f\bar{f})$: decay to SM fermions: full 1L, running m_q at 3L, \mathbf{Z}_{ij}
- $\text{BR}(h_i \rightarrow Z^{(*)}Z^{(*)}, W^{(*)}W^{(*)})$: decay to massive SM gauge bosons: Prophecy4f \oplus coupling factors, \mathbf{U}_{ij}
- $\text{BR}(h_i \rightarrow \gamma\gamma, gg)$: decay to massless SM gauge bosons: NLO QCD, gg : NNLO, NNLL from SM, \mathbf{U}_{ij}
- $\text{BR}(h_i \rightarrow h_j Z^{(*)}, h_j h_k)$: decay to gauge and Higgs bosons: $h_j Z^{(*)}$: \mathbf{U}_{ij} , $h_j h_k$: full 1L, log-resum, \mathbf{Z}_{ij}
- $\text{BR}(h_i \rightarrow \tilde{f}_i \tilde{f}_j)$: decay to sfermions: \mathbf{U}_{ij}
- $\text{BR}(h_i \rightarrow \tilde{\chi}_i^\pm \tilde{\chi}_j^\mp, \tilde{\chi}_i^0 \tilde{\chi}_j^0)$: decay to charginos, neutralinos: \mathbf{U}_{ij}

FeynHiggs “workflow”:



- Renormalization of the CP-conserving Higgs-sector
[Drechsel, Galeta, Heinemeyer, Weiglein, (2016)];
- CP-violating NMSSM, on-shell neutral Higgs *[Drechsel, F.D., Paßehr (2017)];*
- Neutral Higgs decays into SM particles at full one-loop order
[F.D., Heinemeyer, Paßehr, Weiglein, (2018)];
- Higgs-to-Higgs + Higgs-to-SUSY on-going...
- Inclusion within FeynHiggs in an unforeseeable future...
*Learn from MSSM and adapt to the NMSSM.
+ Rejuvenate MSSM from NMSSM.*

In the future: **FeynHiggs 3.0**

⇒ few numerical examples for the Higgs decays

Overall (N)MSSM Higgs decay uncertainty estimates

- $h_i \rightarrow q\bar{q}$: SM-like: SM NNLO QCD, EW NNLO, SUSY 2L: $\sim 5\%$
heavy: as SM-like, Sudakov logs: $\sim 5 - 10\%$
- $h_i \rightarrow \ell\bar{\ell}$: SM-like: $\lesssim 1\%$
heavy: Sudakov logs for very heavy Higgses $\lesssim 10\%$
- $h_i \rightarrow WW^{(*)}, ZZ^{(*)}$: SM-like: $\lesssim 1\%$
heavy: missing 2L (very small width): $\lesssim 50\%$
- $h_i \rightarrow \gamma\gamma, gg, \gamma Z$: $\gamma\gamma$: NNLO QCD, EW: $\lesssim 4\%$
 gg : NNLO QCD, EW: $\lesssim 4\%$
 γZ : NLO: $\sim 5\%$
- $h_i \rightarrow \text{SUSY SUSY}$: [S.H., C. Schappacher '14-'16]
1L effects $10 - 20\%$, 2L?
- all decays: U_{ij}, Z_{ij} : few %, effects close to threshold?

\Rightarrow approaching LC precision for SM-like Higgs (not for heavy Higgses yet)

5. Conclusinos

- High precision prediction for cross sections and branching ratios are crucial for coupling constant determination
- Prediction (SM, MSSM, BMSSM) needed at/below the percent level!
- SUSY Higgs boson masses:
 - hybrid approach is best \Rightarrow FeynHiggs
 - MSSM: uncertainty in M_h reduced to 1-2 GeV (in simple single-scale scenario)
 - NMSSM: uncertainties intrinsically larger, not calculated
- MSSM Higgs production cross sections:

Now available in the cMSSM at the full one-loop level:

$$\sigma(e^+e^- \rightarrow h_i h_j), \sigma(e^+e^- \rightarrow h_i Z), \sigma(e^+e^- \rightarrow h_i \gamma)$$
$$\sigma(e^+e^- \rightarrow H^+ H^-), \sigma(e^+e^- \rightarrow H^\pm W^\mp)$$
 - Tree-level procs: loop corrections crucial ($e^+e^- \rightarrow h_1 h_2, h_1 Z, \dots, H^+ H^-$)
 - Loop induced procs: possibly observable ($e^+e^- \rightarrow h_1 h_1, h_1 \gamma, \dots, H^\pm W^\mp$) \Rightarrow polarization could be crucial!
 \Rightarrow possibly relevant MSSM parameter dependence
- (N)MSSM Higgs decays:

approaching LC precision for SM-like Higgs (not for heavy Higgses yet)

Further Questions?



Charged MSSM Higgs Production at the LC

Charged Higgs production:

$$e^+e^- \rightarrow H^+H^-, H^\pm W^\mp, H^\pm e^\mp \nu, H^\pm tb, \dots$$

Now available in the **cMSSM** at the full one-loop level:

[*S.H., C. Schappacher '17*]

$$\sigma(e^+e^- \rightarrow H^+H^-)$$

$$\sigma(e^+e^- \rightarrow H^\pm W^\mp)$$

In the following:

few examples of each process, relevance of loop corrections

cMSSM parameters:



Table 1: MSSM default parameters for the numerical investigation; all parameters (except of t_β) are in GeV. The values for the trilinear sfermion Higgs couplings, $A_{t,b,\tau}$ are chosen such that charge- and/or color-breaking minima are avoided [64], and $A_{b,\tau}$ are chosen to be real. It should be noted that for the first and second generation of sfermions we chose instead $A_f = 0$, $M_{\tilde{Q},\tilde{U},\tilde{D}} = 1500$ GeV and $M_{\tilde{L},\tilde{E}} = 500$ GeV.

Scen.	\sqrt{s}	t_β	μ	M_{H^\pm}	$M_{\tilde{Q},\tilde{U},\tilde{D}}$	$M_{\tilde{L},\tilde{E}}$	$ A_{t,b,\tau} $	M_1	M_2	M_3
S1	1000	7	200	300	1000	500	$1500 + \mu/t_\beta$	100	200	1500
S2	800	4	200	300	1000	500	$1500 + \mu/t_\beta$	100	200	1500


with \sqrt{s} , M_{H^\pm} , $\tan \beta$, ϕ_{A_t} varied

- Scenario chosen such that many processes are possible at the same time
- not chosen to maximize loop corrections

$$\underline{e^+e^- \rightarrow H^+H^-}$$




$e^+e^- \rightarrow H^+H^-$:




\Rightarrow loop corrections non-negligible!

$e^+e^- \rightarrow H^+H^-$:



\Rightarrow loop corrections non-negligible!


$e^+e^- \rightarrow H^+H^-$:




⇒ loop corrections sizable for large $\tan \beta$

⇒ no relevant complex phase dependence

$e^+e^- \rightarrow H^\pm W^\mp$: purely loop-induced!




$e^+e^- \rightarrow H^\pm W^\mp$:




⇒ small loop-induced cross section . . .

$$\underline{e^+e^- \rightarrow H^\pm W^\mp:}$$




Polarization: $P(e^-) = +80\%$, $P(e^+) = -30\%$
 \Rightarrow crucial to yield detectable cross section!




SM-like Higgs state

- Radiative corrections dominated by QCD-corrections;
- Unitary tree-level approximation works well;
- Difference wrt. FH (small): $h \rightarrow g(g^* \rightarrow b\bar{b})$
(whether $h \rightarrow b\bar{b}$ or $h \rightarrow gg$ is an experimental question).



Heavy doublet Higgs states at ~ 1 TeV


- Sizable EW corrections due to Sudakov logarithms;
- Unitary tree-level approximation ‘fails’ $\sim 10\%$ off;
- Difference wrt. FH (minor): UV scale in Δ_b (higher-order).



Heavy CP-even doublet Higgs state


- black curve: SM 1L prediction of Prophecy4f rescaled (as FH);
- Red curve: full one-loop (on-shell);
- Rescaling procedure fails for a decoupling state

$$g^{HVV} / g^{H_{SM}VV} \simeq 0.$$



- HDECAY provides a QCD/large $\tan\beta$ -corrected width (including SQCD) \simeq our green line;
- Full one-loop shows EW Sudakov logarithms for heavy states.

Here: h_1 SM-like; h_2 (640 GeV) and a_1 (320 GeV) singlet-like; h_3 and a_2 doublet-like (1 TeV).



- HDECAY performs at the same order as us with 5-flavor radiation;
- $\sim 4\%$ deviation due to normalization factor (difference of EW 2-loop and QCD 3-loop order).

Here: h_1 SM-like; h_2 (650 GeV) and a_1 (320 GeV) singlet-like; h_3 and a_2 doublet-like (1 TeV).