



université
PARIS-SACLAY



Update of $t\bar{t}$ Analysis


09/25/20 Y. Okugawa, R. Pöschl

ILC Summer Camp 2020

1. Introduction

Top Quark Pair Production at ILC

- ILC at $\sqrt{s} = 500$ GeV will produce many $t\bar{t}$ pairs, which allows a precision measurement on the heavy quark properties.
 - Play a central role for the indirect searches of the new particle beyond the Standard Model predictions to distinguish them from the various other theories.



u

d

s

c

b

t



1. Introduction

- The couplings between pair of top and are parametrized in terms of form factors. A_{FB}^t is the measure for the level of the parity violation.

- ILC Integrated Luminosity: $\int L dt = 4,000 \text{ fb}^{-1}$

- $t\bar{t}$ cross section

► $\sigma_{unpol} = 572 \text{ fb}$, $\sigma_{eLpR} = 1564 \text{ fb}$, $\sigma_{eRpL} = 724 \text{ fb}$



T. Han 2005 hep-ph/0508097

2. Analysis

- Semi-leptonic process

- 4-jets (final state: $b\bar{b}q\bar{q}'\ell\bar{\nu}$)

- One isolated lepton, e or μ .

(τ is ignored for the time being)



- eLpR process

- Observable:

$$A_{FB}^t = \frac{\overbrace{N(\cos \theta_t > 0)}^{\text{forward}} - \overbrace{N(\cos \theta_t < 0)}^{\text{backward}}}{\overbrace{N(\cos \theta_t > 0)}^{\text{forward}} + \overbrace{N(\cos \theta_t < 0)}^{\text{backward}}}$$

$$\sigma_{\mathcal{P}_{e^-}, \mathcal{P}_{e^+}} = \frac{1}{4} [(1 - \mathcal{P}_{e^-} \mathcal{P}_{e^+})(\sigma_{L,R} + \sigma_{R,L}) + (\mathcal{P}_{e^-} - \mathcal{P}_{e^+})(\sigma_{R,L} - \sigma_{L,R})]$$

| | Final States | # of jets | B.R. |
|---------------|---|-------------------|-------|
| Full Leptonic | $t\bar{t} \rightarrow (b\ell\bar{\nu})(\bar{b}\ell\nu)$ | 2 jets + 2 ℓ | 10.5% |
| Semi Leptonic | $t\bar{t} \rightarrow (b\ell\bar{\nu})(\bar{b}q\bar{q}')$ | 4 jets + 1 ℓ | 43.8% |
| Full Hadronic | $t\bar{t} \rightarrow (bq\bar{q}')(b\bar{q}q')$ | 6 jets | 45.7% |



2. Analysis

- Semi-leptonic process

- 4-jets (final state: $b\bar{b}q\bar{q}'\ell\bar{\nu}$)

- One isolated lepton, e or μ .

(τ is ignored for the time being)


- eLpR process

- Observable:

$$A_{FB}^t = \frac{\overbrace{N(\cos \theta_t > 0)}^{\text{forward}} - \overbrace{N(\cos \theta_t < 0)}^{\text{backward}}}{\overbrace{N(\cos \theta_t > 0)}^{\text{forward}} + \overbrace{N(\cos \theta_t < 0)}^{\text{backward}}}$$

$$\sigma_{\mathcal{P}_{e^-}, \mathcal{P}_{e^+}} = \frac{1}{4} [(1 - \mathcal{P}_{e^-} \mathcal{P}_{e^+})(\sigma_{L,R} + \sigma_{R,L}) + (\mathcal{P}_{e^-} - \mathcal{P}_{e^+})(\sigma_{R,L} - \sigma_{L,R})]$$

| | Final States | # of jets | B.R. |
|---------------|---|-------------------|-------|
| Full Leptonic | $t\bar{t} \rightarrow (b\ell\bar{\nu})(\bar{b}\ell\nu)$ | 2 jets + 2 ℓ | 10.5% |
| Semi Leptonic | $t\bar{t} \rightarrow (b\ell\bar{\nu})(\bar{b}q\bar{q}')$ | 4 jets + 1 ℓ | 43.8% |
| Full Hadronic | $t\bar{t} \rightarrow (bq\bar{q}')(b\bar{q}q')$ | 6 jets | 45.7% |



2. Analysis

- Semi-leptonic process

- 4-jets (final state: $b\bar{b}q\bar{q}'\ell\bar{\nu}$)

- One isolated lepton, e or μ .

(τ is ignored for the time being)


- eLpR process

- Observable:


$$A_{FB}^t = \frac{\overbrace{N(\cos \theta_t > 0)}^{\text{forward}} - \overbrace{N(\cos \theta_t < 0)}^{\text{backward}}}{\overbrace{N(\cos \theta_t > 0)}^{\text{forward}} + \overbrace{N(\cos \theta_t < 0)}^{\text{backward}}}$$

$$\sigma_{\mathcal{P}_{e^-}, \mathcal{P}_{e^+}} = \frac{1}{4} [(1 - \mathcal{P}_{e^-} \mathcal{P}_{e^+})(\sigma_{L,R} + \sigma_{R,L}) + (\mathcal{P}_{e^-} - \mathcal{P}_{e^+})(\sigma_{R,L} - \sigma_{L,R})]$$

| | Final States | # of jets | B.R. |
|---------------|---|-------------------|-------|
| Full Leptonic | $t\bar{t} \rightarrow (b\ell\bar{\nu})(\bar{b}\ell\nu)$ | 2 jets + 2 ℓ | 10.5% |
| Semi Leptonic | $t\bar{t} \rightarrow (b\ell\bar{\nu})(\bar{b}q\bar{q}')$ | 4 jets + 1 ℓ | 43.8% |
| Full Hadronic | $t\bar{t} \rightarrow (bq\bar{q}')(bq\bar{q}')$ | 6 jets | 45.7% |



2. Analysis




Semi-Leptonic Condition

| | Semi-Leptonic |
|-----------------|---------------------------|
| Isolated Lepton | $P_{lep} > 5 \text{ GeV}$ |
| N jets | 4 |
| W^\pm reco | isoLep + q jet |

Cuts

| 種類 | カット |
|------------------------|----------------------------|
| b-tag | $0.8 < b\text{-tag} < 0.3$ |
| Thrust | $\text{Thrust} > 0.9$ |
| Mhad (GeV) | $180 < M_{had} < 420$ |
| Top mass (GeV) | $120 < M_{top} < 270$ |
| W^\pm had mass (GeV) | $50 < M_W < 270$ |

3. Polar Angle




- ▶ Polar angle distribution of top quark for all reconstructed events

Result and Precision



| $(\mathcal{P}_{e^-}, \mathcal{P}_{e^+})$ | $(-1, +1)$ | $(+1, -1)$ |
|--|------------|------------|
| $A_{FB,gen}^t$ | 0.364 | 0.409 |
| $A_{FB,reco}^t$ | 0.345 | 0.369 |
| $\delta_{A_{FB}^t}$ | 0.0025 | 0.0020 |
| Efficiency | 34.6% | 64.1% |

- Precision $A_{FB}^t \approx 0.2\%$ is achieved for both left and right handed full polarized samples.
- Efficiencies reconstructing A_{FB}^t for left-handed is lower due to kinematic constraints.
 - ▶ Left-handed : b -jet follows the top flight path.
 - ▶ Right-handed : W follows the top flight path.

3. Polar Angle




- ▶ Polar angle distribution of top quark for all reconstructed events




- ▶ Polar angle distribution of top quark only using vtx x vtx comparison.

3. Polar Angle




- ▶ Polar angle distribution of top quark for all reconstructed events




Background?

- Mis-combination of b and W?
- Single Top Background?




- ▶ Polar angle distribution of top quark only using vtx x vtx comparison.

3. Polar Angle




► Polar angle distribution of top quark for all reconstructed events



Background?



- Mis-combination of b and W?
- Single Top Background?

Source of systemic error



► Polar angle distribution of top quark only using vtx x vtx comparison.

4. Single Top Analysis



- Two processes are difficult to be distinguished.
 - Share the same final states. ($b\bar{b}q\bar{q}'\ell\bar{\nu}$)
 - Events are mixed in the parton level.


4. Single Top Analysis

- This analysis considered events to correspond to top quark pair production when the following criteria is satisfied for both of W and b pairs.

$$|m_{Wb} - m_t^{MC}| < 15 \text{ GeV}$$

If only one of these pair meet this criterium, the events are labeled as single top quark event.

Fuster, J 2015 1434-6052




4. Single Top Analysis

- This analysis considered events to correspond to top quark pair production when the following criteria is satisfied for both of W and b pairs.

$$|m_{Wb} - m_t^{MC}| < 15 \text{ GeV}$$


If only one of these pair meet this criterium, the events are labeled as single top quark event.

Fuster, J 2015 1434-6052





4. Single Top Analysis

Polar Angle Distribution






- Polar angle distribution of top quark for all reconstructed events **after single top rejection from parton level**



- Polar angle distribution of top quark only using $vtx \times vtx$ comparison **after single top rejection from parton level**

4. Single Top Analysis

b-jet Energy Distribution




- *b*-jet energy distribution of hadronic top for all reconstructed events.


- *b*-jet energy distribution of hadronic top only using $\text{vtx} \times \text{vtx}$ comparison.

4. Single Top Analysis

b -jet Energy Distribution




- ▶ b -jet energy distribution of hadronic top for all reconstructed events.



- ▶ b -jet energy distribution of hadronic top only using vtx x vtx comparison.

➡ vtx x vtx method filters soft b 's?



5. Summary

- **$t\bar{t}$ Pair Production**

- ▶ $t\bar{t}$ production at the ILC at $\sqrt{s} = 500$ GeV for fully-left handed beam polarization using 900,000 events was processed.
- ▶ The ILC is capable of precision measurement of A_{FB}^t up to 0.2% of systematic error.

- **Single Top Analysis**

- ▶ Single top problem emerged as a source of systematic error, thus applied a selection for single top generated events on combined generated mass of b and W .
- ▶ Generated single top events consist 12.5% of overall events.
- ▶ $V_{tx} \times V_{tx}$ comparison scheme seems to eliminate such events by filtering out the soft b-jets.
 - ➡ Might worth to take a look at momentum distributions of tracks from b-jets in single top events to see if one of jets is indeed soft.