

ILC as a SUSY discovery and precision instrument

Teresa Núñez - DESY

On behalf of the ILD Concept Group



Chargino searches

M.T. Núñez Pardo de Vera¹, M. Berggren¹, J. List¹

¹ DESY

From light Higgsinos to test of unification

H. Baer¹, M. Berggren², K. Fujii³, J. List², S.-L. Lehtinen^{4,2}, T. Tanabe⁵, J. Yan³

¹ University of Oklahoma, ² DESY, ³ KEK, ⁴ University of Hamburg, ⁵ ICEPP

Chargino searches

- Compute lighter chargino pair production cross sections in a wide SUSY parameter space (only using MSSM as model and R-parity conservation)
- Determine case with lowest production cross sections
- Compare to cross section detection limits extrapolated from LEP results (in the worst scenario)

One of the prime candidates to be the NLSP

Cross section studies divided depending on:


- chargino mixing: Higgsino-like, Wino-like and Mixed charginos
- sneutrino mass scale: high ($\sim 1\text{TeV}$), low (around kinematic limit)

Produced (in e^+e^- collisions) via Z/γ annihilation in the s-channel and sneutrino exchange in the t-channel

Chargino searches: cross sections

Worst scenario for Wino and Higgsino-like charginos

Cross sections




$P(e^-, e^+) = (-80\%, +30\%)$

Lower efficiencies reached in Wino-like case with sfermions masses close to kinematic limit

Chargino searches: cross sections

Worst scenario for Wino and Higgsino-like charginos

Cross sections




Cross sections compared to limits from LEP extrapolated to 1.6 ab^{-1} integrated luminosity (ILC500, $P(e^-,e^+) = (-80\%, +30\%)$), based on the dependency $\sigma_{\text{LIM}} \sim \frac{1}{\sqrt{L}}$, to obtain mass limits for chargino exclusion/discovery

$P(e^-,e^+) = (-80\%, +30\%)$


Lower efficiencies reached in Wino-like case with sfermions masses close to kinematic limit

Chargino searches: Mass limits

Higgsino




Wino



Chargino searches: Mass limits

Higgsino

Wino




Chargino searches: Mass limits

Higgsino

Wino

Even in the particular case with sfermion masses close to the kinematic limit ...



... for $\Delta M > 3$ GeV exclusion is guaranteed up to $M \tilde{\chi}_1^\pm = 246$ GeV and discovery up to 243 GeV

... for $m_\pi < \Delta M < 3$ GeV exclusion is guaranteed up to $M \tilde{\chi}_1^\pm = 225$ GeV and discovery up to 205 GeV

Chargino searches: Conclusions

Improvements at the ILC

- **Polarisation (increases signal/background ratio)**
- **No trigger (increases detection efficiency and allows 'redundant' analysis) but ...**
ISR needed for gamma-gamma background suppression
- **Smaller beam size (increases detection efficiency by releasing ISR requests -> observation of decay length for soft events)**

General comments

- **Loop corrections are not included (increase chargino pair production cross sections)**
- **Low sfermions masses not taken into account in LEP analysis, they would imply:**
 - changes in chargino branching ratios and decay topology
 - sfermion production and possible discovery

The drop in cross section due to sfermions masses depends on the beam energy, can be shifted.
- **ISR request close to kinematic limits could cause unknown effects**

From Light Higgsinos to test of unification

Electroweak naturalness in simple SUSY models requires a cluster of four light Higgsinos

$\tilde{\chi}_1^\pm, \tilde{\chi}_1^0, \tilde{\chi}_2^0$ compressed spectrum (10-20 GeV) around ~ 100 -300 GeV

Challenging for LHC if other sparticles are heavy... but not for ILC:

- Electron-Positron collider at $\sqrt{s} = 250$ -500 GeV with energy upgradability (1TeV)
- Electrons (80%) and positrons (30%) polarisations
- Well defined initial state: 4-Momentum and spin configuration
- Clean and reconstructable final state
- Hermetic detectors (almost 4π coverage)
- Triggerless operation

From Light Higgsinos to test of unification

Examine the capability of the ILC to make precision measurements of superparticle properties offering the possibility to make important predictions of SUSY breaking and fundamental particle physics and providing insights into the nature of dark matter and cosmology

Three benchmark scenarios

ILC1 & ILC2:

- Natural models from NUHM2
- Gaugino mass unification at GUT scale

Two channels

$$e^+ e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0 q \bar{q} \tilde{\chi}_1^0 e \nu_e (\mu \nu_\mu)$$
$$e^+ e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 e^+ e^- (\mu^+ \mu^-)$$



nGMM1:

- Natural generalized mirage mediation model
- Gaugino mass unification at mirage scale

Light Higgsino masses ~ 150 GeV and mass differences 4-20 GeV

Direct measurement of masses and cross sections



Extracted from kinematic distributions in both channels under study



Chargino channel

Chargino and neutralino masses measured to less than 1% precision

Neutralino channel



$\sigma \times \text{BR}$ up to a few % precision, depending on channel and beam polarisation

Fitting fundamental parameters and testing gaugino mass unification

Extraction of both GUT-scale and weak-scale parameters with two different approaches (4 and 10 free parameters), extrapolation of these to the GUT-scale and dark matter implications

Input parameters:

- Four Higgsino masses
- Four polarised total cross sections from chargino and neutralino
- Higgs observables with ILC precision:
 - > Mass of the lightest CP-even Higgs boson
 - > Higgs decay branching ratios

Fitting fundamental parameters: GUT scale

Strongly dependent on underlying SUSY breaking scheme

None of the benchmarks can be excluded as NUHM2 model

Not underlying models (NUHM1, CMSSM): interpretations ruled out at 95% CL level with only 0.1 total integrated luminosity

Fitting fundamental parameters: Weak scale

Check if it possible to constrain a comprehensive set of parameters from observables of Higgsino sector alone

pMSSM-10 model

Study influence of parameters in which Higgsino sector enters at loop level


pMSSM-4 model

Investigate precision achievable when fitting only tree level Higgsino parameters

- Agreement between fit and model values in the three benchmarks. Less precision in nGMM1 due to worse experimental resolution and large absolute values
- Parameters entering at loop level in Higgsino sector do not have significant effect in Higgsino sector fits
- Improvement fitting only tree level Higgsion parameters

Fitting fundamental parameters: predicting SUSY spectrum

ILC1 (GUT-scale fit)



Clear electroweakinos and heavy Higgs bosons mass predictions


gluino mass firmly predicted (GUT check)

Rest of sfermions masses less constrained but upper limits can be obtained

Provides important information for a future hadron colliders

Fitting fundamental parameters: predicting SUSY spectrum

pMSSM-4 ILC1



Only tree level Higgsino sector parameter:
squark, sleptons, heavy Higgs boson and
gluino parameters fixed to its true value

Heavier electroweakinos mass precisions
between 1.6% and 3%

Changes in fixed parameters keeps
predictions within 1 σ uncertainties


Testing gaugino mass unification

MSSM RGEs used to evolve fitted parameters to higher energy scales and check unification hypothesis

Test gaugino mass unification in different models
Study impact of experimental improvements


- pMSSM-10 weak scale parameters extracted at $Q=10\text{GeV}$
- Scan around extracted values using corresponding PDF within $\pm 1\sigma$
- Calculate running parameters at different energy scales

Testing gaugino mass unification: ILC1




Assuming NUHM2 model, prediction of physical gluino mass can be obtained: $m_g = 1467 \pm 80$ GeV

Q_{unif} and $M_{1/2}$ in agreement with GUT scale model fit (same for ILC2)




Testing gaugino mass unification: nGMM1



Differences with ILC1 & ILC2:

- Underlying model unifies at intermediate scale
- M_1 and M_2 determination at weak scale much less accurate

Unification occurs at the model point after using hypothetical gluino mass



Conclusions

- **It was shown the reach of the ILC for discovering SUSY in the chargino channel within the worst scenario and in very conservative conditions. Mass limits up to few GeV below the kinematic limit have been found**
- **The capability of the ILC for measuring SUSY observables with precision enough to make relevant SUSY related predictions has been studied and confirmed**
- **Improvement on the experimental results are clearly reflected on the predictions**
- **The predictions extracted from direct SUSY measurements can have an important role in future accelerator designs and upgrades**
- **The interplay between ILC and LHC SUSY measurements can improve considerable the results of both analysis**
- **Interplay between ILC results and WIMP detection experiments can provide important information about the nature of dark matter and other dark matter related items**


Backup slides

Chargino searches: cross sections

High sfermion masses

$\sqrt{s} = 500$ GeV, ISR photon
ILC experimental conditions

Lower cross sections for Higgsino-like case




+/- \rightarrow $P(e^-,e^+) = (-80\%, +30\%)$
 -/+ \rightarrow $P(e^-,e^+) = (+80\%, -30\%)$

$P(e^-,e^+) = (-80\%, +30\%)$ with $\mathcal{L} = 1.6 \text{ ab}^{-1}$ will be used for the study

Chargino searches: cross sections

Low sfermion masses


Cross sections



Wino-like charginos

- Affects Wino case via destructive interference of t-channel
- No effect on Higgsino due to weakly coupling to sneutrino

Cross sections



Chargino searches: extrapolated data

Combined LEP chargino studies

- Data taken at up to ~ 208 GeV center-of-mass energy, accumulated luminosity $\sim 800 \text{ pb}^{-1}$
- No signal found, limits derived at 95%CL in the context of MSSM (R parity conservation) focused in the region with small $\Delta M = M_{\tilde{\chi}_1^\pm} - M_{\text{LSP}}$
- Two cases considered:
 - Higgsino-like
 - Wino-like (high sfermion masses)
- Three topologies for the analysis of chargino decays:
 - prompt decays into leptons, leptons + jets, jets via W^* ($\Delta M > 3\text{GeV}$)
 - soft decays with a ISR requested on trigger ($\pi \text{ mass} < \Delta M < 3\text{GeV}$)
 - events with tracks displaying kinks, impact parameters offset or heavy stable charged particles ($\Delta M < \pi \text{ mass}$)

Chargino searches: extrapolated data



Combined LEP chargino studies

- Data taken at up to ~208 GeV center-of-mass energy, accumulated luminosity ~800 pb⁻¹
- No signal found, limits derived at 95%CL in the context of MSSM (R parity conservation) focused in the region with small $\Delta M = M_{\tilde{\chi}_1^\pm} - M_{\text{LSP}}$
- Two cases considered:
 - High
 - Win
- Three topologies for the analysis of chargino decays:
 - prompt decays into leptons, leptons + jets, jets via W^* ($\Delta M > 3\text{GeV}$)
 - soft decays with a ISR requested on trigger ($\pi \text{ mass} < \Delta M < 3\text{GeV}$)
 - events with tracks displaying kinks, impact parameters offset or heavy stable charged particles ($\Delta M < \pi \text{ mass}$)

Cross sections compared to limits from LEP extrapolated to 1.6 fb⁻¹ integrated luminosity (ILC500, P(e-,e+)=(-80%,+30%)) based on the dependency $\sigma_{\text{LIM}} \sim \frac{1}{\sqrt{L}}$

Chargino searches:


Comparison to extrapolated limits



Low efficiency in π mass $< \Delta M < 3$ GeV region due to ISR trigger requirement

Chargino searches:

Comparison to extrapolated limits



Cross section limits extracted and extrapolated to 1.6 fb^{-1} integrated luminosity (ILC500, $P(e-,e+)=(-80\%,+30\%)$) based on the dependency $\sigma_{\text{LIM}} \sim \frac{1}{\sqrt{L}}$

Benchmark scenarios

ILC1 & ILC2:


- Natural models from NUHM2
- Gaugino mass unification at GUT scale

nGMM1:

- Natural generalized mirage mediation model
- Gaugino mass unification at mirage scale (between EW and GUT scales)

Masses (GeV)	ILC1	ILC2	nGMM1
neu1	103	148	151
Chi1+-	117	157.8	159
Neu2	124	158.3	156
Neu3	267	539	1530
Gluino	1560	2830	2860

ICHEP 2020, 29-07-20



Cross sections at $\sqrt{s} = 500$ GeV
several hundreds fb

CLUSTER OF EXCELLENCE
QUANTUM UNIVERSE



Software tools and observables

- Whizard1.95 for event generation
- ILD-specific software based on Geant4 for simulation and reconstruction
- Beam spectrum, ISR and $\gamma\gamma$ “pile-up” included
- $\sqrt{s} = 500$ GeV and $\mathcal{L} = 500$ fb⁻¹ simulation results scaled to operation scenarios

- Processes studied:

$$e^+ e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^- \rightarrow \tilde{\chi}_1^0 q \bar{q} \tilde{\chi}_1^0 e \nu_e (\mu \nu_\mu)$$

$$e^+ e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0 e^+ e^- (\mu^+ \mu^-)$$

- Observables:

- three masses ($\tilde{\chi}_1^\pm$, $\tilde{\chi}_1^0$, $\tilde{\chi}_2^0$)

- four cross sections

- Masses from kinematical distributions (maximum invariant mass -> mass splittings, maximum dilepton/dijet energy -> absolute masses)
- Cross sections from counting events after fitting overall shape

Fitting fundamental parameters: GUT scale




Strongly dependent on underlying SUSY breaking scheme

Fit to NUMH2 model

	ILC1 best fit (true)	ILC2 best fit (true)	nGMM1 best fit
$M_{1/2}$	556.7 (568.3)	1194 (1200)	2407
μ	105.3 (115.0)	150.7 (150.0)	155.6
$\tan\beta$	11.4 (10.0)	16.0 (15.0)	10.0
m_A	968 (1000)	1008 (1000)	1603
M_0	7685 (7025)	4788 (5000)	3422
A_0	-11064 (-10427)	-7663 (-8000)	-7409
χ^2	0.0011 (0.0013)	0.02848 (0.0007)	0.233

Fitting fundamental parameters: predicting SUSY spectrum

Clear electroweakinos and heavy Higgs bosons mass predictions (ILC1 & ILC2)



gluino mass firmly predicted (GUT check)

Rest of sfermions masses less constrained but upper limits can be obtained

Provides important information for a future hadron colliders




Fitting fundamental parameters: Weak scale

- Check if it possible to constrain a comprehensive set of parameters from observables of Higgsino sector alone
- Study influence of parameters in which Higgsino sector enters at loop level
- Investigate precision achievable when fitting only tree level Higgsino parameters

Two models used: pMSSM-10, pMSSM-4

Fit parameters Higgsino sector

ILC1



Fitting fundamental parameters: Weak scale

Weak-scale parameters





- Check if it possible to constrain a comprehensive set of parameters from observables of Higgsino sector alone
- Study influence of parameters in which Higgsino sector enters at loop level
- Investigate precision achievable when fitting only tree level Higgsino parameters

Two models used: pMSSM-10, pMSSM-4

Fit parameters Higgsino sector

ILC2

ILC2 shows same behaviour



Fitting fundamental parameters: Weak scale


Weak-scale parameters

- Check if it possible to constrain a comprehensive set of parameters from observables of Higgsino sector alone
- Study influence of parameters in which Higgsino sector enters at loop level
- Investigate precision achievable when fitting only tree level Higgsino parameters

Two models used: pMSSM-10, pMSSM-4

Fit parameters Higgsino sector


nGMM1



Fitting fundamental parameters: Weak scale



Improvement fit parameters by using mass differences

pMSSM-10 nGMM1





Fitting fundamental parameters: Weak scale

Constraints for parameters of colored sector can be obtained





Hypothetical gluino mass measurement from LHC with 11% precision improves M3 precision accordingly, not significant effect on the rest



Fitting fundamental parameters: Weak scale

ILC2 predictions overestimate sfermions masses but still within 1σ


pMSSM-10 ILC2




M3 only constrained via its loop effects on Higgsino sector, since gluino probably outside LHC reach

Improvement by using mass differences

pMSSM-10 nGMM1



Fit with absolute masses






pMSSM-4 fit predicts heavier electroweakino masses within an 1σ uncertainty (~ 150 GeV) with not improved fit



Dark Matter in Higgsino fits

Fits to MSSM parameters allows extraction of WIMP Dark Matter related observables

Thermally produced WIMP relic density




Dark Matter in Higgsino fits

Fits to MSSM parameters allows extraction of WIMP Dark Matter related observables

Thermally produced WIMP relic density

Agreement model with fit values




Such measurement would confirm possible underabundance of Higgsino-like WIMPs





Dark Matter in Higgsino fits

Fits to MSSM parameters allows extraction of WIMP Dark Matter related observables


Thermally produced WIMP relic density



Testing gaugino mass unification: ILC1




Testing gaugino mass unification: ILC2





Assuming NUHM2 model and that M_3 unifies, prediction of physical gluino mass can be obtained: $m_g = 2872 \pm 605$ GeV

Possible to verify that M_1 and M_2 unify at GUT scale





Testing gaugino mass unification: ILC2

Assuming NUHM2 model and that M_3 unifies, prediction of physical gluino mass can be obtained: $m_g = 2872 \pm 605$ GeV



Possible to verify that M_1 and M_2 unify at GUT scale



Testing gaugino mass unification: nGMM1

