

Contribution ID: 127

Type: Oral presentation using Zoom

Di-Higgs production ($\gamma\gamma \rightarrow hh$ **) in Composite Models**

Thursday, 28 October 2021 14:20 (20 minutes)

In Standard Model (SM) Higgs Boson pair production initiated by photons $(\gamma \gamma \rightarrow hh)$ loop generated process and thereby is very sensitive to any new couplings and particles that may come in loops. Composite Higgs Models (CHMs) provide an alternate mechanism to address the hierarchy problem of SM where Higgs could be a bound state of a strongly interacting sector instead of being an elementary field. These set of models apart from modifying the SM Higgs couplings could also introduce new effective couplings that can have substantial impact on the loop processes. In this work we have studied the impact of Composite Higgs models in $\gamma \gamma \rightarrow hh$ (Di-Higgs) production process.

1st preferred time slot for your oral presentation

19:00-21:00 JST (12:00-14:00 CEST, 6:00-8:00 EDT, 3:00-5:00 PDT)

2nd preferred time slot for your oral presentation

15:30-17:30 JST (8:30-10:30 CEST, 2:30-4:30 EDT, 23:30-1:30 PDT)

Primary author: HARADA, Daisuke (KEK / Graduate University for Advanced Studies)
Presenter: HARADA, Daisuke (KEK / Graduate University for Advanced Studies)
Session Classification: F-2: Higgs properties

Track Classification: Parallel sessions: Topical Groups: Session F: Higgs properties