

Contribution ID: 354

Type: not specified

Let's measure the F2-gamma Structure -function

Wednesday, 27 October 2021 20:15 (20 minutes)

A higher energy eplus-eminus collider than LEP should have a better luminometer , optimised to measure the acollinearity and energies of the leptons in Bhabha scattering events. Building upon what we did with OPAL, such a luminometer could be used to make even more accurate measurements of Q-squared in deep-inelastic gamma-gamma scattering up to much higher values than LEP could reach. A new central hadron detector would give better measurements of the invariant-mass-squared W-squared of the recoiling hadrons and give better accuracy for x=Q-squared/(Q-squared + W-squared).

The work still to be done includes writing and testing a Monte Carlo generator for fragmentation of the final state hadrons, so that the structure function can be unfolded from the observed distributions of x and Q-squared, in the way used by my former student Edmund Clay in the last of our published OPAL F2-gamma analyses. His thesis is available as a ps.gz file on the UCL HEP group website. The only other extra work would be to write a simulation program for the performance of the luminometer as a tagger. The hard work for this may already have been done in optimising the luminosity measurement, maybe by Sailer from DESY, forCLIC. ILC ,CLIC or FCC eplus-eminus should all be similar.

Primary author: MILLER, David (UCL)Presenter: MILLER, David (UCL)Session Classification: B-3: Calorimeters