

Kinematic fitting for arbitrary resolution functions

Yu Kato, Tomohiko Tanabe, Masakazu Kurata,
Shogo Kajiwara, Takanori Mogi, Satoru Yamashita

The University of Tokyo, KEK, ICEPP

The 71st General Meeting of ILC Physics Subgroup

May 12, 2021

Overview

- We are developing kinematic fitter based on the log-likelihood method.
 - in order to deal with non-Gaussian distributions
 - by implementation of numerical differentiation
- For proof of principle, we try to fit $e^+e^- \rightarrow ZH \rightarrow \mu\mu bb$ at 250 GeV.
 - This process is selected for technical study of kinematic fitter.
 - No significant invisible particles \rightarrow good for ISR fit
 - The b-jet has asymmetric energy distribution.
 - We evaluated b-jet resolutions and compared fit result with uds-jet resolution.
- The study of Higgs light mediator decay has been started.
 - We expect our fitter is effective to this channel.
 - Sample generation is finished.


Introduction: Kinematic fit

- Kinematic fit:
 - one of the constrained optimization method
 - adjustment of measured kinematic parameters under certain constraints
 - distributions of parameters e.g. energy resolution
 - kinematic relations among the parameters e.g. energy conservation
- Purposes:
 - improve accuracy of measurements (reconstruction)
 - estimate how well a given event matches a signal model (event selection)
- Standard procedure: minimize χ^2

$$\chi^2(\boldsymbol{\eta}, \boldsymbol{\xi}, \boldsymbol{\lambda}) = (\mathbf{y} - \boldsymbol{\eta})^T \mathbf{V}^{-1} (\mathbf{y} - \boldsymbol{\eta}) - 2\boldsymbol{\lambda}^T \mathbf{h}(\boldsymbol{\eta}, \boldsymbol{\xi})$$

\mathbf{y} : measured variables
 $\boldsymbol{\eta}$: fit parameters
 \mathbf{V} : covariance matrix

$\boldsymbol{\xi}$: unmeasured parameters
 $\boldsymbol{\lambda}$: Lagrange multipliers
 \mathbf{h} : constraint functions



Our approach for non-Gaussian distributions

- The basic method assumes that the measured parameters would have Gaussian error against the true value.
- In order to treat arbitrary error distributions, the chi-square term is re-defined as the log-likelihood function;

$$\chi^2(\boldsymbol{\eta}, \boldsymbol{\xi}, \boldsymbol{\lambda}) = -2\ln L_{fo}(\boldsymbol{\eta}) - 2\boldsymbol{\lambda}^T \mathbf{h}(\boldsymbol{\eta}, \boldsymbol{\xi}) - 2\ln L_{sc}(\boldsymbol{\eta}, \boldsymbol{\xi})$$
$$L_{fo}(\boldsymbol{\eta}) = \prod_{i=1}^n f_i(y_i; \eta_i) \quad L_{sc}(\boldsymbol{\eta}, \boldsymbol{\xi}) = \prod_{i=1}^m s_i(\boldsymbol{\eta}, \boldsymbol{\xi})$$

f_i : error distributions s_i : soft constraint distributions

Note:

- The error distributions are normalized as the peak position returns 1.
 - The soft constraint term is applied optionally.
- In the case of Gaussian distributions, the basic method is reproduced.

Notes on implementation

Requirements

- Numerical differentiation
 - Although the Gaussian case can be solved analytically, the arbitrary case needs numerical calculation.
- Resolution information
 - It is necessary to prepare the error distribution functions for each measured parameters.

Fitter algorithm


- Based on Sequential Quadratic Programming (SQP) method
- Hessian matrix is approximated by damped-BFGS method. (quasi-Newton method)
- The size of the iteration step (α) is adjusted by Armijo condition.

B-jet energy resolution

- The b-jet has asymmetric energy distribution due to neutrinos from semi-leptonic decay.
- We need to know the true energy distribution when a particular measured energy is obtained.
- The definition of the true jet:
 - Sum of the MCParticles which direction is close to reconstructed jet
 - Including neutrinos
- The resolutions are evaluated as the function of (E_{rec} , $\cos\theta_{\text{rec}}$) for each jet.

B-jet energy resolution: Evaluation setup


- Sample: b-jet pair
 - ILD DBD full simulation
 - E_{cm} : 20 - 240 GeV
 - PandoraPFA -> Durham jet clustering (LCFIPlus)
- Workflow:
 1. prepare data set of (E_{mc} , E_{rec}) in specific $\cos\theta_{rec}$ window
 2. generate E_{mc} histogram in specific E_{rec} window
 - normalized by all E_{rec} histogram
 - Each E_{mc} entry is shifted according as E_{rec} value.
 3. fit the spectrum
- Fit function: Crystal Ball (Gaussian & quartic polynomial)
 - p1: Gaussian mean
 - p2: Gaussian sigma
 - p3: Connection boundary in sigma unit



↑ True jet energy distribution
for $E_{rec} = 45.5 \pm 2.5$ GeV, $\cos\theta_{rec} = [0., 0.05]$

Test process: $e^+e^- \rightarrow ZH \rightarrow \mu\mu b\bar{b}$

- This process is selected for technical study of our kinematic fitter.



Simulation setup

- $\sqrt{s} = 250$ GeV
- $(P_{e^-}, P_{e^+}) = (-1, +1)$
- ILD DBD sample, $\sim 10k$ event
- Main background: 4f_ZZ_semileptonic

Event reconstruction

1. Particle reconstruction: PandoraPFA
2. Muon selection: IsolatedLeptonTaggingProcessor
3. Jet clustering & Flavor tagging : LCFIPlus
Durham (forced 2 jets)


Setup of kinematic fit for $e^+e^- \rightarrow ZH \rightarrow \mu\mu b\bar{b}$

Fit Objects:




- JetFitObject (JFO) x 2
 - parameter: (E, θ, φ) with b-jet resolution
 E : Crystal Ball, θ : Gaus, φ : Gaus
 - $\text{mass}^{\text{fit}} \equiv E^{\text{fit}}/E^{\text{meas.}} \times \text{mass}^{\text{meas.}}$
 - Resolutions are adjusted by $(E, \cos\theta)$ for each jet
- MuonFitObject (MFO) x 2
 - parameter: (P_t, θ, φ) with Gaussian error from track parameters
- ISRPhotonFitObject
 - parameter: P_z ($E_{\text{max}} = 31.5$ GeV)
$$\mathcal{P}(p_{z,\gamma}) = \frac{\beta}{2E_{\text{max}}} \cdot \left| \frac{p_{z,\gamma}}{E_{\text{max}}} \right|^{\beta-1} \quad \beta = \frac{2\alpha}{\pi} \left(\ln \frac{s}{m_e^2} - 1 \right)$$

Constraints:

- Hard:
 - Total Energy/ $P_x/P_y/P_z$ for all FOs
 - Higgs mass = 125 GeV for 2 JFOs
- Soft:
 - Z mass w/ Breit-Wigner for 2 MFOs with mean 91.2 GeV and width 2.5 GeV





Fit Result: Hard constraint





- 9120 / 9128 events are converged.
- All the hard constraints are working correctly.
 - Energy momentum constraint is slightly deviated.

Fit Result: $\Delta E_j = E_{j\text{rec/fit}} - E_{j\text{mc}}$ ($E_{j_1\text{rec}} > E_{j_2\text{rec}}$)



- The jet energies are significantly recovered!
- The error distributions after fit looks symmetry.




Fit Result: Soft constraint, ISR



- ISR effect is also recovered!
- M_Z hasn't changed so much because of good muon resolution.


Fit Result: for Event Selection

- Using χ^2 distribution, event selection is tested.
- Event number is just sample number, not exact rate.
- These results are very preliminary, but b-jet case looks better than uds case.




ZH-> $\mu\mu\phi\phi$ -> $\mu\mu bbbb$ at 250 GeV

- Motivation:
 - Higgs exotic decay ($H \rightarrow \phi\phi \rightarrow bbbb$)
 - To apply b-jet kinematic fitter
- Setup:
 - WHIZARD 2.8.5
 - $m\phi$: 15, 20, 30, 40, 50, 60 [GeV]
 - 20,000 events for each $m\phi$, polarization{ (-0.8,+0.3), (+0.8,-0.3) }
 - mc-2020, $\sqrt{s} = 250$ GeV
- Status:
 - ✓ Sample preparation
 - Generate WHIZARD sample and some check at MC level
 - Simulate with DDSim, Reconstruct with MarlinStdReco
 - ❑ Fast analysis
 - ❑ Test fitting
 - ❑ Detailed analysis



S. Matsumoto(Kavli IPMU), ILC summer camp 2020



[arxiv.org/1612.09284](https://arxiv.org/abs/1612.09284)








Fig. 12. The 95% C.L. upper limit on selected Higgs exotic decay branching fractions at HL-LHC, CEPC, ILC and FCC-ee. The benchmark parameter choices are the same as in Table 3. We put several vertical lines in this figure to divide different types of Higgs exotic decays.

Summary and Outlook

- We are developing kinematic fitter based on the log-likelihood method.
 - to deal with non-Gaussian distributions
 - by implementation of numerical differentiation
- For proof of principle, we try to fit $e^+e^- \rightarrow ZH \rightarrow \mu\mu b\bar{b}$ at 250 GeV.
 - This process is selected for technical study of kinematic fitter.
 - We evaluated b-jet resolutions using true jet definition.
- Currently, we have confirmed a series of operations.
 - Fit results are obtained almost as we expected.
 - We also compared the results from b/uds resolutions and briefly obtained the benefit of b-resolution in event selection test.
- The study of Higgs light mediator decay has been started.
 - We expect our fitter is effective to this channel.
 - Sample generation is finished.


backup

Fit Result: χ^2 , Fit probability



Test of χ^2 and degrees of freedom





- Fast simulation
 1. Generate pseudo samples of $ZH \rightarrow 4$ particles which parameters $\{E, \theta, \phi\}$ have Gaussian errors;
 $\sigma_E = 1$ [GeV], $\sigma_\theta = 0.1$ [rad.], $\sigma_\phi = 0.1$ [rad.]
 2. Perform the kinematic fit under the 4 jets assumption which parameter errors are Gaussian above.
 3. Estimate the degrees of freedom from the χ^2 distribution when each constraint is applied;
4C: Energy momentum, 5C: Energy momentum & Higgs mass, 1C: Energy
- Results: fit probability



The χ^2 distributions show that the d.o.f. equals the number of constraints.
Our kinematic fitter evaluates the χ^2 output correctly in the simplest case.





B-jet energy resolution: Energy dependence

- Energy scan in the barrel region
 - $\cos\theta_{\text{rec}} = [0., 0.05]$
- In the higher edge the spectrum varies due to the lack of statistics.
- Parameters between points are interpolated.





B-jet energy resolution: Angle dependence


- Angle scan at $E_{\text{rec}} = 45.5 \text{ GeV}$
- JER is worse for forward jet as expected.







Fit Result: comparison of b/uds resolution




- We also obtained the result which use uds-jet resolutions.
- χ^2 is better for b-jet case.
- Fit probability flatness is slightly worse for b-jet case.
- A detailed check is in progress.







Fit Result: $\log_{10}\chi^2$, Fit probability, #Iteration




ISR spectrum


M. Beckmann, "Treatment of Photon Radiation in Kinematic Fits at Future e+e- Colliders"
F.A. Berends and R. Kleiss, Nucl. Phys. B177 (1981) 237

- ISR: $\mathcal{P}(p_{z,\gamma}) = \frac{\beta}{2E_{\max}} \cdot \left| \frac{p_{z,\gamma}}{E_{\max}} \right|^{\beta-1}$ $\beta = \frac{2\alpha}{\pi} \left(\ln \frac{s}{m_e^2} - 1 \right)$
- beamstrahlung: ?

ISR



beamstrahlung



Ecm - Z - H

