

Electroweak phase transition triggered by fermion sector

Katsuya Hashino (CHEP@Peking University)

Collaborators: Qing-Hong Cao^{1,2,3}, Xu-Xiang Li¹, Zhe Ren^{4,5}, Jiang-Hao Yu^{2,4,5,6,7}


(1. Department of Physics and SKLNPT at PKU, 2. CHEP at PKU, 3. CICQM, 4. ITP at CAS, 5. SPS at CAS, 6. HIAS at CAS, 7. ICTPAP)

[arXiv:2103.05688]

Introduction

- ★ The shape of Higgs potential is still undetermined...

$$V_{SM}(\phi) = \mu^2 |\phi|^2 + \lambda |\phi|^4 \quad (\text{The SM case})$$



- ★ The dynamics of electroweak phase transition (EWPT) is governed by the shape of the Higgs potential.

If the model realizes the first-order EWPT...

- Baryon asymmetry of the universe could be explained by electroweak baryogenesis scenario.
- The model could be tested by the measurement of gravitational wave from first-order PT.

How can we realize the first-order EWPT? What is a source of the EWPT?

First-order EWPT

- ★ Effective potential with high temperature approximation:

$$V_{\text{eff}}(\varphi, T) = D(T^2 - T_0^2)\varphi^2 - ET\varphi^3 + \frac{\lambda_T}{4}\varphi^4$$

E : Thermal loop effect of bosons

- ★ To realize first-order EWPT, it is necessary to develop a sizable barrier in the thermal potential.


The SM does not generate such a sizable barrier.

[Y. Aoki, F. Csikor, Z. Fodor and A. Ukawa, Phys. Rev. D 60, 013001 (1999)]

$$V_{\text{eff}}^{SM}(\varphi, T) \simeq D(T^2 - T_0^2)\varphi^2 + \frac{\lambda_T}{4}\varphi^4 \quad (\text{at } T \sim \varphi)$$

Usually, the sizable barrier could be developed by additional contribution to E term.

Although the fermion does not enhance the E term, the extended fermion model can generate a sizable barrier in the potential.



EWPT triggered by fermion sector

- ★ The fermion model could have additional reductions in φ^2 and φ^4 terms through new fermion effects.

$$V_{\text{eff}}(\varphi, T) = D(T^2 - T_0^2)\varphi^2 - ET\varphi^3 + \frac{\lambda_T}{4}\varphi^4$$


The new fermion effects make φ^2 and φ^4 terms comparable to φ^3 term.

- ★ The model with heavy fermion (around TeV scale) could generate a barrier through the new fermion effects.

$$V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\phi^2 - \frac{1}{4}\lambda\phi^4 + \frac{1}{6}\gamma\phi^6 \quad (\mu^2, \lambda, \gamma > 0)$$

Not only φ^3 term but also φ^4 term are negative.

The simple model with extended fermion sector could develop a sizable barrier by above two scenarios.




EWPT triggered by fermion sector

- ★ The fermion model could have additional reductions in φ^2 and φ^4 terms through new fermion effects.

$$V_{\text{eff}}(\varphi, T) = D(T^2 - T_0^2)\varphi^2 - ET\varphi^3 + \frac{\lambda_T}{4}\varphi^4$$

The new fermion effects make φ^2 and φ^4 terms




In this talk, we will discuss the detail in a simple fermion model with one isospin doublet fermion and one singlet neutral fermion.

- ★ generate a barrier through the new fermion effects.

$$V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\phi^2 - \frac{1}{4}\lambda\phi^4 + \frac{1}{6}\gamma\phi^6 \quad (\mu^2, \lambda, \gamma > 0)$$

Not only φ^3 term but also φ^4 term are negative.

The simple model with extended fermion sector could develop a sizable barrier by above two scenarios.



Extended fermion model

$$L = \begin{pmatrix} N \\ E \end{pmatrix}$$

- ★ The simple Lagrangian for new fermions

$$-\mathcal{L}_{VLL} = y_N (\bar{L}_L \tilde{H} N'_R + \bar{N}'_L \tilde{H}^\dagger L_R) + m_N \bar{L}_L L_R + m_L \bar{N}'_L N'_R + \text{h.c.}$$

Double lepton: L

Singlet neutral lepton: N'

SM-like Higgs doublet field : H

- ★ New parameters in the model: y_N, m_L, m_N

Mass parameter
region

A. $m_L \sim m_N \sim y_N v$ (Both fermions are at EW scale)

[M. Carena, A. Megevand, M. Quiros and C. E. Wagner, Nucl. Phys. B 716 (2005), 319, M. Fairbairn and P. Grothaus, JHEP 10 (2013), 176, A. Aranda, E. Jiménez and C. A. Vaquera-Araujo, JHEP01 (2015), 070, D. Egana-Ugrinovic, JHEP 12 (2017), 064, A. Angelescu and P. Huang, Phys. Rev. D 99 (2019) no.5, 055023]

B. $m_L \gg m_N \gg y_N v, m_L \sim m_N \gg y_N v$, (Both are at TeV scale)

C. $m_L \gg m_N \sim y_N v$ (One is at TeV scale, another is at EW scale)

[H. Davoudiasl, I. Lewis and E. Ponton, Phys. Rev. D 87 (2013) no.9, 093001, O. Matsedonskyi and G. Servant, JHEP 09 (2020), 012]

The light m_L region is already prohibited by collider experiments.

[A. M. Sirunyan et al. [CMS Collaboration], Phys. Rev. D 100, no. 5, 052003 (2019)]

How to generate a barrier?

★ Effective couplings

$$\lambda_{n,eff} \equiv \frac{1}{(n-1)!} \left. \frac{\partial^n}{\partial \varphi^n} (V_{eff} + \Delta V_T) \right|_{\varphi=0}$$

We extract the temperature dependence of the coefficient of φ^n in potential.

$$V_{eff}^P = \lambda_{1,eff} \varphi + \frac{\mu_{eff}^2}{2} \varphi^2 + \frac{\lambda_{3,eff}}{3} \varphi^3 + \frac{\lambda_{eff}}{4} \varphi^4 + \frac{\lambda_{5,eff}}{5} \varphi^5 + \frac{\gamma_{eff}}{6} \varphi^6 + \frac{\lambda_{7,eff}}{7} \varphi^7 + \frac{\delta_{eff}}{8} \varphi^8 + \frac{\lambda_{9,eff}}{9} \varphi^9 + \frac{\epsilon_{eff}}{10} \varphi^{10} + \mathcal{O}(\varphi^{11}).$$

★ Mass region (A) ($m_L \sim m_N \sim y_N v$)

$$(\mu_{eff}^{2 \text{ fermions}})^2 \simeq \frac{y_N^2}{12} \left(2T^2 - \frac{9m_L^2}{\pi^2} \left(\ln \frac{\alpha_F T^2}{v^2} - \frac{3}{2} \right) \right),$$

$$\lambda_{eff}^{2 \text{ fermions}} \simeq -\frac{y_N^4}{8\pi^2} \left(\ln \frac{\alpha_F T^2}{v^2} - \frac{3}{2} \right),$$

Dominant contributions at the critical temperature for EWPT.

Both terms have the same signs.

(- μ^2 - Fermion loop) φ^2
 (+ λ - Fermion loop) φ^4

Typically, there are no reduction effects in φ^2 and φ^4 terms at the same time.

How to generate a barrier?

★ Effective couplings

$$\lambda_{n,eff} \equiv \frac{1}{(n-1)!} \left. \frac{\partial^n}{\partial \varphi^n} (V_{eff} + \Delta V_T) \right|_{\varphi=0}$$

We extract the temperature dependence of the coefficient of φ^n in potential.

$$V_{eff}^P = \lambda_{1,eff} \varphi + \frac{\mu_{eff}^2}{2} \varphi^2 + \frac{\lambda_{3,eff}}{3} \varphi^3 + \frac{\lambda_{eff}}{4} \varphi^4 + \frac{\lambda_{5,eff}}{5} \varphi^5 + \frac{\gamma_{eff}}{6} \varphi^6 + \frac{\lambda_{7,eff}}{7} \varphi^7 + \frac{\delta_{eff}}{8} \varphi^8 + \frac{\lambda_{9,eff}}{9} \varphi^9 + \frac{\epsilon_{eff}}{10} \varphi^{10} + \mathcal{O}(\varphi^{11}).$$

It is difficult to generate a sizable barrier in this region.

★ Mass region (A) ($m_L \sim m_N \sim y_N v$)

$$(\mu_{eff}^{2 \text{ fermions}})^2 \simeq \frac{y_N^2}{12} \left(2T^2 - \frac{9m_L^2}{\pi^2} \left(\ln \frac{\alpha_F T^2}{v^2} - \frac{3}{2} \right) \right),$$

$$\lambda_{eff}^{2 \text{ fermions}} \simeq -\frac{y_N^4}{8\pi^2} \left(\ln \frac{\alpha_F T^2}{v^2} - \frac{3}{2} \right),$$

Both terms have the same signs.

(- μ^2 - Fermion loop) φ^2

(+ λ - Fermion loop) φ^4

Dominant contributions at the critical temperature for EWPT.


Typically, there are no reduction effects in φ^2 and φ^4 terms at the same time.

How to generate a barrier?

- ★ Mass region (B) and (C) have heavy fermion with TeV scale.

We used “matching method” to treat the multi-scale effective potential.

[Qing-Hong Cao, K. H., Xuxiang Li, Zhe Ren, Jiang-Hao Yu, arXiv:2103.05688]



$$\left. \begin{aligned} (\mu^2)^{LE} &= (1 + 2Z_\varphi^{\text{heavy}}) \mu^2 + \frac{\partial^2}{\partial \varphi^2} \sum_{i=\text{heavy fields}} \frac{n_i}{64\pi^2} M_i(\varphi)^4 \left(\log \frac{M_i(\varphi)^2}{Q_M^2} - c_i \right) \Bigg|_{\varphi=0} \\ \lambda^{LE} &= (1 + 4Z_\varphi^{\text{heavy}}) \lambda + \frac{1}{6} \frac{\partial^4}{\partial \varphi^4} \sum_{i=\text{heavy fields}} \frac{n_i}{64\pi^2} M_i(\varphi)^4 \left(\log \frac{M_i(\varphi)^2}{Q_M^2} - c_i \right) \Bigg|_{\varphi=0} \\ \lambda_{(n)}^{LE} &= \frac{1}{(n-1)!} \frac{\partial^n}{\partial \varphi^n} \sum_{i=\text{heavy fields}} \frac{n_i}{64\pi^2} M_i(\varphi)^4 \left(\log \frac{M_i(\varphi)^2}{Q_M^2} - c_i \right) \Bigg|_{\varphi=0} \end{aligned} \right\}$$

- ★ This treatment can be extended to other new physics models with heavy fields and light fields not limited to the fermion degree of freedom.

How to generate a barrier?

- ★ Mass region (B) ($m_L \gg m_N \gg y_N v$, $m_L \sim m_N \gg y_N v$)

$$\lambda_{eff}^{T=0} \sim \lambda_{eff}^{SM} - \frac{y_N^6 v^2}{8\pi^2 M_L^2}$$

$$(M_L \gg m_N \gg y_N v)$$

$$\lambda_{eff}^{T=0} \sim \lambda_{eff}^{SM} - \frac{y_N^6 v^2}{160\pi^2 m_L^2}$$

$$(m_L \sim m_N \gg y_N v)$$

New fermion effects show up in the potential through the high dimensional operator, like second terms.

The contributions to ϕ^n terms are small...

It is difficult to generate a sizable barrier in this region.

- ★ Mass region (C) ($m_L \gg m_N \sim y_N v$)

$$(\mu_{eff}^{new\ fermion})^2 \simeq \boxed{\gamma_{eff}^{T=0} v^4} + \frac{y_N^2 X}{2(1-X)} \left(-\frac{T^2}{3} + \frac{X^2 m_L^2}{2\pi^2} \left(\ln \frac{\alpha_F T^2}{v^2} - \frac{3}{2} \right) \right),$$

$$\lambda_{eff}^{new\ fermion} \simeq \boxed{-2\gamma_{eff}^{T=0} v^2} + \frac{4y_N^4(1+X)}{16m_L^2(1-X)^3} \left(\frac{T^2}{3} - \frac{X^2 m_L^2(3-X)}{2\pi^2(1+X)} \left(\ln \frac{\alpha_F T^2}{v^2} - \frac{3}{2} \right) \right)$$

$$X = m_N/m_L$$



The contributions to ϕ^2 term are positive, while ones to ϕ^4 term are negative.

→ **The sizable barrier could be developed in the potential in this region.**

Mass region (C)

- ★ Temperature dependence of normalized effective couplings

Scenario (I) : $V_{\text{eff}}(\varphi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{3}\lambda_3\varphi^3 + \frac{1}{4}\lambda\varphi^4$




$y_N = 2.5, m_L = 2000 \text{ GeV}, m_N = 450 \text{ GeV}$

Mass region (C)


- ★ Temperature dependence of normalized effective couplings

Scenario (I) : $V_{\text{eff}}(\varphi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{3}\lambda_3\varphi^3 + \frac{1}{4}\lambda\varphi^4$



$y_N = 2.5, m_L = 2000 \text{ GeV}, m_N = 450 \text{ GeV}$

Scenario (II) : $V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{4}\lambda\varphi^4 + \frac{1}{6}\gamma\varphi^6$




$y_N = 2.5, m_L = 1750 \text{ GeV}, m_N = 500 \text{ GeV}$

Mass region (C)

- ★ Temperature dependence of normalized effective couplings


Scenario (I) : $V_{\text{eff}}(\varphi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{3}\lambda_3\varphi^3 + \frac{1}{4}\lambda\varphi^4$



$$y_N = 2.5, m_L = 2000 \text{ GeV}, m_N = 450 \text{ GeV}$$

$$\mu_{\text{eff}}^2 \supset \frac{y_N^2 m_L^2 X^3}{4\pi^2(1-X)} \left(\ln \frac{\alpha T^2}{v^2} - \frac{3}{2} \right), \lambda_{\text{eff}} \supset -\frac{y_N^4 X^2(3-X)}{8\pi^2(1-X)^3} \left(\ln \frac{\alpha T^2}{v^2} - \frac{3}{2} \right) \quad X = m_N/m_L$$

Scenario (II) : $V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{4}\lambda\varphi^4 + \frac{1}{6}\gamma\varphi^6$



$$y_N = 2.5, m_L = 1750 \text{ GeV}, m_N = 500 \text{ GeV}$$

Fermionic reduction contributions depends on the value of m_N/m_L .

Mass region (C)

- ★ Parameter region where a sizable barrier could be developed in mass region $m_L \gg m_N \sim y_N v$.

Orange (Scenario I)


$$V_{\text{eff}}(\varphi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{3}\lambda_3\varphi^3 + \frac{1}{4}\lambda\varphi^4 \left(+ \frac{\gamma}{6}\varphi^6 \right)$$

Green (Scenario II)

$$V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{4}\lambda\varphi^4 + \frac{1}{6}\gamma\varphi^6$$

Magenta and blue contours correspond to λ_{eff} and γ_{eff} at $T=0$.

A sizable barrier could be generated by large m_N/m_L value.



The first-order EWPT could be realized in the simple extended fermion model, especially, one is heavy (TeV scale) and another is light (EW scale).

Mass region (C)

- ★ Parameter region where a sizable barrier could be developed in mass region $m_L \gg m_N \sim y_N v$.

Orange (Scenario I)


$$V_{\text{eff}}(\varphi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{3}\lambda_3\varphi^3 + \frac{1}{4}\lambda\varphi^4 \left(+ \frac{\gamma}{6}\varphi^6 \right)$$

Green (Scenario II)

$$V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\phi^2 - \frac{1}{4}\lambda\phi^4 + \frac{1}{6}\gamma\phi^6$$

Magenta and blue contours correspond to λ_{eff} and γ_{eff} at $T=0$.

A sizable barrier could be generated by large m_N/m_L value.



The first-order EWPT could be realized in the simple extended fermion model, especially, one is heavy (TeV scale) and another is light (EW scale).

hhh coupling

★ Typically, the triple Higgs boson coupling in the model with strongly first-order EWPT is enhanced from the SM prediction value. [S. Kanemura, Y. Okada and E. Senaha, Phys. Lett. B 606 (2005), 361]

★ Triple Higgs boson coupling $\lambda_{hhh} \equiv \left. \frac{\partial^3 V_{eff}}{\partial \varphi^3} \right|_{\varphi=v}$

$$\lambda_{hhh}^{\text{new fermion}} \sim 8\gamma v^3 + \frac{y_N^6 v^3 X}{\pi^2 m_L^2 (1-X)^2 \left(1 - X - \frac{y_N^2 v^2}{X m_L^2}\right)}$$


The value of hhh coupling could be enhanced by large m_N/m_L value.

In these parameter region being able to generate a barrier, the value of hhh coupling is 10% larger than the SM prediction value.

Future collider experiments can measure the hhh coupling at 10% accuracy. [arXiv:1506.05992, PRD 97 (2018) no.11, 113004, PRD 100 (2019) no.9, 096001, EPJ. C 80 (2020) no.11, 1010, CERN Yellow Rep. Monogr. 7 (2019), 221]

We could check whether a first-order EWPT can be realized or not by measurements of the hhh coupling.

[Qing-Hong Cao, K. H., Xuxiang Li, Zhe Ren, Jiang-Hao Yu, arXiv:2103.05688]




Summary

- ★ In this time, we discuss the phase transition patterns in the simple fermion model, one isospin doublet and one singlet neutral fermions.
- ★ Although the fermion does not contribute to φ^3 term, a sizable barrier could be developed by the fermionic reduction effects.
- ★ Especially, the model with one heavy and one light fermions can realize the first-order electroweak phase transition by following two scenarios:

(I) scenario $V_{\text{eff}}(\varphi, T) \simeq \frac{1}{2}\mu^2\varphi^2 - \frac{1}{3}\lambda_3\varphi^3 + \frac{1}{4}\lambda\varphi^4 \quad (\mu^2, \lambda_3, \lambda > 0)$

(II) scenario $V_{\text{eff}}(\phi, T) \simeq \frac{1}{2}\mu^2\phi^2 - \frac{1}{4}\lambda\phi^4 + \frac{1}{6}\gamma\phi^6 \quad (\mu^2, \lambda, \gamma > 0)$



We could check whether a first-order EWPT can be realized in the extended fermion model or not by measurements of the $h h h$ coupling.