


S O K E N D A I



# Study of photon-associated Higgs production at the ILC

---

Yumi Aoki(SOKENDAI), Junping Tian(Tokyo Univ.)

, Keisuke Fujii(KEK),

Sunghoon Jung(Seoul National Univ.),

Junghwan Lee(Seoul National Univ.)

2021.10.20(Wed) @Software&Analysis mtg


# 1. Motivation

---

Find new physics via  $h\gamma\gamma$  and  $h\gamma Z$  couplings

$h\gamma Z$  coupling in the Standard Model (SM) is a loop induced coupling.  
 → We expect BSM amplitude can be larger than SM amplitude.

We are the first to study the  $h\gamma Z$  coupling bond using  $e^+e^- \rightarrow h\gamma$




Any deviation of the coupling constants from SM signals new physics.

In this talk, we will report the simulation results of cross-section at ILC and the constraints such as model independent and  $h\gamma Z$  coupling constant.

## 2. Theoretical framework for our analysis

### SM one-loop predictions

\*For unpolarized beam



SM cross sections by one loop calculation:  $\sqrt{s} = 250$  GeV

$$\sigma_{\text{SM}}^L = 0.35 \text{ fb for } (-100\%, +100\%)$$

$$\sigma_{\text{SM}}^R = 0.016 \text{ fb for } (+100\%, -100\%)$$

$$\sigma_{\text{SM}} = 0.20 \text{ fb for } (-80\%, +30\%)$$

This analysis  
is very challenging.

### 3. Experimental Method

---

The effective field theory (EFT) Lagrangian (model-independent)

$$L_{\gamma H} = L_{SM} + \frac{\zeta_{AZ}}{v} A_{\mu\nu} Z^{\mu\nu} H + \frac{\zeta_A}{2v} A_{\mu\nu} A^{\mu\nu} H$$

effective  $h\gamma Z$  coupling      effective  $h\gamma\gamma$  coupling

$A_{\mu\nu}, Z_{\mu\nu}$  : field strength tensors

$v$ : vacuum expectation value

cross section of  $e^+e^- \rightarrow \gamma H$


Phys.Rev. D94 (2016) 095015

$$\frac{\sigma_{\gamma H}}{\sigma_{SM}} = 1 - 273\zeta_A - 201\zeta_{AZ} \quad (\text{eLpR})$$

$$\frac{\sigma_{\gamma H}}{\sigma_{SM}} = 1 + 492\zeta_A - 311\zeta_{AZ} \quad (\text{eRpL})$$

Since  $\zeta_A$  is already constrained by measurement of  $h \rightarrow \gamma\gamma$  branching ratio at LHC, we can extract  $\zeta_{AZ}$  parameter by just measuring cross section for a single beam polarization.

## 4. Simulation framework



- $\sqrt{s}=250 \text{ GeV}$   
Integrated Luminosity:  $2000 \text{ fb}^{-1}$   
**( $900 \text{ fb}^{-1}$  each for Left / Right handed pol.)**
- background : 2f,4f (DBD sample)
- ISR and Beamstrahlung effects are included
- ILD full simulation (Mokka)
- Geant4 based, realistic detailed detector model
- Full reconstruction chain from detector signals  
to 4-vectors  
(iLCSoft v01-16-02/ MarlinReco, PandoraPFA, LCFI+,  
Isolated photon finder, jet clustering )

$h \rightarrow bb$


$h \rightarrow WW^*$   
fully hadronic

$h \rightarrow WW^*$   
semi leptonic

## 5. Pre-Event selection

### Isolated photon

- Photon ID
- $E_\gamma > 50 \text{ GeV}$



※ The split photon clusters within a small cone are recovered

- $(\cos\theta_{\text{cone}}=0.998)$


### Remaining particles other than photon

- Jet clustering (Durham)
- Flavor tagged (LCFI+)

For h $\rightarrow$ WW\* semi-leptonic : number of decay w to qq=1

For h $\rightarrow$ WW\* fully-hadronic : number of decay w to qq=2

## 5. Event selection



# 5. Signal Significance

---


$$\text{significance} = \frac{N_s}{\sqrt{N_s + N_B}}$$

N<sub>s</sub>: # of signal  
N<sub>B</sub>: # of bg

Left	2f	4f	total bg	Signal	Significance
Expected	$1.0 \times 10^8$	$3.7 \times 10^7$	$1.4 \times 10^8$	$1.1 \times 10^2$	$9.0 \times 10^{-3}$
Pre selection	$2.8 \times 10^7$	$1.6 \times 10^6$	$2.9 \times 10^7$	$9.9 \times 10^1$	$1.8 \times 10^{-2}$
b likeliness > 0.77	$2.2 \times 10^6$	$2.1 \times 10^4$	$2.2 \times 10^6$	$9.0 \times 10^1$	$6.0 \times 10^{-2}$
Emis < 35	$1.9 \times 10^6$	$1.6 \times 10^4$	$1.9 \times 10^6$	$8.2 \times 10^1$	$5.9 \times 10^{-2}$
mvabdt > 0.025	$1.9 \times 10^4$	$3.2 \times 10^2$	$2.0 \times 10^4$	$3.4 \times 10^1$	$2.4 \times 10^{-1}$
-0.92 < cosθ <sub>γ</sub> < 0.92	$1.2 \times 10^4$	$1.3 \times 10^2$	$1.2 \times 10^4$	$2.9 \times 10^1$	$2.6 \times 10^{-1}$

Right	2f	4f	total bg	Signal	Significance
Expected	$7.3 \times 10^7$	$4.6 \times 10^6$	$7.8 \times 10^7$	$1.1 \times 10^1$	$1.3 \times 10^{-3}$
Pre selection	$2.3 \times 10^7$	$4.7 \times 10^5$	$2.3 \times 10^7$	$1.0 \times 10^1$	$2.1 \times 10^{-3}$
b likeliness > 0.77	$1.4 \times 10^6$	$9.3 \times 10^3$	$1.5 \times 10^6$	$9.4 \times 10^0$	$7.8 \times 10^{-3}$
Emis < 35	$1.3 \times 10^6$	$7.7 \times 10^3$	$1.3 \times 10^6$	$8.4 \times 10^0$	$7.5 \times 10^{-3}$
mvabdt > 0.025	$1.0 \times 10^4$	$2.1 \times 10^2$	$1.0 \times 10^4$	$3.4 \times 10^0$	$3.4 \times 10^{-2}$
-0.92 < cosθ <sub>γ</sub> < 0.92	$5.9 \times 10^3$	$5.7 \times 10^1$	$5.9 \times 10^3$	$3.0 \times 10^0$	$3.9 \times 10^{-2}$

## 5. Event selection



# 5. Signal Significance



Left	2f	4f	total bg	Signal	Significance
Expected	$1.0 \times 10^8$	$3.7 \times 10^7$	$1.4 \times 10^8$	$1.8 \times 10^1$	$3.4 \times 10^{-3}$
Pre selection	$1.3 \times 10^7$	$7.5 \times 10^5$	$1.3 \times 10^7$	$1.0 \times 10^1$	$3.6 \times 10^{-3}$
of charged particle > 3	$7.8 \times 10^4$	$2.3 \times 10^5$	$3.1 \times 10^5$	5.4	$9.8 \times 10^{-3}$
$ m_{W1} - 80.4  < 10 \text{ GeV}$	$2.5 \times 10^4$	$1.6 \times 10^5$	$1.9 \times 10^5$	3.7	$8.6 \times 10^{-3}$
$ m_{W2} - 80.4  < 9.4 \text{ GeV}$					
b likeliness < 0.77	$1.7 \times 10^4$	$1.6 \times 10^5$	$1.8 \times 10^5$	3.7	$8.7 \times 10^{-3}$
mvabdt > 0.1	3.1	$3.8 \times 10^1$	$4.1 \times 10^1$	1.0	$1.6 \times 10^{-1}$
$-0.93 < \cos\theta_\gamma < 0.93$	0.0	8.4	8.4	$9.5 \times 10^{-1}$	$3.1 \times 10^{-1}$

Right	2f	4f	total bg	Signal	Significance
Expected	$7.3 \times 10^7$	$4.6 \times 10^6$	$7.8 \times 10^7$	1.9	$4.8 \times 10^{-4}$
Pre selection	$1.2 \times 10^7$	$3.1 \times 10^5$	$1.2 \times 10^7$	2.0	$3.9 \times 10^{-4}$
of charged particle > 3	$5.0 \times 10^4$	$3.6 \times 10^4$	$8.6 \times 10^4$	$1.5 \times 10^0$	$1.9 \times 10^{-3}$
$ m_{W1} - 80.4  < 10 \text{ GeV}$	$1.7 \times 10^4$	$1.5 \times 10^4$	$3.2 \times 10^4$	$3.8 \times 10^{-1}$	$2.1 \times 10^{-3}$
$ m_{W2} - 80.4  < 9.4 \text{ GeV}$					
b likeliness < 0.77	$1.2 \times 10^4$	$1.4 \times 10^4$	$2.6 \times 10^5$	$3.7 \times 10^{-1}$	$2.3 \times 10^{-3}$
mvabdt > 0.1	$5.3 \times 10^1$	$2.1 \times 10^1$	$7.4 \times 10^1$	$1.0 \times 10^{-1}$	$1.2 \times 10^{-2}$
$-0.93 < \cos\theta_\gamma < 0.93$	0.0	4.7	4.7	$9.3 \times 10^{-2}$	$4.2 \times 10^{-2}$

## 6. cos $\theta_\gamma$ Distribution

We assume no difference between SM and BSM cos  $\theta_\gamma$  distribution  
 → Is this assumption reasonable?

$$\frac{d\sigma}{d \cos \theta} = \frac{d\sigma_{SM}}{d \cos \theta} + \zeta_A \frac{d\sigma_{BSM}}{d \cos \theta} (\zeta_A = 1, \zeta_{AZ} = 0) + \zeta_{AZ} \frac{d\sigma_{BSM}}{d \cos \theta} (\zeta_A = 0, \zeta_{AZ} = 1)$$



$$\text{Efficiency} = \frac{\# \text{ of signal after cut}}{\# \text{ of signal before cut}}$$

There are no cos  $\theta$  dependence

## 7. Combined result - Each polarization

(Pe, Pp)=(-80%, +30%)

95% C.L upper limit

$$\sigma_{h\gamma} = \sigma_{SM} + \frac{1.64}{\text{significance}} \sigma_{SM}$$

H $\rightarrow$ bb

Significance = 0.26 for SM

H $\rightarrow$ WW (Semi-leptonic)

Significance = 0.31 for SM

Combined

Significance = 0.40 for SM

$$\frac{\sigma_{h\gamma}^L}{\sigma_{SM}^L} < 5.1 \quad \sigma_{h\gamma}^L < 1.0 \text{ fb} \quad (95\% \text{ C.L upper limit})$$

(+80%, -30%)

H $\rightarrow$ bb

Significance = 0.039 for SM

H $\rightarrow$ WW (Semi-leptonic)

Significance = 0.042 for SM

Combined

Significance = 0.06 for SM

$$\frac{\sigma_{h\gamma}^R}{\sigma_{SM}^R} < 28.3$$

$$\sigma_{h\gamma}^R < 0.6 \text{ fb} \quad (95\% \text{ C.L upper limit})$$

## 7. Combined result - Each polarization

---

(Pe, Pp)=(-100%, +100%)

95% C.L upper limit

$$\sigma_{h\gamma} = \sigma_{SM} + \frac{1.64}{\text{significance}} \sigma_{SM}$$

**Combined**

Significance = 0.41 for SM

$\sigma^L_{SM}=0.35$  fb for (-100%,+100%)

$$\frac{\sigma_{h\gamma}^L}{\sigma_{SM}^L} < 5.0$$

$$\sigma_{h\gamma}^L < 1.8 \text{ fb}$$

(95% C.L upper limit)

(Pe, Pp)=(+100%, -100%)

**Combined**

Significance = 0.027 for SM

$\sigma^R_{SM}=0.016$  fb for (+100%,-100%)

$$\frac{\sigma_{h\gamma}^R}{\sigma_{SM}^R} < 61.7$$

$$\sigma_{h\gamma}^R < 0.99 \text{ fb}$$

(95% C.L upper limit)

## 8. Constraints from sigma

$\sigma$  limit from experiment

$$\sigma_{hy}^L < 1.8 \text{ fb}$$

$$\sigma_{hy}^R < 0.99 \text{ fb}$$


Constraint on  
 $\zeta_{AZ}$  coupling

Model Independent

Constraint on  
concrete models

Inert Triplet Model

Constraint on  
electron Yukawa coupling



## 8. Constraint on $\zeta_{AZ}$ coupling

---

$$L_{\gamma H} = L_{\text{SM}} + \frac{\zeta_{AZ}}{v} A_{\mu\nu} Z^{\mu\nu} H + \frac{\zeta_A}{2v} A_{\mu\nu} A^{\mu\nu} H$$

Left handed      **Combined**      Significance = 0.41 for SM

$$5.0 > \frac{\sigma_{\gamma H}}{\sigma_{SM}} = 1 - 273\zeta_A - 201\zeta_{AZ} > 0 \quad \text{assume } \zeta_A = 0$$


$$-0.020 < \zeta_{AZ} < 0.005$$

Right handed      **Combined**      Significance = 0.027 for SM

$$61.7 > \frac{\sigma_{\gamma H}}{\sigma_{SM}} = 1 + 492\zeta_A - 311\zeta_{AZ} > 0 \quad \text{assume } \zeta_A = 0$$

$$-0.195 < \zeta_{AZ} < 0.0032$$

## 8. Comparison with LHC




# 8. Constraint on concrete models

$$\frac{\sigma_{h\gamma}^L}{\sigma_{SM}^L} < 5.0$$

(95% C.L. upper limit)

$$\frac{\sigma_{h\gamma}^R}{\sigma_{SM}^R} < 61.7$$

Inert Triplet Model




Kanemura, et al, arXiv:1808.10268

This is the unpolarization plot →

I asked theorist to provide new version of HCOUP, which can calculate cross-section to prepare similar plot for each polarization.

$$\Delta R(e^+e^- \rightarrow h\gamma) = \frac{\sigma(e^+e^- \rightarrow h\gamma)}{\sigma_{SM}(e^+e^- \rightarrow h\gamma)} - 1$$



Kanemura, et al, arXiv:1808.10268



## 8. Constraint on electron Yukawa coupling

The differential cross section is calculated as :

$$\frac{d\sigma_{LL}}{d\cos\theta} = \frac{d\sigma_{RR}}{d\cos\theta} =$$


$$\frac{1}{2s\beta_e} 2e^2 \left(\frac{m_e}{v}\right)^2 D_t D_u s^2 \left[ \left(1 - \frac{m_h^2}{s}\right)^2 t u D_t D_u + 2 \left(\frac{m_h^2}{s}\right) \right] \frac{\bar{\beta}}{16\pi}$$

$$\frac{1}{q^2 - m_e^2} = D_t, \frac{1}{r^2 - m_e^2} = D_u$$



The cross-section is small in the small  $|\cos\theta|$  region.

→ It is very difficult to capture the photon and h in this region because it is the pipeline of the detector.




## 8. cosθ Integration

95% C.L. upper limit

$(e^-, e^+) = (-80, +30)$ ,  $(-80, +30)$

$$\sigma_{hy}^L < 1.0 \text{ fb} \quad \sigma_{hy}^R < 0.6 \text{ fb}$$


$|cos\theta| < 0.92$



$$\sigma_{hy}^L \text{ (Electron-Yukawa)} = 3.6 \times 10^{-7} \text{ [fb]}$$

$$y_e/y_{e(SM)} < 1.6 \times 10^3$$

$$\sigma_{\gamma H} = \sigma_{SM} + \frac{1.64}{\text{significance}} \sigma_{SM}$$



$$\sigma_{hy}^R \text{ (Electron-Yukawa)} = 5.9 \times 10^{-7} \text{ [fb]}$$

$$y_e/y_{e(SM)} < 1.0 \times 10^3$$

## 9. Summary

---

We have performed a full simulation study of  $e^+e^- \rightarrow h\gamma$  at 250 GeV ILC, using ILD detector and full 1-loop SM amplitudes.

Upper limit of  $e^+e^- \rightarrow \gamma h$  at  $\sqrt{s}=250$  GeV,  $900 \text{ fb}^{-1}$

(Left handed)  $\sigma_{h\gamma}^L < 1.0 \text{ fb}$  (95% C.L upper limit)

(Right handed)  $\sigma_{h\gamma}^R < 0.6 \text{ fb}$  (95% C.L upper limit)

Constraint on  $\zeta_{AZ}$  coupling

(Left handed)  $-0.020 < \zeta_{AZ} < 0.005$

(Right handed)  $-0.195 < \zeta_{AZ} < 0.0032$

Constraint on electron-Yukawa coupling

$\sigma_{h\gamma}^L (\text{Electron-Yukawa}) = 3.6 \times 10^{-7} \text{ [fb]}$      $\sigma_{h\gamma}^R (\text{Electron-Yukawa}) = 5.9 \times 10^{-7} \text{ [fb]}$

Future work

Constraint on concrete models (ITM)