



AFB studies at 500 GeV (update)



ILD Top/HF group meeting 25/01/22

A. Irles*, J. P. Márquez*



Orsay/Tohoku/Valencia HQ@ILC Team

AITANA group at IFIC-CSIC/UV



Observables for BSM studies

- Forward-backward asymmetry:

$$A_{FB} = \frac{\int_0^1 \frac{d\sigma}{d \cos \theta} d \cos \theta - \int_{-1}^0 \frac{d\sigma}{d \cos \theta} d \cos \theta}{\int_{-1}^1 \frac{d\sigma}{d \cos \theta} d \cos \theta}$$

- From theory to experiment, i.e., from cross-section to events:

$$N = L \cdot \sigma \cdot \epsilon$$



- Experimental definition:

- Reduce bias from systematic errors from the luminosity and efficiency w.r.t. the cross-section observable

$$A_{FB}^{Exp} = \frac{N_F - N_B}{N_{Total}}$$

Gauge-Higgs Unification (GHU) Models

- The GHU unified all the force carriers under a single gauge group:




- A-Models[1]: Stronger deviations in the right-handed case ($e^-_{RE} e^+_{L} \rightarrow q\bar{q}$).
- B-Models[2]: Stronger deviations in the left-handed case ($e^-_{LE} e^+_{R} \rightarrow q\bar{q}$).
 - The gauge group of GHU is also related to Grand Unification Theory (GUT), embedded in the $SO(11)$ group.

[1] Shuichiro Funatsu, Hisaki Hatanaka, Yutaka Hosotani, Yuta Oriksa, and Naoki Yamatsu. *Fermion pair production at e^-e^+ linear collider experiments in GUT inspired gauge-Higgs unification*. Phys. Rev. D, 102(1):015029, 2020.
[2] Shuichiro Funatsu, Hisaki Hatanaka, Yutaka Hosotani, and Yuta Oriksa. *Distinct signals of the gauge-Higgs unification in e^-e^+ collider experiments*. Phys. Lett. B, 775:297–302, 2017.


Gauge-Higgs Unification (GHU) Models

- A-Model cross-section deviation examples (c-quark) 500 GeV:



Preselection of $q\bar{q}$ signals

- Once we have the reconstructed pfos of the events with different targets:
 - We cluster the signal in jets (VLC algorithm):
 - The algorithm packs together the PFOs into two jets.
 - Signal is expected in a back-to-back topology (but not the backgrounds!)
 - Most of the background is **radiative return (yqq)**
 - And most of the data is background!
 - x3 for $e^-_L e^+_R$ and x6 for $e^-_R e^+_L$ at 250 GeV
 - x4 for $e^-_L e^+_R$ and x7 for $e^-_R e^+_L$ at 500 GeV
 - Then we apply different cuts to the signal to remove the background processes



Summary from the last meeting ($e_L p_R$)


- Cuts:
- $K_{reco} < 20 \text{ GeV}$
 - $m_{2jets} > 250 \text{ GeV}$
 - Charged N pfos > 0.5
 - Neutral N pfos > 4.5
 - Photon veto ($E < 240 \text{ GeV}; |\cos\theta_q| < 0.97$)
 - $y_{23} < 0.006$

- VLC Algorithm parameters:
- $R = 1.0$
 - $\gamma = 0.0$
 - $\beta = 1.0$

	Background/Signal	ISR	WW	ZZ
No Cut	4.044	1.193	0.106	
+ Cut 1	0.871	0.784	0.059	
+ Cut 2	0.863	0.784	0.059	
+ Cut 3	0.243	0.785	0.059	
+ Cut 4	0.182	0.787	0.059	
+ Cut 5	0.134	0.615	0.051	
+ Cut 6	0.102	0.123	0.009	

X2 than at 250 GeV!

Too high!



$K_{reco} < 20 \text{ GeV} \& m_{j_1 j_2} > 250 \text{ GeV} \& NpfosVeto$

$\& Npfos Veto \& Photon Veto 1 \& y_{23} < 0.006$

Efficiency of the preselection for the different quark flavours vs the angular distribution of the two jet system (up to cut 6)


Summary from the last meeting ($e_R p_L$)

- Cuts:
- $K_{reco} < 20 \text{ GeV}$
 - $m_{2\text{jets}} > 250 \text{ GeV}$
 - Charged N pfos > 0.5
 - Neutral N pfos > 4.5
 - Photon veto ($E < 240 \text{ GeV}$; $|\cos\theta_q| < 0.97$)
 - $y_{23} < 0.006$

- VLC Algorithm parameters:
- $R = 1.0$
 - $\gamma = 0.0$
 - $\beta = 1.0$

	Background/Signal		
	ISR	WW	ZZ
No Cut	7.416	0.016	0.127
+ Cut 1	1.597	0.014	0.072
+ Cut 2	1.582	0.014	0.072
+ Cut 3	0.350	0.014	0.072
+ Cut 4	0.227	0.014	0.073
+ Cut 5	0.140	0.015	0.062
+ Cut 6	0.103	0.001	0.013

↑
X2 than at 250 GeV!



$K_{reco} < 20 \text{ GeV} \& m_{j_1 j_2} > 250 \text{ GeV} \& N\text{pfosVeto}$
 $\& N\text{pfos Veto} \& \text{Photon Veto } 1 \& y_{23} < 0.006$

Efficiency of the preselection for the different quark flavours vs the angular distribution of the two jet system (up to cut 6)

Samples (500 GeV)

e_Lp_R

Luminosity (fb^{-1})			
$q\bar{q}$ + ISR	WW	ZZ	$t\bar{t}_1$
46.4	49.0	56.6	7704.9

Cross-Section (fb)			
$q\bar{q}$ + ISR	WW	ZZ	$t\bar{t}_1$
32470.5	7680	680.2	165.0

Luminosity (fb^{-1})			
$q\bar{q}$ + ISR	WW	ZZ	$t\bar{t}_1$
47.0	500	72.5	8354.1

e_Rp_L

Cross-Section (fb)			
$q\bar{q}$ + ISR	WW	ZZ	$t\bar{t}_1$
17994.7	33.5	271.9	63.7

K_{ISR}<20 GeV

	Cross-Section (fb)				
	$b\bar{b}$	$c\bar{c}$	$s\bar{s}$	$u\bar{u}$	$d\bar{d}$
$q\bar{q}$	1051.6	1633.1	1051.5	1643.5	1058.2
ISR	5391.9	4933.3	5389.0	4951.9	5366.6
Ratio	5.1	3	5.1	3	5.1

K_{ISR}<70 GeV

	Cross-Section (fb)				
	$b\bar{b}$	$c\bar{c}$	$s\bar{s}$	$u\bar{u}$	$d\bar{d}$
$q\bar{q}$	1231.3	1917.3	1232.2	1923.5	1239.9
ISR	5212.2	4649.1	5208.4	4671.8	5184.9
Ratio	4.2	2.4	4.2	2.4	4.2

K_{ISR}<20 GeV


	Cross-Section (fb)				
	$b\bar{b}$	$c\bar{c}$	$s\bar{s}$	$u\bar{u}$	$d\bar{d}$
$q\bar{q}$	226.6	733.0	221.7	732.8	224.1
ISR	3233.5	3092.5	3222.5	3075.0	3243
Ratio	14.2	4.2	14.5	4.2	14.5

K_{ISR}<70 GeV


	Cross-Section (fb)				
	$b\bar{b}$	$c\bar{c}$	$s\bar{s}$	$u\bar{u}$	$d\bar{d}$
$q\bar{q}$	264.7	857.9	260.1	857.5	263.7
ISR	3185.4	2967.7	3184.1	2950.3	3203.5
Ratio	12.0	3.5	12.2	3.4	12.1

1: There are 4 different samples (for different processes), this is the average number for each of them

We stepped back to study K_{reco}



We stepped back to study K_{reco}



$K_{\text{reco}} < 20 \text{ GeV}$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	4.04	1.19	0.11	0.11
+ Cut 1	70.9	70.5	70.8	0.87	0.78	0.06	0.007
+ Cut 2	70.8	70.5	70.8	0.86	0.78	0.06	0.007
+ Cut 3	70.7	70.4	70.5	0.24	0.79	0.06	0.007
+ Cut 4	70.7	70.3	70.0	0.20	0.79	0.06	0.007
+ Cut 5	68.5	68.1	67.6	0.14	0.61	0.05	0.007

We stepped back to study K_{reco}

$K_{\text{reco}} < 70 \text{ GeV}$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	3.30	1.19	0.11	0.11
+ Cut 1	94.6	94.3	94.3	0.95	1.07	0.09	0.062
+ Cut 2	94.2	94.1	94.2	0.91	1.07	0.09	0.057
+ Cut 3	94.1	93.9	93.7	0.24	1.07	0.09	0.058
+ Cut 4	94.0	93.8	93.0	0.18	1.07	0.09	0.058
+ Cut 5	90.0	89.6	88.6	0.11	0.69	0.07	0.060

Higher statistic
+
Higher flavour
differences

Similar B/S




We retouched the cuts for $K_{\text{reco}}=70$ ($e_L p_R$)

- Cuts:
- $K_{\text{reco}} < 70 \text{ GeV}$
 - $m_{\text{2jets}} > 250 \text{ GeV}$
 - Charged N pfos > 0.5
 - Neutral N pfos > 3.5
 - Photon veto

- VLC Algorithm parameters:
- $R = 1.0$
 - $\gamma = 0.0$
 - $\beta = 1.0$


	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	3.30	1.02	0.09	0.09
+ Cut 1	94.6	94.3	94.3	0.95	0.92	0.08	0.05
+ Cut 2	94.2	94.1	94.2	0.91	0.91	0.08	0.05
+ Cut 3	94.1	93.9	93.7	0.24	0.91	0.08	0.05
+ Cut 4	94.1	93.8	93.4	0.19	0.91	0.08	0.05
+ Cut 5	90.1	89.7	89.1	0.11	0.59	0.06	0.05

Rebalanced





Now, we carry on the analysis looking in the $e_L p_R$ case

Looking for a sixth cut (y_{23})



Cut $y_{23} < 0.005$ for $K_{\text{reco}} = 20 \text{ & } 70 \text{ GeV}$



Cut $y_{23} < 0.005$ for $K_{\text{reco}} = 20 \text{ & } 70 \text{ GeV}$



$K_{\text{reco}} < 20 \text{ GeV}$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
$y_{23} < 0.005$	48.6	48.7	48.7	0.10	0.09	0.007	0.0001

$K_{\text{reco}} < 70 \text{ GeV}$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
$y_{23} < 0.005$	52.9	52.6	52.6	0.08	0.11	0.010	0.0002

Looking for a seventh cut ($m_{j_1}+m_{j_2}$)



Seventh cut: $m_{j_1} + m_{j_2}$ ($k_{\text{reco}} < 20$ GeV)

$m_{j_1} + m_{j_2} < X$ GeV	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
120	45.9	45.9	45.8	0.10	0.05	0.004	2e-06
130	47.0	47.1	47.1	0.10	0.05	0.004	3e-06
140	47.6	47.8	47.8	0.10	0.06	0.005	5e-06
150	48.0	48.3	48.2	0.10	0.06	0.005	8e-06



$|\cos(\theta)| < 0.9$

$m_{j_1} + m_{j_2} < X$ GeV	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
120	48.4	48.5	48.3	0.10	0.04	0.003	2e-06
130	49.5	49.9	49.7	0.10	0.04	0.004	4e-06
140	50.2	50.6	50.5	0.10	0.05	0.004	6e-06
150	50.6	51.1	50.9	0.10	0.05	0.004	9e-06



Seventh cut: $m_{j_1} + m_{j_2}$ ($\mathbf{k}_{\text{reco}} < 70$ GeV)

$m_{j_1} + m_{j_2} < X$ GeV	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
120	49.7	49.5	49.5	0.08	0.07	0.006	4e-06
130	50.8	50.8	50.9	0.08	0.08	0.007	7e-06
140	51.5	51.6	51.7	0.08	0.08	0.007	9e-06
150	51.9	52.0	52.1	0.10	0.08	0.007	1e-05



$|\cos(\theta)| < 0.9$

$m_{j_1} + m_{j_2} < X$ GeV	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
120	51.7	51.8	51.7	0.08	0.04	0.004	3e-06
130	52.9	53.2	53.1	0.08	0.05	0.005	7e-06
140	53.7	53.9	53.9	0.08	0.05	0.005	8e-06
150	54.1	54.4	54.4	0.08	0.05	0.005	1e-05



Final selection for $K_{\text{reco}}=20 \text{ GeV}$ ($e_L p_R$)


Cuts:

- $K_{\text{reco}} < 20 \text{ GeV}$
- $m_{2\text{jets}} > 250 \text{ GeV}$
- Charged N pfos > 0.5
- Neutral N pfos > 3.5
- Photon veto
- $y_{23} < 0.005$
- $m_{j_1} + m_{j_2} < 140 \text{ GeV}$

VLC Algorithm parameters:

- $R = 1.0$
- $\gamma = 0.0$
- $\beta = 1.0$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	4.04	1.19	0.11	0.11
+ Cut 1	70.9	70.5	70.8	0.87	0.78	0.06	0.007
+ Cut 2	70.8	70.5	70.8	0.86	0.78	0.06	0.007
+ Cut 3	70.7	70.4	70.5	0.24	0.79	0.06	0.007
+ Cut 4	70.7	70.4	70.0	0.20	0.79	0.06	0.007
+ Cut 5	68.5	68.1	67.6	0.14	0.61	0.05	0.007
+ Cut 6	48.6	48.7	48.7	0.10	0.09	0.007	1e-04
+ Cut 7	47.6	47.8	47.8	0.10	0.06	0.005	5e-06



Final selection for $K_{\text{reco}}=20 \text{ GeV}$ ($e_R p_L$)


Cuts:

- $K_{\text{reco}} < 20 \text{ GeV}$
- $m_{2\text{jets}} > 250 \text{ GeV}$
- Charged N pfos > 0.5
- Neutral N pfos > 3.5
- Photon veto
- $y_{23} < 0.005$
- $(m_{j_1} + m_{j_2} < 140 \text{ GeV})$ Not needed

VLC Algorithm parameters:

- $R = 1.0$
- $\gamma = 0.0$
- $\beta = 1.0$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	7.42	0.02	0.13	0.11
+ Cut 1	70.6	70.7	71.0	1.60	0.01	0.07	0.008
+ Cut 2	70.5	70.7	71.0	1.58	0.01	0.07	0.008
+ Cut 3	70.5	70.6	70.7	0.35	0.01	0.07	0.008
+ Cut 4	70.5	70.6	70.5	0.26	0.01	0.07	0.008
+ Cut 5	68.4	68.3	67.9	0.15	0.01	0.06	0.008
+ Cut 6	48.8	48.9	48.6	0.10	5e-04	0.01	8e-05
+ Cut 7	47.7	48.0	47.8	0.10	2e-04	0.007	4e-06



Final selection for $K_{\text{reco}}=70 \text{ GeV}$ ($e_L p_R$)


Cuts:

- $K_{\text{reco}} < 70 \text{ GeV}$
- $m_{2\text{jets}} > 250 \text{ GeV}$
- Charged N pfos > 0.5
- Neutral N pfos > 3.5
- Photon veto
- $y_{23} < 0.005$
- $m_{j_1} + m_{j_2} < 140 \text{ GeV}$

VLC Algorithm parameters:

- $R = 1.0$
- $\gamma = 0.0$
- $\beta = 1.0$

	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	3.30	1.02	0.09	0.09
+ Cut 1	94.6	94.3	94.3	0.95	0.92	0.08	0.05
+ Cut 2	94.2	94.1	94.2	0.91	0.91	0.08	0.05
+ Cut 3	94.1	93.9	93.7	0.24	0.91	0.08	0.05
+ Cut 4	94.1	93.8	93.4	0.19	0.91	0.08	0.05
+ Cut 5	90.1	89.7	89.1	0.11	0.59	0.06	0.05
+ Cut 6	52.6	52.5	52.6	0.08	0.11	0.01	2e-04
+ Cut 7	51.5	51.6	51.7	0.08	0.08	0.007	9e-06



We have to check this weird behavior at high angles with new samples

Final selection for $K_{\text{reco}}=70 \text{ GeV}$ ($e_L p_R$)


Cuts:

- $K_{\text{reco}} < 70 \text{ GeV}$
- $m_{2\text{jets}} > 250 \text{ GeV}$
- Charged N pfos > 0.5
- Neutral N pfos > 3.5
- Photon veto
- $y_{23} < 0.005$
- $(m_{j_1} + m_{j_2} < 140 \text{ GeV})$ Not needed

VLC Algorithm parameters:

- $R = 1.0$
- $\gamma = 0.0$
- $\beta = 1.0$


	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
No cut	100	100	100	6.19	0.01	0.11	0.10
+ Cut 1	93.9	94.4	94.3	1.75	0.01	0.09	0.06
+ Cut 2	93.6	94.3	94.2	1.68	0.01	0.09	0.06
+ Cut 3	93.6	94.0	93.7	0.37	0.01	0.09	0.06
+ Cut 4	93.5	94.0	93.4	0.28	0.01	0.09	0.06
+ Cut 5	89.5	89.9	89.2	0.13	0.01	0.07	0.06
+ Cut 6	52.1	52.6	52.5	0.09	5e-04	0.01	1e-04
+ Cut 7	51.0	51.7	51.6	0.09	3e-04	0.01	5e-06





Back-Up slides

Gauge-Higgs Unification (GHU) Models

- B-Model cross-section deviation examples (b-quark) 500 GeV:




Prospects for b-quark in GHU (A_{FB})




Prospects for c-quark in GHU (A_{FB})

$e^-e^+ \rightarrow c\bar{c}$ (Left-Handed)



$e^-e^+ \rightarrow c\bar{c}$ (Right-Handed)



Efficiency up to cut 6 (previous meeting)

$e_L p_R$

	Efficiency (%)					
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ
Cut 1	70.9	70.5	70.8	15.2	46.5	39.3
+ Cut 2	70.8	70.5	70.8	15.1	46.4	39.3
+ Cut 3	70.7	70.4	70.5	4.2	46.4	39.2
+ Cut 4	70.7	70.3	70.0	3.2	46.3	39.1
+ Cut 5	68.5	68.0	67.3	2.2	34.9	32.5
+ Cut 6	51.1	51.2	50.9	1.3	5.3	4.3

$e_R p_L$

	Efficiency (%)					
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ
Cut 1	70.6	70.7	71.0	15.3	63.8	40.4
+ Cut 2	70.5	70.7	71.0	15.1	63.7	40.3
+ Cut 3	70.5	70.6	70.7	3.3	63.7	40.3
+ Cut 4	70.4	70.6	70.2	2.2	63.6	40.1
+ Cut 5	68.3	68.2	67.6	1.28	62.8	33.0
+ Cut 6	51.4	51.3	50.9	0.7	2.7	5.1

Optimization of the cuts: K_{reco}


- K_{reco} is a good estimator of E_γ :

- Definition of acolinearity:

$$\sin \Psi_{\text{acol}} = \frac{\vec{p}_{j_1} \times \vec{p}_{j_2}}{|\vec{p}_{j_1}| \cdot |\vec{p}_{j_2}|}$$

- Momentum of the collinear photon in the ultrarelativistic limit ($m_{\text{jets}} \ll p_{\text{jets}}$):

$$|\vec{k}| \approx K_{\text{reco}} = \frac{250 \text{ GeV} \cdot \sin \Psi_{\text{acol}}}{\sin \Psi_{\text{acol}} + \sin \theta_1 + \sin \theta_2}$$




Kinematics of a two jets system reconstruction with ISR

Samples (500 GeV)





- The samples names are:
 - $\bar{q}q$ +ISR: 2f_hadronic
 - WW: 4f_WW_hadronic
 - ZZ: 4f_ZZ_hadronic
 - $t\bar{t}$:
 - 6f_ttbar_yycyyc
 - 6f_ttbar_yycyyu
 - 6f_ttbar_yyuuyyc
 - 6f_ttbar_yyuuyyu



Looking for a sixth cut (d_{23})



Looking for a sixth cut (d_{23})





Looking for a sixth cut (d_{23})



d_{23} always induce flavour dependance

Jesús P. Márquez Hernández - ILD Top/HF group meeting 25/01/21



Looking for a sixth cut (y_{23})



Low y_{23} induces a positive slope

Jesús P. Márquez Hernández - ILD Top/HF group meeting 25/01/21



Looking for a sixth cut (y_{23})



High y_{23} barely reduces background

Jesús P. Márquez Hernández - ILD Top/HF group meeting 25/01/21

Cut $y_{23} < 0.02$ for $K_{\text{reco}} = 20 \text{ & } 70 \text{ GeV}$



Cut $y_{23} < 0.02$ for $K_{\text{reco}} = 20 \text{ & } 70 \text{ GeV}$

$K_{\text{reco}} < 20 \text{ GeV}$



	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
$y_{23} < 0.020$	63.7	63.4	63.2	0.13	0.52	0.034	0.001

$K_{\text{reco}} < 70 \text{ GeV}$



	Efficiency (%)			Background/Signal			
	$b\bar{b}$	$c\bar{c}$	$q\bar{q}$	ISR	WW	ZZ	$t\bar{t}$
$y_{23} < 0.020$	74.7	74.5	74.3	0.09	0.52	0.04	0.0034

If we try to remove the slope using a milder cut in y_{23} we left way too much background



Seventh cut: $m_{j_1}+m_{j_2}<120$ GeV



Seventh cut: $m_{j_1}+m_{j_2}<130$ GeV



Seventh cut: $m_{j_1} + m_{j_2} < 140$ GeV



Seventh cut: $m_{j_1}+m_{j_2}<150$ GeV

