

# **15pA561-8**


## **ILCにおける超対称性粒子測定を用いた ミューオンg-2への寄与の再構成の検証**

川田真一、遠藤基、浜口幸一<sup>A</sup>、岩本祥<sup>B</sup>、北原鉄平<sup>C</sup>、諸井健夫<sup>A</sup>、末原大幹<sup>D</sup>  
KEK素核研、東京大理<sup>A</sup>、エトヴェシュ・ロランド大<sup>B</sup>、名古屋大KMI<sup>C</sup>、九州大理<sup>D</sup>


# The International Linear Collider (ILC)

- $e^+e^-$  collider,  $\sqrt{s} = 250$  GeV (upgradable to 500 GeV, 1 TeV)
- polarized beam ( $e^-$ :  $\mp 80\%$ ,  $e^+$ :  $\pm 30\%$ )
- clean environment, known initial state
- matured technology, TDR published

2000  $\text{fb}^{-1}$  @ 250 GeV  
200  $\text{fb}^{-1}$  @ 350 GeV  
4000  $\text{fb}^{-1}$  @ 500 GeV



# Introduction: muon g-2 anomaly



4.2 $\sigma$  discrepancy from the SM prediction  
---> New physics?


Now the discrepancy between the experimental and theoretical values amounts to

$$\Delta a_\mu \equiv a_\mu^{\text{BNL+FNAL}} - a_\mu^{\text{SM}} = (25.1 \pm 5.9) \times 10^{-10}, \quad (5)$$

whose significance is equivalent to  $4.2\sigma$  level, and the muon  $g-2$  anomaly is reconfirmed.<sup>#3</sup>

Many models proposed to explain.  
This talk will pick up the interpretation of [2104.03217]: SUSY interpretation  
(pure-Bino-contribution scenario)


	BLR1	BLR2	BLR3	BLR4
$M_1$	100	100	150	150
$m_L = m_R$	150	150	200	200
$\tan \beta$	5	10	5	10
$\mu$	1323	678	1922	973
$m_{\tilde{\mu}_1}$	154	154	202	202
$m_{\tilde{\mu}_2}$	159	159	207	208
$m_{\tilde{\tau}_1}$	113	113	159	158
$m_{\tilde{\tau}_2}$	190	191	242	243
$m_{\tilde{\nu}_{\mu,\tau}}$	137	136	190	190
$m_{\tilde{\chi}_1^0}$	99	99	150	149
$m_{\tilde{\chi}_2^0}, m_{\tilde{\chi}_3^0}, m_{\tilde{\chi}_1^\pm}$	1323–1324	678–680	1922–1923	973–975
$a_\mu^{\text{SUSY}} \times 10^{10}$	27	27	17	17
$\Omega_{\text{DM}} h^2$	0.120	0.120	0.120	0.120
$\sigma_p^{\text{SI}} \times 10^{47} [\text{cm}^2]$	1.7	3.7	0.8	1.9
$\mu_{\gamma\gamma}$	1.01	1.01	1.01	1.01



$\Omega_{\tilde{\chi}_1^0} = \Omega_{\text{dark matter}} = 0.12$  in this plane  
Can explain muon g-2 with  $1\sigma(2\sigma)$

# Muon g-2 and ILC

## neutralino



At ILC500 (or even at ILC250), we can reconstruct the contribution of this loop-diagram.


Table 2: Observables necessary for the reconstruction of  $a_\mu^{(\text{ILC})}$ , and their uncertainties with  $\sqrt{s} = 500 \text{ GeV}$  and  $\mathcal{L} \sim 500\text{--}1000 \text{ fb}^{-1}$ . Processes relevant to determine each observable are also shown. The second and third rows are the information to determine  $m_{\tilde{\mu}LR}^2$ . For the determination of  $m_{\tilde{\chi}_1^0}$ , analyses of the productions of selectrons and smuons are combined. The uncertainties in  $\tilde{g}_{1,L}^{(\text{eff})}$  are those from the experiment and theory, respectively.

$X$	$\delta X$	$\delta_X a_\mu^{(\text{ILC})}$	Process	
$m_{\tilde{\mu}LR}^2$	12 %	13 %	$e^+e^- \rightarrow \tilde{\tau}^+\tilde{\tau}^-$	(cross section, endpoint)
$(\sin 2\theta_{\tilde{\tau}})$	(9 %)	—	$e^+e^- \rightarrow \tilde{\tau}_1^+\tilde{\tau}_1^-$	(cross section)
$(m_{\tilde{\tau}2})$	(3 %)	—	$e^+e^- \rightarrow \tilde{\tau}_2^+\tilde{\tau}_2^-$	(endpoint)
$m_{\tilde{\mu}1}, m_{\tilde{\mu}2}$	200 MeV	0.3 %	$e^+e^- \rightarrow \tilde{\mu}^+\tilde{\mu}^-$	(endpoint)
$m_{\tilde{\chi}_1^0}$	100 MeV	< 0.1 %	$e^+e^- \rightarrow \tilde{\mu}^+\tilde{\mu}^-/\tilde{e}^+\tilde{e}^-$	(endpoint)
$\tilde{g}_{1,L}^{(\text{eff})}$	a few + 1 %	a few + 1 %	$e^+e^- \rightarrow \tilde{e}_L^+\tilde{e}_R^-$	(cross section)
$\tilde{g}_{1,R}^{(\text{eff})}$	1 %	0.9 %	$e^+e^- \rightarrow \tilde{e}_R^+\tilde{e}_R^-$	(cross section)

**ALL measurable**

# This study: Stau measurement at the ILC

neutralino



Approximately  $\Delta a_\mu^{(\tilde{B})} \propto m_{\tilde{\mu}LR}^2$



Need smuon left-right mixing measurement  
Generally, it is difficult to measure directly,

but we also have:  $m_{\tilde{\mu}LR}^2 = \frac{m_\mu}{m_\tau} m_{\tilde{\tau}LR}^2$



Need stau mass and mixing measurement

# Analysis setup

- ILC500 with BLR1 benchmark point (p4)
- eLpR ( $P(e^-,e^+) = (-0.8,+0.3)$ ) and eRpL ( $P(e^-,e^+) = (+0.8,-0.3)$ ):  $1.6 \text{ ab}^{-1}$  both
- SUSY MC sample production: DELPHES + ILC generic detector card
- SM background (~210M MC events in total)
  - aa\_2f (2-photon process): SGV sample due to huge cross-section but old (~8 years)
  - others: ALL ILD-IDR 500 GeV full simulation samples
- Tau reconstruction: TaJetClustering with default settings

# Statistics (no cuts, 1.6 ab<sup>-1</sup>)

eLpR	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$	SM bkg	aa_2f
No cuts	$4.593 \times 10^4$	$8.570 \times 10^4$	$2.205 \times 10^5$	$1.586 \times 10^5$	$4.314 \times 10^4$	$1.488 \times 10^5$	$4.647 \times 10^4$	$2.621 \times 10^4$	$9.663 \times 10^7$	$4.283 \times 10^9$

eRpL	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$	SM bkg	aa_2f
No cuts	$3.569 \times 10^4$	$8.751 \times 10^5$	$1.852 \times 10^5$	$4.151 \times 10^4$	$1.480 \times 10^5$	$1.386 \times 10^5$	$4.211 \times 10^4$	$2.075 \times 10^4$	$4.727 \times 10^7$	$4.283 \times 10^9$

$O(10^4-10^5)$  stau events vs  $O(10^9)$  SM bkg + aa\_2f

Clearly, we need to design cuts to reject background.

First, we design cuts to select stau-like events and apply veto.

# Design of precuts

- pre1:  $N_{\text{tau}} == 2$
  - pre2:  $E_{\text{tau+}} != 0, E_{\text{tau-}} != 0$  equivalent to require opposite charged tau
  - pre3:  $N_e \text{ in taus} == 0$  reject leptonic events and apply veto both tau->1-prong+no photon mainly for rejecting SUSY background
  - pre4:  $N_{\mu} \text{ in taus} == 0$
  - pre5:  $N_{\text{photon}} \text{ in taus} >= 1$  or  $N_{\text{chargedPFO}} \text{ in taus} >= 3$
  - pre6:  $N_{\text{chargedPFO}} \text{ except tau} <= 1$
  - pre7:  $N_{\text{neutralPFO}} \text{ except tau} <= 5$
- } reject high multiplicity events

# After precuts (1.6 ab<sup>-1</sup>)

eLpR	$\widetilde{e_L}\widetilde{e_L}$	$\widetilde{e_R}\widetilde{e_R}$	$\widetilde{e_L}\widetilde{e_R}$	$\widetilde{\mu_L}\widetilde{\mu_L}$	$\widetilde{\mu_R}\widetilde{\mu_R}$	$\widetilde{\tau}_1^+\widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+\widetilde{\tau}_2^-$	$\widetilde{\tau}_1\widetilde{\tau}_2$	SM bkg	aa_2f
No cuts	$4.593*10^4$	$8.570*10^4$	$2.205*10^5$	$1.586*10^5$	$4.314*10^4$	$1.488*10^5$	$4.647*10^4$	$2.621*10^4$	$9.663*10^7$	$4.283*10^9$
precuts	571.2	1081	2703	234.9	62.47	$2.157*10^4$	$1.340*10^4$	5176	$1.209*10^5$	$3.047*10^7$

eRpL	$\widetilde{e_L}\widetilde{e_L}$	$\widetilde{e_R}\widetilde{e_R}$	$\widetilde{e_L}\widetilde{e_R}$	$\widetilde{\mu_L}\widetilde{\mu_L}$	$\widetilde{\mu_R}\widetilde{\mu_R}$	$\widetilde{\tau}_1^+\widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+\widetilde{\tau}_2^-$	$\widetilde{\tau}_1\widetilde{\tau}_2$	SM bkg	aa_2f
No cuts	$3.569*10^4$	$8.751*10^5$	$1.852*10^5$	$4.151*10^4$	$1.480*10^5$	$1.386*10^5$	$4.211*10^4$	$2.075*10^4$	$4.727*10^7$	$4.283*10^9$
precuts	441.7	$1.081*10^4$	2272	64.01	215.7	$2.004*10^4$	$1.213*10^4$	4128	$7.292*10^4$	$3.047*10^7$

$O(10^3\text{-}10^4)$  stau events vs  $O(10^7)$  SM bkg + aa\_2f  
 Still lots of SM bkg remain, especially aa\_ll.  
 We then design kinematical cuts to reduce backgrounds.

# Design of kinematical cuts

- Cut1:  $\frac{\theta_{\text{acop}}}{\pi} > 0.05$
- Cut2:  $20 < E_{\text{vis}} < 300 \text{ GeV}$
- Cut3:  $M_{\text{inv}} > 200 \text{ GeV}$
- Cut4:  $|\cos \theta_{\text{miss}}| < 0.9$
- Cut5: missing  $P_t > 20 \text{ GeV}$
- Cut6:  $|\cos \theta_{\tau^\pm}| < 0.9$

# After Cuts1-6 (1.6 ab<sup>-1</sup>)

eLpR	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$	SM bkg	aa_2f
No cuts	$4.593 \times 10^4$	$8.570 \times 10^4$	$2.205 \times 10^5$	$1.586 \times 10^5$	$4.314 \times 10^4$	$1.488 \times 10^5$	$4.647 \times 10^4$	$2.621 \times 10^4$	$9.663 \times 10^7$	$4.283 \times 10^9$
precuts	571.2	1081	2703	234.9	62.47	$2.157 \times 10^4$	$1.340 \times 10^4$	5176	$1.209 \times 10^5$	$3.047 \times 10^7$
Cuts1-6	394.1	736.9	1607	176.1	46.85	4456	9457	3397	7681	1764


eRpL	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$	SM bkg	aa_2f
No cuts	$3.569 \times 10^4$	$8.751 \times 10^5$	$1.852 \times 10^5$	$4.151 \times 10^4$	$1.480 \times 10^5$	$1.386 \times 10^5$	$4.211 \times 10^4$	$2.075 \times 10^4$	$4.727 \times 10^7$	$4.283 \times 10^9$
precuts	441.7	$1.081 \times 10^4$	2272	64.01	215.7	$2.004 \times 10^4$	$1.213 \times 10^4$	4128	$7.292 \times 10^4$	$3.047 \times 10^7$
Cuts1-6	322.2	7068	1345	47.32	157.4	4091	8564	2706	1001	1764

$O(10^3)$  stau events vs  $O(10^3)$  SM bkg: similar level!!!  
 Less SM backgrounds in eRpL  
 This time we only focus eRpL.


# Stau measurement (eRpL)

- We assume:
  - Neutralino mass can be precisely determined using other SUSY particle measurement ( $\chi_1^0 = 99.3 \pm 0.1$  GeV)
  - Ignore aa\_2f by using BeamCal information based on previous research (not possible to access in SGV)
- We can obtain directly from the plot
  - Number of events
  - Endpoint of staus
- We then can calculate/reconstruct
  - Stau masses
  - Mixing angle
  - Muon g-2 contribution

Figure: Reconstructed tau energy after all cuts. Higher E(tau) is plotted (tau+ or tau-) to see the endpoints more clearly.




# Event counting



- count number of events in [60 - 150] GeV for SM bkg, SUSY bkg,  $\tilde{\tau}_2\tilde{\tau}_2$ , and  $\tilde{\tau}_1\tilde{\tau}_2$
- $N(\text{SMbkg}) = 595.2$
- $N(\text{SUSY}) = 7215$
- $N(\tilde{\tau}_2\tilde{\tau}_2) = 5803$
- $N(\tilde{\tau}_1\tilde{\tau}_2) = 1354$


# $\tilde{\tau}_2$ endpoint



- **fit SM bkg** using straight line [0]\*x+[1] with the range 130 - 170 GeV with log-likelihood option (assume we can determine SM bkg nicely)
- **fit all** using double straight line [0]\*x+[1]+[2]\*(x-[3]) with the range 136 - 150 GeV
- obtain endpoint [3] from the fit  
 $[3] = 150.4 \pm 1.2 \text{ GeV}$  (theory: 149.9 GeV)

$M_{\tilde{\tau}_2} = 188.6^{+4.9}_{-3.9} \text{ GeV}$   
(model = 189.8 GeV)

# $\tilde{\tau}_1$ endpoint



- fit all using 2nd order polynomial  $[0]*x^2+[1]*x+[2]$  with the range 60 - 120 GeV
- fit all using 2nd order polynomial + straight line  $[0]*x^2+[1]*x+[2]+[3]*(x-[4])$  with the range 40 - 54 GeV
- obtain endpoint [4] from the fit  
 $[4] = 53.19 \pm 0.66$  GeV (theory: 54.5 GeV)



$$M_{\tilde{\tau}_1} = 112.8 \pm 0.2 \text{ GeV}$$

(model = 113.2 GeV)

# Other observables

- Mixing angle:  $\cos \theta_{\tilde{\tau}} = 0.680^{+0.070}_{-0.061}$
- Stau left-right mixing parameter:  $m_{\tilde{\tau}LR}^2 = (1.21^{+0.03}_{-0.12}) \times 10^4 \text{ GeV}^2$
- Smuon left-right mixing parameter:  $m_{\tilde{\mu}LR}^2 = 720^{+17}_{-70} \text{ GeV}^2$
- Muon g-2 contribution:  $a_\mu^{(\tilde{B})} = 26.4^{+2.1}_{-1.7} \times 10^{-10}$  [-7%, +8%]
  - model point:  $a_\mu^{(\tilde{B})} = 27.5 \times 10^{-10}$ ,  $a_\mu^{\text{SUSY}} = 27.1 \times 10^{-10}$

# Summary

- Muon g-2 anomaly is a window to new physics
- SUSY model [2104.03217] can explain this anomaly
- Generated MC samples for realistic estimation at the ILC500
- Designed cuts to reject huge amount of SM background
- We can determine ~0.2%/2.5% for stau1/stau2 masses.
- The mixing angle (~10%), left-right mixing parameter (~10%), and muon g-2 anomaly contribution (~8%) can be determined nicely as well.
- Submitted as a Snowmass White Paper [2203.07056]

# **BACKUP**

# Muon g-2 anomaly + SUSY interpretation

The SUSY contributions to the muon  $g - 2$  can be sizable when at least *three* SUSY multiplets are as light as  $\mathcal{O}(100)$  GeV. They are classified into four types: “WHL”, “BHL”, “BHR”, and “BLR”, where W, B, H, L, and R stand for wino, bino, higgsino, left-handed and right-handed smuons, respectively. Under the mass-insertion approximation, these four types are given as [23]<sup>#4</sup>

$$a_\mu^{\text{WHL}} = \frac{\alpha_2}{4\pi} \frac{m_\mu^2}{M_2 \mu} \tan \beta \cdot f_C \left( \frac{M_2^2}{m_{\tilde{\nu}_\mu}^2}, \frac{\mu^2}{m_{\tilde{\nu}_\mu}^2} \right) - \frac{\alpha_2}{8\pi} \frac{m_\mu^2}{M_2 \mu} \tan \beta \cdot f_N \left( \frac{M_2^2}{m_{\tilde{\mu}_L}^2}, \frac{\mu^2}{m_{\tilde{\mu}_L}^2} \right), \quad (6)$$

$$a_\mu^{\text{BHL}} = \frac{\alpha_Y}{8\pi} \frac{m_\mu^2}{M_1 \mu} \tan \beta \cdot f_N \left( \frac{M_1^2}{m_{\tilde{\mu}_L}^2}, \frac{\mu^2}{m_{\tilde{\mu}_L}^2} \right), \quad (7)$$

$$a_\mu^{\text{BHR}} = -\frac{\alpha_Y}{4\pi} \frac{m_\mu^2}{M_1 \mu} \tan \beta \cdot f_N \left( \frac{M_1^2}{m_{\tilde{\mu}_R}^2}, \frac{\mu^2}{m_{\tilde{\mu}_R}^2} \right), \quad (8)$$

$$a_\mu^{\text{BLR}} = \frac{\alpha_Y}{4\pi} \frac{m_\mu^2 M_1 \mu}{m_{\tilde{\mu}_L}^2 m_{\tilde{\mu}_R}^2} \tan \beta \cdot f_N \left( \frac{m_{\tilde{\mu}_L}^2}{M_1^2}, \frac{m_{\tilde{\mu}_R}^2}{M_1^2} \right), \quad (9)$$

# Two-body decay kinematics (1)

- In the end, we have

$$\begin{aligned} \bullet E^+ &= \frac{\sqrt{s}}{4} \left[ 1 - \left( \frac{m_\chi}{m_{\text{SUSY}}} \right)^2 \right] \left[ 1 + \sqrt{1 - 4 \left( \frac{m_{\text{SUSY}}}{\sqrt{s}} \right)^2} \right] \\ \bullet E^- &= \frac{\sqrt{s}}{4} \left[ 1 - \left( \frac{m_\chi}{m_{\text{SUSY}}} \right)^2 \right] \left[ 1 - \sqrt{1 - 4 \left( \frac{m_{\text{SUSY}}}{\sqrt{s}} \right)^2} \right] \end{aligned}$$

- where  $E^+/E^-$  is the maximum/minimum energy of lepton (electron/positron/muon/tau),  $m_{\text{SUSY}}$  is the mass of SUSY particle (selectron/smugon/stau),  $\sqrt{s} = 500$  GeV in this analysis, and  $m_\chi$  is the neutralino mass and equals to 99 GeV on BLR1 parametrization
- Ignored lepton masses

# Two-body decay kinematics (2)

$\sqrt{s} = 500 \text{ GeV}$ ,  $\widetilde{\chi}_1^0 = 99 \text{ GeV}$ , ignored lepton masses

SUSY particle	mass (GeV)	$E^+$ (GeV)	$E^-$ (GeV)
$\widetilde{e}_L$	157	133.9	16.7
$\widetilde{e}_R$	156	133.0	16.3
$\widetilde{\mu}_L$	158	134.8	17.1
$\widetilde{\mu}_R$	154	131.1	15.6
$\widetilde{\tau}_1$	113	55.0	3.1
$\widetilde{\tau}_2$	190	150.2	31.9

# Produced events (1)

Process $e^+e^- \rightarrow$	Pol (e-, e+) (%)	Xsec (fb)	N = L*Xsec (Assume L = 4 ab <sup>-1</sup> )	N = L*Xsec (Assume L = 1.6 ab <sup>-1</sup> )	N_generated	process ID
$\tilde{e}_L^+\tilde{e}_L^-$	-80/+30	28.7091 +- 0.0012	114836	45935	500K	1
$\tilde{e}_L^+\tilde{e}_L^-$	+80/-30	22.30497 +- 0.00071	89220	35688	500K	2
$\tilde{e}_R^+\tilde{e}_R^-$	-80/+30	53.5626 +- 0.0019	214250	85700	1M	3
$\tilde{e}_R^+\tilde{e}_R^-$	+80/-30	546.909 +- 0.022	2187636	875054	10M	4
$\tilde{\mu}_L^+\tilde{\mu}_L^-$	-80/+30	99.1388 +- 0.0079	396555	158622	1.5M	5
$\tilde{\mu}_L^+\tilde{\mu}_L^-$	+80/-30	25.9426 +- 0.0021	103770	41508	500K	6
$\tilde{\mu}_R^+\tilde{\mu}_R^-$	-80/+30	26.9622 +- 0.0021	107849	43140	500K	7
$\tilde{\mu}_R^+\tilde{\mu}_R^-$	+80/-30	92.4999 +- 0.0072	370000	148000	1.5M	8

1.6 ab<sup>-1</sup> is the integrated luminosity of ILC500 with -80/+30 and +80/-30

# Produced events (2)

Process $e^+e^- \rightarrow$	Pol (e-, e+) (%)	Xsec (fb)	$N = L^*Xsec$ (Assume $L = 4 \text{ ab}^{-1}$ )	$N = L^*Xsec$ (Assume $L = 1.6 \text{ ab}^{-1}$ )	N_generated	process ID
$\tilde{\tau}_1^+\tilde{\tau}_1^-$	-80/+30	92.9890 +- 0.0063	371956	148782	1.5M	9
$\tilde{\tau}_1^+\tilde{\tau}_1^-$	+80/-30	86.6444 +- 0.0059	346578	138631	1.5M	10
$\tilde{\tau}_2^+\tilde{\tau}_2^-$	-80/+30	29.0410 +- 0.0033	116164	46466	500K	11
$\tilde{\tau}_2^+\tilde{\tau}_2^-$	+80/-30	26.3214 +- 0.0029	105286	42114	500K	12
$\tilde{\tau}_1^+\tilde{\tau}_2^-$	-80/+30	8.18989 +- 0.00062	32760	13104	200K	13
$\tilde{\tau}_1^+\tilde{\tau}_2^-$	+80/-30	6.48573 +- 0.00050	25943	10377	200K	14
$\tilde{\tau}_2^+\tilde{\tau}_1^-$	-80/+30	8.19128 +- 0.00062	32765	13106	200K	15
$\tilde{\tau}_2^+\tilde{\tau}_1^-$	+80/-30	6.48553 +- 0.00050	25942	10377	200K	16

$1.6 \text{ ab}^{-1}$  is the integrated luminosity of ILC500 with -80/+30 and +80/-30

# Produced events (3)

Process $e^+e^- \rightarrow$	Pol (e-, e+) (%)	Xsec (fb)	$N = L \times Xsec$ (Assume $L = 4 \text{ ab}^{-1}$ )	$N = L \times Xsec$ (Assume $L = 1.6 \text{ ab}^{-1}$ )	N_generated	process ID
$\tilde{e}_L^+ \tilde{e}_R^-$	-80/+30	23.5750 +- 0.0011	94300	37720	500K	17
$\tilde{e}_L^+ \tilde{e}_R^-$	+80/-30	114.248 +- 0.0051	456992	182797	1.5M	18
$\tilde{e}_R^+ \tilde{e}_L^-$	-80/+30	114.248 +- 0.0051	456992	182797	1.5M	19
$\tilde{e}_R^+ \tilde{e}_L^-$	+80/-30	23.575 +- 0.0011	94300	37720	500K	20

$1.6 \text{ ab}^{-1}$  is the integrated luminosity of ILC500 with -80/+30 and +80/-30

# Potential problem

- The spin information is not stored in stau events
  - This might affect to the decay products of tau
  - It is OK for SM world (e.g.: Keita's study)
  - So far, no special treatment applied

# Physics analysis

- Made everything luminosity-weighted
  - Considered MC statistics
  - $eLpR/eRpL$  for ( $e^-$ ,  $e^+$ ) = (-80%, +30%)/(+80%, -30%)
  - $1.6 \text{ ab}^{-1}$  for both polarization (ILC500 full statistics)
- Included **ALL** available SM background MC samples: in total ~210M MC samples

# SM background (1)

- Added **ALL** available IDR samples
  - `/gpfs/group/ilc/soft/samples/mc-opt-3/ild/dst-merged/500-TDR_ws/PROCESS/ILD_I5_o1_v02/v02-00-01/~~~~~.slcio`
  - processes (h = hadronic, l = leptonic, sl = semileptonic)
    - all 2f (bhabha, h, l)
    - all 4f (singleW\_l/sl, singleZee\_l/sl, singleZnunu\_l/sl, singleZsingleWMix\_l, WW\_h/l/sl, ZZ\_h/l/sl, ZZWWMix\_h/l)
    - all 5f
    - all 6f (eeWW, llWW, ttbar, vvWW, xxWW, xxxxZ, yyyyZ)
    - all aa\_4f
    - all higgs\_ffh (qqh/llhnnh, no specific decays)

# SM background (2)

- Also added **ALL** aa\_2f created by SGV
  - `/ghi/fs02/orig_root_fs02/ilc/grid/storm/users/berggren/mc-dbd/sgv-dst_6/500-TDR_ws/aa_2f/~~~~~.slcio`
  - ~8 years old samples (even used in my PhD thesis)
  - 4 types of processes: aa\_ee, aa\_ll, aa\_xx, aa\_yy
- Since the cross-section is huge, there are no full simulation samples of aa\_2f @ 500 GeV.
- SGV is pretty much faster, but not enough MC samples (event weight ~ 20, which means 1 MC event corresponds to > 20 real events)

# Tau clustering: TaJetClustering

- Originally developed for tau reconstruction under the jet environment
- Treat inclusively, no special treatments for different tau decay
- Used with all default values
  - MinimumJetEnergy = 3 GeV: minimum energy for reconstructed tau
  - MinimumTrackEnergy = 2 GeV: minimum energy for tau seed
  - MinimumTrackEnergyAssoc = 2 GeV: minimum energy for associate particle for tau seed
- This setting might be problematic for  $\tilde{\tau}_1$ 
  - Theoretical  $E_+ = 55.0$  GeV,  $E_- = 3.1$  GeV for  $\tau$ . Its decay products have even lower energy.

# PID information

- Now using `getParticleIDs` instead of `getType`
- In analysis, DELPHES and full simulation samples information are changed to PID information, not `getType` information anymore.
  - DELPHES only have 2 algorithms, picked up higher probability one
  - Full simulation: pick up LikelihoodPID
- SGV can only use PID information (due to old?), but performance of PID maybe not so good.
  - e.g.: 2muons + missing in MC truth, 2pions in PID
  - Only one PID is available

# Statistics (eLpR)

SUSY	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$			
No cuts	$4.593 \times 10^4$	$8.570 \times 10^4$	$2.205 \times 10^5$	$1.586 \times 10^5$	$4.314 \times 10^4$	$1.488 \times 10^5$	$4.647 \times 10^4$	$2.621 \times 10^4$			
SM bkg (1)	Bhabha	2f_I	2f_h	4f_sw_I	4f_sw_sl	4f_sze_I	4f_sze_sl	4f_szn_I	4f_szn_sl	4f_szsw_I	
No cuts	$5.401 \times 10^6$	$5.436 \times 10^6$	$3.140 \times 10^7$	$2.593 \times 10^6$	$7.765 \times 10^6$	$1.144 \times 10^7$	$3.012 \times 10^6$	$2.618 \times 10^5$	$8.941 \times 10^5$	$1.043 \times 10^6$	
SM bkg (2)	4f_WW_h	4f_WW_I	4f_WW_sl	4f_ZZ_h	4f_ZZ_I	4f_ZZ_sl	4f_ZZWW_h	4f_ZZWW_I	5f	eeWW	IIWW
No cuts	$7.191 \times 10^6$	$7.403 \times 10^5$	$8.915 \times 10^6$	$6.519 \times 10^5$	$5.824 \times 10^4$	$5.858 \times 10^5$	$5.995 \times 10^6$	$7.684 \times 10^5$	$1.237 \times 10^5$	$4.612 \times 10^4$	$1.943 \times 10^4$
SM bkg (3)	vvWW	xxWW	xxxxZ	yyyyZ	ttbar	AA4f	AAee	AAll	AAqq	higgs	
No cuts	$3.227 \times 10^4$	$3.650 \times 10^4$	1293	2803	$1.470 \times 10^6$	$3.356 \times 10^5$	$1.146 \times 10^9$	$2.246 \times 10^9$	$8.909 \times 10^8$	$4.123 \times 10^5$	

stau events:  $O(10^4\text{-}10^5)$   
 SUSY background:  $O(10^4\text{-}10^5)$   
 SM background:  $O(10^7)$   
 aa\_2f:  $O(10^9)$

# Precuts (eLpR)

SUSY	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$
No cuts	$4.593 \times 10^4$	$8.570 \times 10^4$	$2.205 \times 10^5$	$1.586 \times 10^5$	$4.314 \times 10^4$	$1.488 \times 10^5$	$4.647 \times 10^4$	$2.621 \times 10^4$
pre1	$4.308 \times 10^4$	$8.028 \times 10^4$	$2.038 \times 10^5$	$1.492 \times 10^5$	$4.068 \times 10^4$	$5.892 \times 10^4$	$3.282 \times 10^4$	$1.350 \times 10^4$
pre2	$4.308 \times 10^4$	$8.028 \times 10^4$	$2.038 \times 10^5$	$1.492 \times 10^5$	$4.068 \times 10^4$	$5.866 \times 10^4$	$3.281 \times 10^4$	$1.346 \times 10^4$
pre3	641.0	1205	3011	$1.492 \times 10^5$	$4.068 \times 10^4$	$3.960 \times 10^4$	$2.273 \times 10^4$	9164
pre4	641.0	1205	3011	433.4	118.2	$2.318 \times 10^4$	$1.400 \times 10^4$	5474
pre5	571.2	1081	2703	234.9	62.47	$2.158 \times 10^4$	$1.341 \times 10^4$	5178
pre6	571.2	1081	2703	234.9	62.47	$2.158 \times 10^4$	$1.340 \times 10^4$	5176
pre7	571.2	1081	2703	234.9	62.47	$2.157 \times 10^4$	$1.340 \times 10^4$	5176

# Precuts (eLpR)

SM bkg (1)	Bhabha	2f_I	2f_h	4f_sw_I	4f_sw_sl	4f_sze_I	4f_sze_sl	4f_szn_I	4f_szn_sl	4f_szsw_I
No cuts	$5.401 \times 10^6$	$5.436 \times 10^6$	$3.140 \times 10^7$	$2.593 \times 10^6$	$7.765 \times 10^6$	$1.144 \times 10^7$	$3.012 \times 10^6$	$2.618 \times 10^5$	$8.941 \times 10^5$	$1.043 \times 10^6$
pre1	$2.605 \times 10^6$	$3.092 \times 10^6$	$8.230 \times 10^4$	$1.495 \times 10^6$	$3.721 \times 10^5$	$1.857 \times 10^6$	$2.549 \times 10^5$	$7.508 \times 10^4$	4212	$6.027 \times 10^5$
pre2	$2.581 \times 10^6$	$3.063 \times 10^6$	$5.775 \times 10^4$	$1.477 \times 10^6$	$2.659 \times 10^5$	$1.624 \times 10^6$	$2.313 \times 10^5$	$7.404 \times 10^4$	2838	$5.948 \times 10^5$
pre3	1665	$1.151 \times 10^6$	2010	$2.451 \times 10^4$	1249	$2.978 \times 10^5$	810	$2.999 \times 10^4$	222.9	668.5
pre4	1382	$1.513 \times 10^5$	750.5	5789	399.1	$1.654 \times 10^4$	316.2	3955	86.74	508.8
pre5	772.5	$1.185 \times 10^5$	460.8	4360	289.7	$1.201 \times 10^4$	228.8	3483	49.66	293.0
pre6	614.9	$1.015 \times 10^5$	0	3733	0	$1.021 \times 10^4$	129.5	2967	12.36	237.2
pre7	546.0	$8.984 \times 10^4$	0	3457	0	6840	50.41	2665	12.36	209.2

2f\_I, 4f\_singleW\_I, 4f\_singleZee\_I, 4f\_singleZnunu\_I:  $O(10^3\text{-}10^4)$   
 semileptonic events:  $< O(10^2)$

# Precuts (eLpR)

SM bkg (2)	4f_WW_h	4f_WW_I	4f_WW_sl	4f_ZZ_h	4f_ZZ_I	4f_ZZ_sl	4f_ZZWW_h	4f_ZZWW_I	5f	eeWW	IIWW
No cuts	$7.191 \times 10^6$	$7.403 \times 10^5$	$8.915 \times 10^6$	$6.519 \times 10^5$	$5.824 \times 10^4$	$5.858 \times 10^5$	$5.995 \times 10^6$	$7.684 \times 10^5$	$1.237 \times 10^5$	$4.612 \times 10^4$	$1.943 \times 10^4$
pre1	$2.850 \times 10^4$	$4.455 \times 10^5$	$4.122 \times 10^5$	2504	$2.155 \times 10^4$	$1.072 \times 10^5$	$2.457 \times 10^4$	$4.682 \times 10^5$	$3.048 \times 10^4$	$1.352 \times 10^4$	5487
pre2	$1.670 \times 10^4$	$4.413 \times 10^5$	$2.915 \times 10^5$	1472	$2.051 \times 10^4$	$1.048 \times 10^5$	$1.477 \times 10^4$	$4.644 \times 10^5$	$2.033 \times 10^4$	9312	4263
pre3	439.8	$1.514 \times 10^5$	$2.758 \times 10^4$	16.53	6990	7648	422.5	$1.766 \times 10^5$	587.5	61.65	217.5
pre4	76.48	1622	3271	16.53	696.2	465.4	115.7	$2.134 \times 10^4$	39.35	5.995	7.175
pre5	76.48	1187	2617	16.53	616.5	401.9	96.54	$1.846 \times 10^4$	30.68	5.215	5.639
pre6	0	1007	18.68	0	510.4	3.744	0	$1.579 \times 10^4$	20.15	3.247	0.8555
pre7	0	928.4	18.68	0	443.1	0	0	$1.456 \times 10^4$	15.54	1.955	0.5277

4f\_WW\_I, 4f\_ZZ\_I, 4f\_ZZWW\_I:  $O(10^3\text{-}10^4)$

semileptonic events:  $< O(10^2)$

hadronic events: 0

# Precuts (eLpR)

SM bkg (3)	vvWW	xxWW	xxxxZ	yyyyZ	ttbar	AA4f	AAee	AAII	AAqq	higgs
No cuts	$3.227 \times 10^4$	$3.650 \times 10^4$	1293	2803	$1.470 \times 10^6$	$3.356 \times 10^5$	$1.146 \times 10^9$	$2.246 \times 10^9$	$8.909 \times 10^8$	$4.123 \times 10^5$
pre1	3849	4200	59.39	500.1	$1.415 \times 10^5$	$7.629 \times 10^4$	$8.937 \times 10^8$	$1.116 \times 10^9$	$5.456 \times 10^6$	$4.070 \times 10^4$
pre2	3510	3612	54.16	400.9	$1.266 \times 10^5$	$5.738 \times 10^4$	$8.923 \times 10^8$	$1.109 \times 10^9$	$3.643 \times 10^6$	$3.875 \times 10^4$
pre3	514.3	123.4	0.9239	16.87	2700	4958	$6.863 \times 10^5$	$1.063 \times 10^9$	$2.529 \times 10^6$	4813
pre4	52.38	4.970	0.1071	0.9115	65.30	280.3	$1.017 \times 10^5$	$1.059 \times 10^9$	$1.473 \times 10^6$	1765
pre5	44.88	4.399	0.05906	0.7031	52.64	230.1	850.5	$3.040 \times 10^7$	$1.123 \times 10^6$	1600
pre6	34.65	0	0	0.2749	0	133.3	850.5	$3.026 \times 10^7$	$3.119 \times 10^5$	1284
pre7	32.39	0	0	0.1283	0	114.2	850.5	$3.021 \times 10^7$	$2.561 \times 10^5$	1150

aa\_ll:  $O(10^7)$

aa\_qq:  $O(10^5)$

6f high multiplicity events: negligible

# Summary of precuts

- Already stau1-pair process is rejected by 64%, still order of  $O(10^4)$  statistics.
  - Due to default setting of TaJetClustering and its lower energy of decay products
- High multiplicity events are now almost negligible.
- SGV-based samples cannot reject by requiring  $N_{(e/\mu\text{-PFO})}$  because such information is not stored in reconstructed PFO. This is maybe due to the performance of PID.
- 209M  $\rightarrow$  1.35M MC events

# Reject more aa\_2f and save stau events

- Stau events:  $O(10^3\text{-}10^4)$  for all channels
- aa\_2f:  $O(10^7)$  at maximum
- Need to design some cuts to reduce the background level

# After Cut7 (eLpR)

SUSY	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$
No cuts	$4.593*10^4$	$8.570*10^4$	$2.205*10^5$	$1.586*10^5$	$4.314*10^4$	$1.488*10^5$	$4.647*10^4$	$2.621*10^4$
precuts	571.2	1081	2703	234.9	62.47	$2.157*10^4$	$1.340*10^4$	5176
Cut1	518.3	982.0	2514	212.8	56.69	$1.703*10^4$	$1.230*10^4$	4536
Cut2	518.3	982.0	2514	212.8	56.69	$1.608*10^4$	$1.229*10^4$	4499
Cut3	518.3	982.0	2514	212.8	56.69	$1.608*10^4$	$1.229*10^4$	4499
Cut4	482.1	909.3	2236	202.2	52.46	$1.475*10^4$	$1.141*10^4$	4141
Cut5	470.1	882.5	2158	198.1	51.51	4798	$1.091*10^4$	3675
Cut6	394.1	736.9	1607	176.1	46.85	4456	9457	3397

# After Cut7 (eLpR)

SM bkg (1)	Bhabha	2f_I	2f_h	4f_sw_I	4f_sw_sl	4f_sze_I	4f_sze_sl	4f_szn_I	4f_szn_sl	4f_szsw_I
No cuts	$5.401 \times 10^6$	$5.436 \times 10^6$	$3.140 \times 10^7$	$2.593 \times 10^6$	$7.765 \times 10^6$	$1.144 \times 10^7$	$3.012 \times 10^6$	$2.618 \times 10^5$	$8.941 \times 10^5$	$1.043 \times 10^6$
precuts	546.0	$8.984 \times 10^4$	0	3457	0	6840	50.41	2665	12.36	209.2
Cut1	189.5	$1.577 \times 10^4$	0	3053	0	5752	50.41	2455	12.36	192.3
Cut2	33.87	3833	0	2454	0	3940	25.69	2438	12.36	154.5
Cut3	3.605	2164	0	1822	0	1986	25.69	2405	12.36	104.0
Cut4	0	778.1	0	1193	0	466.5	0	1892	12.36	48.22
Cut5	0	615.2	0	1125	0	15.85	0	1673	12.36	40.74
Cut6	0	403.9	0	783.2	0	12.94	0	1456	12.36	24.34

hadronic and semileptonic events are now negligible

4f\_singleZnunu\_leptonic: 1456

4f\_singleW\_leptonic: 783.2

2f\_leptonic: 403.9

# After Cut7 (eLpR)

SM bkg (2)	4f_WW_h	4f_WW_I	4f_WW_sl	4f_ZZ_h	4f_ZZ_I	4f_ZZ_sl	4f_ZZWW_h	4f_ZZWW_I	5f	eeWW	IIWW
No cuts	$7.191 \times 10^6$	$7.403 \times 10^5$	$8.915 \times 10^6$	$6.519 \times 10^5$	$5.824 \times 10^4$	$5.858 \times 10^5$	$5.995 \times 10^6$	$7.684 \times 10^5$	$1.237 \times 10^5$	$4.612 \times 10^4$	$1.943 \times 10^4$
precuts	0	928.4	18.68	0	443.1	0	0	$1.456 \times 10^4$	15.54	1.955	0.5277
Cut1	0	757.4	18.68	0	346.8	0	0	$1.153 \times 10^4$	13.98	1.878	0.3952
Cut2	0	542.5	0	0	344.9	0	0	$1.090 \times 10^4$	12.25	1.482	0.3952
Cut3	0	444.8	0	0	323.1	0	0	$1.035 \times 10^4$	11.05	1.359	0.3952
Cut4	0	342.2	0	0	190.1	0	0	7790	5.581	0.5022	0.1360
Cut5	0	332.5	0	0	186.1	0	0	6622	4.350	0.4617	0.06906
Cut6	0	210.4	0	0	147.9	0	0	3859	3.879	0.3587	0.06906

hadronic and semileptonic events are now negligible  
 4f\_ZZWW\_leptonic: 3859

# After Cut7 (eLpR)

SM bkg (3)	vvWW	xxWW	xxxxZ	yyyyZ	ttbar	AA4f	AAee	AAll	AAqq	higgs
No cuts	$3.227 \times 10^4$	$3.650 \times 10^4$	1293	2803	$1.470 \times 10^6$	$3.356 \times 10^5$	$1.146 \times 10^9$	$2.246 \times 10^9$	$8.909 \times 10^8$	$4.123 \times 10^5$
precuts	32.39	0	0	0.1283	0	114.2	850.5	$3.021 \times 10^7$	$2.561 \times 10^5$	1150
Cut1	29.57	0	0	0.1240	0	104.9	277.1	$1.303 \times 10^7$	$7.263 \times 10^4$	1036
Cut2	29.39	0	0	0.1025	0	99.68	259.5	$7.011 \times 10^6$	$4.625 \times 10^4$	1032
Cut3	29.26	0	0	0.1025	0	96.82	216.0	$6.029 \times 10^6$	$3.953 \times 10^4$	1016
Cut4	24.18	0	0	0.02483	0	64.46	35.09	$5.950 \times 10^5$	1242	867.8
Cut5	21.06	0	0	0	0	56.68	0	2788	0	809.4
Cut6	17.30	0	0	0	0	46.03	0	1764	0	703.6

hadronic and semileptonic events are now negligible  
 AA\_ll: 1764

# Statistics (eRpL)

SUSY	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$
No cuts	$3.569 \times 10^4$	$8.751 \times 10^5$	$1.852 \times 10^5$	$4.151 \times 10^4$	$1.480 \times 10^5$	$1.386 \times 10^5$	$4.211 \times 10^4$	$2.075 \times 10^4$

SM bkg (1)	Bhabha	2f_I	2f_h	4f_sw_I	4f_sw_sl	4f_sze_I	4f_sze_sl	4f_szn_I	4f_szn_sl	4f_szsw_I
No cuts	$5.159 \times 10^6$	$4.377 \times 10^6$	$1.866 \times 10^7$	$3.070 \times 10^5$	$9.148 \times 10^5$	$1.131 \times 10^7$	$2.807 \times 10^6$	$2.951 \times 10^4$	$1.085 \times 10^5$	$1.787 \times 10^5$

SM bkg (2)	4f_WW_h	4f_WW_I	4f_WW_sl	4f_ZZ_h	4f_ZZ_I	4f_ZZ_sl	4f_ZZWW_h	4f_ZZWW_I	5f	eeWW	IIWW
No cuts	$4.615 \times 10^5$	$4.818 \times 10^4$	$5.759 \times 10^5$	$2.926 \times 10^5$	$3.784 \times 10^4$	$3.040 \times 10^5$	$4.321 \times 10^5$	$6.372 \times 10^4$	$7.201 \times 10^4$	$1.932 \times 10^4$	1634

SM bkg (3)	vvWW	xxWW	xxxxZ	yyyyZ	ttbar	AA4f	AAee	AAII	AAqq	higgs
No cuts	2489	2947	308.6	1591	$6.372 \times 10^5$	$3.356 \times 10^5$	$1.146 \times 10^9$	$2.246 \times 10^9$	$8.909 \times 10^8$	$1.303 \times 10^5$

stau events:  $O(10^4\text{-}10^5)$   
 SUSY background:  $O(10^4\text{-}10^5)$   
 SM background:  $O(10^7)$   
 aa\_2f:  $O(10^9)$

# Precuts (eRpL)

SUSY	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$
No cuts	$3.569 \times 10^4$	$8.751 \times 10^5$	$1.852 \times 10^5$	$4.151 \times 10^4$	$1.480 \times 10^5$	$1.386 \times 10^5$	$4.211 \times 10^4$	$2.075 \times 10^4$
pre1	$3.358 \times 10^4$	$8.173 \times 10^5$	$1.712 \times 10^5$	$3.906 \times 10^4$	$1.395 \times 10^5$	$5.485 \times 10^4$	$2.970 \times 10^4$	$1.071 \times 10^4$
pre2	$3.358 \times 10^4$	$8.173 \times 10^5$	$1.712 \times 10^5$	$3.906 \times 10^4$	$1.395 \times 10^5$	$5.460 \times 10^4$	$2.970 \times 10^4$	$1.068 \times 10^4$
pre3	495.1	$1.207 \times 10^4$	2530	$3.906 \times 10^4$	$1.395 \times 10^5$	$3.687 \times 10^4$	$2.060 \times 10^4$	7289
pre4	495.1	$1.207 \times 10^4$	2530	113.7	381.0	$2.155 \times 10^4$	$1.268 \times 10^4$	4367
pre5	441.7	$1.081 \times 10^4$	2272	64.01	215.7	$2.005 \times 10^4$	$1.214 \times 10^4$	4129
pre6	441.7	$1.081 \times 10^4$	2272	64.01	215.7	$2.004 \times 10^4$	$1.213 \times 10^4$	4128
pre7	441.7	$1.081 \times 10^4$	2272	64.01	215.7	$2.004 \times 10^4$	$1.213 \times 10^4$	4128

# Precuts (eRpL)

SM bkg (1)	Bhabha	2f_l	2f_h	4f_sw_l	4f_sw_sl	4f_sze_l	4f_sze_sl	4f_szn_l	4f_szn_sl	4f_szsw_l
No cuts	$5.159 \times 10^6$	$4.377 \times 10^6$	$1.866 \times 10^7$	$3.070 \times 10^5$	$9.148 \times 10^5$	$1.131 \times 10^7$	$2.807 \times 10^6$	$2.951 \times 10^4$	$1.085 \times 10^5$	$1.787 \times 10^5$
pre1	$2.464 \times 10^6$	$2.396 \times 10^6$	$4.863 \times 10^4$	$1.368 \times 10^5$	$3.526 \times 10^4$	$1.827 \times 10^6$	$1.900 \times 10^5$	8974	418.0	$7.211 \times 10^4$
pre2	$2.440 \times 10^6$	$2.373 \times 10^6$	$3.410 \times 10^4$	$1.340 \times 10^5$	$2.526 \times 10^4$	$1.598 \times 10^6$	$1.697 \times 10^5$	8873	289.2	$7.028 \times 10^4$
pre3	1721	$8.750 \times 10^5$	993.1	2618	185.4	$2.968 \times 10^5$	763.4	3432	20.80	97.60
pre4	1366	$1.112 \times 10^5$	441.7	621.7	40.66	$1.594 \times 10^4$	352.6	445.5	8.920	75.17
pre5	797.0	$8.347 \times 10^4$	305.3	487.1	30.75	$1.133 \times 10^4$	235.5	378.1	6.702	39.55
pre6	745.5	$7.164 \times 10^4$	0	409.6	0	9728	81.06	330.5	0.7395	32.91
pre7	590.8	$6.333 \times 10^4$	0	379.0	0	6578	33.15	301.2	0.7395	31.23

2f\_l, 4f\_singleZee\_l:  $O(10^3-10^4)$   
 semileptonic events:  $< O(10^2)$

# Precuts (eRpL)

SM bkg (2)	4f_WW_h	4f_WW_I	4f_WW_sl	4f_ZZ_h	4f_ZZ_I	4f_ZZ_sl	4f_ZZWW_h	4f_ZZWW_I	5f	eeWW	IIWW
No cuts	4.615*10 <sup>5</sup>	4.818*10 <sup>4</sup>	5.759*10 <sup>5</sup>	2.926*10 <sup>5</sup>	3.784*10 <sup>4</sup>	3.040*10 <sup>5</sup>	4.321*10 <sup>5</sup>	6.372*10 <sup>4</sup>	7.201*10 <sup>4</sup>	1.932*10 <sup>4</sup>	1634
pre1	1849	2.961*10 <sup>4</sup>	2.739*10 <sup>4</sup>	845.0	1.338*10 <sup>4</sup>	5.363*10 <sup>4</sup>	1796	3.540*10 <sup>4</sup>	1.628*10 <sup>4</sup>	4868	465.1
pre2	1085	2.934*10 <sup>4</sup>	1.928*10 <sup>4</sup>	384.2	1.266*10 <sup>4</sup>	5.233*10 <sup>4</sup>	1099	3.510*10 <sup>4</sup>	1.113*10 <sup>4</sup>	3169	367.7
pre3	26.31	1.014*10 <sup>4</sup>	1895	0.9889	4371	3811	50.03	1.338*10 <sup>4</sup>	509.6	52.43	19.82
pre4	4.576	108.2	231.1	0.9889	380.9	233.3	21.07	1639	30.66	6.164	0.7124
pre5	4.576	78.50	184.5	0.9889	338.7	200.7	19.92	1419	25.90	5.382	0.5849
pre6	0	65.81	1.118	0	275.1	0.2240	0	1212	16.98	3.716	0.1119
pre7	0	61.14	1.118	0	245.4	0	0	1106	13.65	2.519	0.06642

4f\_ZZ\_I, 4f\_ZZWW\_I: O(10<sup>2</sup>-10<sup>3</sup>)

semileptonic events: < O(10)

hadronic events: 0

# Precuts (eRpL)

SM bkg (3)	vvWW	xxWW	xxxxZ	yyyyZ	ttbar	AA4f	AAee	AAII	AAqq	higgs
No cuts	2489	2947	308.6	1591	$6.372 \times 10^5$	$3.356 \times 10^5$	$1.146 \times 10^9$	$2.246 \times 10^9$	$8.909 \times 10^8$	$1.303 \times 10^5$
pre1	296.3	354.3	26.18	358.1	$6.172 \times 10^4$	$7.629 \times 10^4$	$8.937 \times 10^8$	$1.116 \times 10^9$	$5.456 \times 10^6$	$1.771 \times 10^4$
pre2	271.3	308.3	24.18	279.2	$5.536 \times 10^4$	$5.738 \times 10^4$	$8.923 \times 10^8$	$1.109 \times 10^9$	$3.643 \times 10^6$	$1.699 \times 10^4$
pre3	38.72	10.69	0.3570	13.18	1223	4958	$6.863 \times 10^5$	$1.063 \times 10^9$	$2.529 \times 10^6$	1095
pre4	4.170	0.4392	0.02543	0.6309	34.88	280.3	$1.017 \times 10^5$	$1.059 \times 10^9$	$1.473 \times 10^6$	269.6
pre5	3.585	0.3783	0.01577	0.4996	28.96	230.1	850.5	$3.040 \times 10^7$	$1.123 \times 10^6$	227.9
pre6	2.724	0	0	0.1851	0	133.3	850.5	$3.026 \times 10^7$	$3.119 \times 10^5$	153.1
pre7	2.524	0	0	0.09603	0	114.2	850.5	$3.021 \times 10^7$	$2.561 \times 10^5$	133.6

aa\_ll:  $O(10^7)$

aa\_qq:  $O(10^5)$

6f high multiplicity events: negligible

# After Cut7 (eRpL)

SUSY	$\widetilde{e}_L \widetilde{e}_L$	$\widetilde{e}_R \widetilde{e}_R$	$\widetilde{e}_L \widetilde{e}_R$	$\widetilde{\mu}_L \widetilde{\mu}_L$	$\widetilde{\mu}_R \widetilde{\mu}_R$	$\widetilde{\tau}_1^+ \widetilde{\tau}_1^-$	$\widetilde{\tau}_2^+ \widetilde{\tau}_2^-$	$\widetilde{\tau}_1 \widetilde{\tau}_2$
No cuts	$3.569 \times 10^4$	$8.751 \times 10^5$	$1.852 \times 10^5$	$4.151 \times 10^4$	$1.480 \times 10^5$	$1.386 \times 10^5$	$4.211 \times 10^4$	$2.075 \times 10^4$
precuts	441.7	$1.081 \times 10^4$	2272	64.01	215.7	$2.004 \times 10^4$	$1.213 \times 10^4$	4128
Cut1	397.7	9912	2114	57.53	194.7	$1.581 \times 10^4$	$1.113 \times 10^4$	3616
Cut2	397.7	9912	2114	57.53	194.7	$1.493 \times 10^4$	$1.112 \times 10^4$	3584
Cut3	397.7	9912	2114	57.53	194.7	$1.493 \times 10^4$	$1.112 \times 10^4$	3584
Cut4	374.9	9058	1874	55.04	182.9	$1.369 \times 10^4$	$1.032 \times 10^4$	3301
Cut5	365.3	8764	1806	53.71	178.8	4396	9868	2930
Cut6	322.2	7068	1345	47.32	157.4	4091	8564	2706

# After Cut7 (eRpL)

SM bkg (1)	Bhabha	2f_l	2f_h	4f_sw_l	4f_sw_sl	4f_sze_l	4f_sze_sl	4f_szn_l	4f_szn_sl	4f_szsw_l
No cuts	$5.159 \times 10^6$	$4.377 \times 10^6$	$1.866 \times 10^7$	$3.070 \times 10^5$	$9.148 \times 10^5$	$1.131 \times 10^7$	$2.807 \times 10^6$	$2.951 \times 10^4$	$1.085 \times 10^5$	$1.787 \times 10^5$
precuts	590.8	$6.333 \times 10^4$	0	379.0	0	6578	33.15	301.2	0.7395	31.23
Cut1	342.2	$1.113 \times 10^4$	0	341.0	0	5624	33.15	277.4	0.7395	30.22
Cut2	68.73	2239	0	288.1	0	3813	12.69	274.5	0.7395	22.54
Cut3	17.47	1259	0	217.0	0	1949	12.69	252.1	0.7395	11.66
Cut4	0	396.5	0	111.9	0	509.6	0	171.0	0.7395	5.024
Cut5	0	341.6	0	106.4	0	11.95	0	156.1	0.7395	4.576
Cut6	0	204.8	0	62.70	0	8.851	0	135.6	0.7395	1.456

hadronic and semileptonic events are now negligible

2f\_leptonic: 204.8

4f\_singleZnunu\_leptonic: 135.6

# After Cut7 (eRpL)

SM bkg (2)	4f_WW_h	4f_WW_I	4f_WW_sl	4f_ZZ_h	4f_ZZ_I	4f_ZZ_sl	4f_ZZWW_h	4f_ZZWW_I	5f	eeWW	IIWW
No cuts	$4.615 \times 10^5$	$4.818 \times 10^4$	$5.759 \times 10^5$	$2.926 \times 10^5$	$3.784 \times 10^4$	$3.040 \times 10^5$	$4.321 \times 10^5$	$6.372 \times 10^4$	$7.201 \times 10^4$	$1.932 \times 10^4$	1634
precuts	0	61.14	1.118	0	245.4	0	0	1106	13.65	2.519	0.06642
Cut1	0	50.91	1.118	0	184.5	0	0	894.7	12.35	2.356	0.05850
Cut2	0	38.05	0	0	184.3	0	0	851.7	11.01	1.752	0.05850
Cut3	0	32.21	0	0	175.1	0	0	804.0	9.474	1.538	0.05850
Cut4	0	26.07	0	0	113.9	0	0	604.1	4.952	0.5664	0.03403
Cut5	0	25.49	0	0	111.7	0	0	512.2	4.299	0.5259	0.02680
Cut6	0	18.18	0	0	85.77	0	0	339.1	3.764	0.4816	0.02680

hadronic and semileptonic events are now negligible  
 4f\_ZZWW\_leptonic: 339.1

# After Cut7 (eRpL)


SM bkg (3)	vvWW	xxWW	xxxxZ	yyyyZ	ttbar	AA4f	AAee	AAII	AAqq	higgs
No cuts	2489	2947	308.6	1591	$6.372 \times 10^5$	$3.356 \times 10^5$	$1.146 \times 10^9$	$2.246 \times 10^9$	$8.909 \times 10^8$	$1.303 \times 10^5$
precuts	2.524	0	0	0.09603	0	114.2	850.5	$3.021 \times 10^7$	$2.561 \times 10^5$	133.6
Cut1	2.317	0	0	0.04830	0	104.9	277.1	$1.303 \times 10^7$	$7.263 \times 10^4$	125.8
Cut2	2.301	0	0	0.03463	0	99.68	259.5	$7.011 \times 10^6$	$4.625 \times 10^4$	123.6
Cut3	2.287	0	0	0.03463	0	96.82	216.0	$6.029 \times 10^6$	$3.953 \times 10^4$	119.9
Cut4	1.901	0	0	0.001486	0	64.46	35.09	$5.950 \times 10^5$	1242	108.2
Cut5	1.692	0	0	0	0	56.68	0	2788	0	104.7
Cut6	1.415	0	0	0	0	46.03	0	1764	0	91.87

hadronic and semileptonic events are now negligible  
 AA<sub>II</sub>: 1764

# **Stau Measurement (eLpR)**


All plots are reconstructed higher tau energy between tau+ and tau- to see the endpoint more clearly.

# Event counting



- count number of events with  $[60 - 150]$  GeV for SM bkg, SUSY bkg,  $\tilde{\tau}_2\tilde{\tau}_2$ , and  $\tilde{\tau}_1\tilde{\tau}_1$
- $N(\text{SMbkg}) = 4873$
- $N(\text{SUSY}) = 2365$
- $N(\tilde{\tau}_2\tilde{\tau}_2) = 6413$
- $N(\tilde{\tau}_1\tilde{\tau}_1) = 1705$


# $\tilde{\tau}_2$ endpoint



- fit SM bkg using straight line  $[0]*x+[1]$  with the range 130 - 170 GeV (assume we can determine SM bkg nicely)
- fit all using double straight line  $[0]*x+[1]+[2]*(x-[3])$  with the range 136 - 150 GeV
- obtain endpoint [3] from the fit  
 $[3] = 149.5 \pm 1.7 \text{ GeV}$

$M_{\tilde{\tau}_2} = XXX \pm XXX \text{ GeV}$   
(model = 189.8 GeV)

# $\tilde{\tau}_1$ endpoint



- fit all using 2nd order polynomial  $[0]*x^2+[1]*x+[2]$  with the range 60 - 120 GeV
  - fit all using 2nd order polynomial + straight line  $[0]*x^2+[1]*x+[2]+[3]*(x-[4])$  with the range 40 - 54 GeV
  - obtain endpoint [4] from the fit
- $[4] = 51.73 \pm 0.53 \text{ GeV}$



$$M_{\tilde{\tau}_1} = XX \pm XX \text{ GeV}$$

(model = 113.2 GeV)