


arXiv:1912.08403


arXiv:2203.07668

Measuring the tau polarisation at ILC

Keita Yumino, Daniel Jeans



KEK, SOKENDAI



Motivation 1

At the ILC, forward-backward asymmetry $A_{FB} = \frac{3}{4} A_e \cdot A_f$ can be measured

Left- and right-handed coupling g_R, g_L to Z boson are different

⇒ Left- and right-handed polarisation asymmetry is expected.

$$A_f = \frac{g_R^2 - g_L^2}{g_R^2 + g_L^2}$$

Thanks to ILC's polarised beams, A_e can be measured

⇒ A_f can be extracted from A_{FB}

By measuring A_{FB} precisely and looking for deviations from SM predictions, it is possible to search for new physics, such as heavy gauge boson Z'

Motivation 2

Tau has extra information

tau is the only particle that can measure the polarisation of the final state in the ILC250

We can also directly measure A_τ by using tau polarisation $P(\tau)$

$$\frac{dP(\tau)}{d \cos \theta} = \frac{3}{8} A_\tau (1 + \cos^2 \theta) + \frac{3}{4} A_e \cos \theta$$

The aim of this study


The reconstruction of tau spin orientation (“**Polarimeter**”)
in order to measure polarisation to investigate new physics.

Polarimeter

Reconstruction of tau polarisation $P(\tau)$ depends on tau decay mode.

Polarimeter vectors of $\tau \rightarrow \pi\nu$ in τ rest frame

$$h(\tau^\pm \rightarrow \pi^\pm\nu) \propto p_{\pi^\pm}$$



Polarimeter vectors of $\tau \rightarrow \rho\nu$ in τ rest frame

$$h(\tau^\pm \rightarrow \pi^\pm\pi^0\nu) \propto m_\tau(E_{\pi^\pm} - E_{\pi^0})(p_{\pi^\pm} - p_{\pi^0}) + \frac{1}{2}(p_{\pi^\pm} + p_{\pi^0})^2 p_\nu$$

“Polarimeter”



The cosine of the angle this polarimeter vector makes to the tau flight direction

Previous study

Extract polarimeter without using neutrino information

"Approximate" polarimeters based only on the momenta of visible tau decay products

"Optimal" polarimeters including the neutrino component



We explicitly extract the neutrino momentum and reconstruct polarimeters

Simulation setup

- Signal event sample with 100 % $e_L^- e_R^+$ beam polarisations were generated using WHIZARD ver 2.8.5.
- The decay of the polarised tau was done using TAUOLA.
- MC truth information was used.


currently

- only look at

$$\tau \rightarrow \pi\nu \quad (\text{BR} \sim 10 \%)$$

$$\tau \rightarrow \rho\nu \quad (\text{BR} \sim 26 \%)$$


τ reconstruction method



Assume

- Two taus are produced along the beam line ($x = y = 0$),
- Two taus are back-to-back in x-y plane,
 - any ISR photons have negligible p_T
- Charged particle travels along a straight line.


τ reconstruction method



Assume

- Primary interaction occurs along the beam line($x = y = 0$),
 - Two taus are back-to-back in x-y plane,
 - Charged particle travels along a straight line, and
 - Two taus produced at somewhere on z-axis.
- Two taus momenta lie in a plane containing z-axis defined by azimuthal angle ϕ


τ reconstruction method



★ The intersection between plane and trajectory : the decay points of τ

For a plane with given azimuthal angle ϕ ,
the intersection of trajectories with this plane can be calculated .


τ reconstruction method



then choice of z_{IP} gives direction of tau momenta

\Rightarrow How can we choose ϕ, z_{IP} ?

τ reconstruction method



(ϕ, z_{IP}) determined tau direction

with assumptions

- $p_T^{\tau_1} = p_T^{\tau_2}$
- Single ISR photon
- E_{CM}


\Rightarrow tau 4-momentum

By comparing with the visible tau 4-momentum
 \Rightarrow the invariant mass of the missing (neutrino) momentum for each tau can be calculated



We choose the values of z and ϕ which result in neutrino masses closest to zero

Find solutions

We choose the values of z and ϕ which result in neutrino masses closest to zero





Find solutions



Impact parameter method efficiency is $> 80\%$ for events with $m_{\tau\tau} \sim 250$ GeV

Comparison with MC

Reconstructed $m_{\tau\tau}$ based on new method solutions




Reasonable agreement between MC and reconstructed $m_{\tau\tau}$

Polarimeter


Polarimeter vectors of $\tau \rightarrow \pi\nu$ in τ rest frame

$$h(\tau^\pm \rightarrow \pi^\pm\nu) \propto p_{\pi^\pm}$$



$\tau \rightarrow \rho\nu$ in τ rest frame

$$m_\tau(E_{\pi^\pm} - E_{\pi^0})(p_{\pi^\pm} - p_{\pi^0}) + \frac{1}{2}(p_{\pi^\pm} + p_{\pi^0})^2 p_\nu$$



Polarimeter

Polarimeter vectors of $\tau \rightarrow \pi\nu$ in τ rest frame



$\tau \rightarrow \rho\nu$ in τ rest frame

$$h(\tau^\pm \rightarrow \pi^\pm\nu) \propto p_{\pi^\pm}$$

$$m_\tau(E_{\pi^\pm} - E_{\pi^0})(p_{\pi^\pm} - p_{\pi^0}) + \frac{1}{2}(p_{\pi^\pm} + p_{\pi^0})^2 p_\nu$$

$\tau \rightarrow \pi\nu$ New method vs MC eLpR(100%)

$\tau \rightarrow \rho\nu$ New method vs MC eLpR(100%)



Polarimeter using reconstructed ν is in reasonable agreement with MC one.

Summary

- The reconstruction of neutrino momentum at ILC-250 was investigated.
- For events with $m_{\tau\tau} \sim 250$ GeV, new method efficiency is $> 80\%$.
- Polarimeters were reconstructed in the $\tau \rightarrow \pi\nu$ and $\tau \rightarrow \rho\nu$ decay modes, and used to estimate the tau polarisation.
- Reasonable agreement between MC truth polarimeter value and the one from “Impact parameter method” for both $\tau \rightarrow \pi\nu$ and $\tau \rightarrow \rho\nu$ decay were found.



Future plan

- Investigate search for new physics by using the tau polarisation.
- Investigate the effect of full detector simulation and reconstruction.
- Quantify the precision with which the tau polarisation can be measured at ILC-250.
- Understand the structure of the method's efficiency around the Z peak.
- Consider how to handle up to four possible solutions that can be found.

Future plan

- Investigate search for new physics by using the tau polarisation.
- Investigate the effect of full detector simulation and reconstruction.
- Quantify the precision with which the tau polarisation can be measured at ILC-250.
- Understand the structure of the method's efficiency around the Z peak.
- Consider how to handle up to four possible solutions that can be found.

Find solutions



Efficiency improved even for low $m_{\tau\tau}$ region



There is an interesting structure around Z peak which is not yet fully understood

Future plan

- Investigate search for new physics by using the tau polarisation.
- Investigate the effect of full detector simulation and reconstruction.
- Quantify the precision with which the tau polarisation can be measured at ILC-250.
- Understand the structure of the method's efficiency around the Z peak.
- Consider how to handle up to four possible solutions that can be found.

Future plan




Comparison with MC



The reconstructed direction is typically within a few mrad of the true direction.

reasonable agreement between MC and reconstructed tau


π polarimeter



Correlations are found in 4 possible solutions.

Consider how to handle these solutions to extract the polarimeter

on going



For a given (ϕ, z_{IP}) ,
 know two tau's momentum direction

- visible daughter momentum
- tau mass constraint
- 1-neutrino assumption


⇒ 2 solutions for each tau's energy

⇒ 4 solutions / event

τ reconstruction method


Assume

- 1 neutrino per tau
- $m_\tau = 1.776 \text{ GeV}$
- $E_\tau = \frac{E_{cm}}{2}$



from these assumptions, τ direction must make an angle β to the visible τ momentum

$$\cos \beta = f(\vec{P}_{\text{vis}}^\tau, m_\tau, E_\tau)$$



$\vec{P}_{\text{vis}}^\tau$: tau visible daughter momentum

\vec{P}_ν : neutrino momentum

\vec{P}_τ : tau momentum

The same for the other tau


τ reconstruction method

We further assume

- two taus are back-to-back

To reconstruct tau momentum, flip one of the cones and find the **intersections**.




The intersection of the cones are the candidate τ momentum directions.



We call this “Cone method”

Find solutions

If at least one intersection point was found, there is a solution.




Cone method efficiency is $\sim 80\%$ for events with $m_{\tau\tau} \sim 250$ GeV

Midpoint method

For events for which “Cone Method” cannot find a solution

$$\vec{P}_{vis}^{\tau^+}$$



take a midpoint of the closest approach points of the two cone edges

and use this **new vector** as a solution

We call this “Midpoint method”

Various levels of “cheating” and methods


Two levels of cheating

1. Using true neutrino momentum from MC.
2. Using true MC visible tau daughters.
 - 2.1 “Cone method” to estimate the neutrino momentum.
 - 2.2 If Cone method fails, “Midpoint method”


Polarimeter: single pi decay

Polarimeter vectors of $\tau \rightarrow \pi\nu$ in τ rest frame $h(\tau^\pm \rightarrow \pi^\pm\nu) \propto p_{\pi^\pm}$

true vs Cone method



true vs Midpoint method



Polarimeter using reconstructed ν is in reasonable agreement with MC one.

Cone method works better than Midpoint method.



Polarimeter: rho decay

Polarimeter vectors of $\tau \rightarrow \rho\nu$ in τ rest frame

$$h(\tau^\pm \rightarrow \pi^\pm \pi^0 \nu) \propto m_\tau (E_{\pi^\pm} - E_{\pi^0})(p_{\pi^\pm} - p_{\pi^0}) + \frac{1}{2}(p_{\pi^\pm} + p_{\pi^0})^2 p_\nu$$

true vs Cone method

true vs Midpoint method




Extracted polarimeter is less precise than $\tau \rightarrow \pi\nu$

Tau Polarisation Accuracy

Q. How accurately can tau polarisation be measured?

Reconstructed polarimeter templates for $\tau \rightarrow \pi\nu$ and $\tau \rightarrow \rho\nu$ decays.

Scaled to the integrated luminosity of 1000 fb^{-1} with 100% $e_L^- e_R^+$.



Polarisation

$$P = \frac{N_R - N_L}{N_R + N_L}$$

Use likelihood method to analytically estimate polarisation uncertainty.

Tau Polarisation Accuracy

Scaled to the luminosity of 1000 fb^{-1}

Sample with 100 % $e_L^- e_R^+$ beam polarisations

N_τ : the expected total number of taus, σ_P : the expected polarisation uncertainty

$\tau \rightarrow \pi\nu$	N_τ	σ_P
MC	0.58 M	0.27 %
Cone	0.36 M	0.35 %
Mid	0.22 M	0.55 %
Combined	0.58 M	0.30 %

$\tau \rightarrow \rho\nu$	N_τ	σ_P
MC	1.31 M	0.18 %
Cone	0.70 M	0.28 %
Mid	0.59 M	0.42 %
Combined	1.29 M	0.23 %

Precision on the polarisation σ_P of “Cone method” + “Midpoint method”

$$\tau \rightarrow \pi\nu : \sim 0.30 \% , \quad \tau \rightarrow \rho\nu : \sim 0.23 \%$$