


Exploring Right Handed Neutrinos at ILC500

Work in progress

Jurina Nakajima, Daniel Jeans^A, Arindam Das^B, Keisuke Fujii^A,
Nobuchika Okada^C, Satomi Okada^C, Ryo Yonamine^A

SOKENDAI, KEK^A, Hokkaido Univ.^B, Arabama Univ.^C



Motivation and introduction

The Right Handed Neutrino (RHN) can address the following big questions

- Why does matter dominate anti-matter in our universe?
- Do quarks and leptons unify?
- Why is neutrino mass so small?


RHN is assumed to be

[arXiv\[1812.11931\]](https://arxiv.org/abs/1812.11931)

- a **Majorana** particle ($N = \bar{N}$)
- minimal $U(1)_{B-L}$ model

→ RHN **pair** production

$$G_{B-L} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L} \rightarrow \text{gauge boson : } Z'$$



Benchmark points with $M_N = 100, 150, 200, 225$ GeV

Benchmark points

- Pol(e⁻, e⁺) = (-0.8, +0.3), (+0.8, -0.3): $\mathcal{L} = 1600$ [fb⁻¹]
- Pol(e⁻, e⁺) = (-0.8, -0.3), (+0.8, +0.3): $\mathcal{L} = 400$ [fb⁻¹]


M_N [GeV] RHN mass	$M_{Z'}$ [TeV] Z' mass	$g_{1'}$ U(1) _{B-L} coupling	$ V_{eN} ^2$ mixing angle	$\sigma_0(e_L^- e_R^+ \rightarrow NN)$ 100% polarization [fb]	BR ($N \rightarrow e^+ W^-$)	Event # at ILC500 [4000fb ⁻¹]
100	7	1	0.0009	0.55	0.44	1446
150	7	1	0.0009	0.36	0.33	925
200	7	1	0.0009	0.14	0.30	349
225	7	1	0.0009	0.046	0.29	112

► minimal U(1)_{B-L} model

► ILC 500 with initial state radiation (ISR) and beamstrahlung (BS)


Current limits - Z' mass

SM like Z' coupling



ATLAS-TDR-LHCC2017-2018


RHN pair production crosssection at ILC250 for expected HL-LHC limits on $M_{Z'}/g'$



arXiv[1812.11931]

The heavier Z' mass less constrained by LHC

Same sign cross-section vs M_N




$$\begin{aligned} \text{cross - section} &= \sigma(ee \rightarrow NN \rightarrow e^\pm e^\pm W^\mp W^\mp) \\ &= \sigma_0(ee \rightarrow NN) \times 2(BR(N \rightarrow e^- W^+))^2 \end{aligned}$$

Signal process and analysis tools


ILC500

Signal process:



s-channel via Z'

+ (Destructive interference)



t-channel via W
(only e_L)

UFO model files

WHIZARD ver 2.8.5

Make Events

ILD Full Simulation
& (Geant4)

Reconstruction

miniDST

Events format


Background samples

We consider 6f and 4f background samples
All processes with at least 1 electron and 2 quarks


- eeqqqq
- 6f_ttbar
- 4f_singleW_semileptonic
- 4f_singleZee_semileptonic

IDR samples
ILC500
miniDST


4 fermions singleW
semileptonic



4 fermions singleZee
semileptonic



6 fermions ttbar
1electron





Cut conditions to select signal events

Related to
isolated electron

- ▶ 2 isolated electron && 0 isolated γ , μ
- ▶ **Same sign isolated electrons**
- ▶ Isolated electron energies $E_{iso} < 200$ [GeV]
- ▶ Isolated electron polar angles $|\cos\theta_{isoel}| < 0.95$
- ▶ **IsolatedLepTagging(min) > 0.9**
 - ▶ Jet clustering with Durham $\log_{10}(y_{12}) > -1$
 - ▶ $P_{miss} < 100$ [GeV] && ($P_{miss} < 40$ [GeV] || $|\cos\theta_{Pmiss}| > 0.95$)

Electron Charge


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$



We use only same sign samples $e_1 \times e_2 = 1$

Distribution of IsolatedLepTagging

- ILC 500 with ISR / BS
- **Pol(e⁻, e⁺) = (+0.8, -0.3)**
- Isolated e # = 2 && Isolated γ # = 0 && Isolated μ # = 0
- Isolated e is same sign (e₁ × e₂ = 1)



Isolated lepton tagging
... output parameter of MVA
to identify isolated lepton

→ Output for e is **near 1**

IsolatedLepTagging_{min} > 0.9

Cut flow (eLpR)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$ $\mathcal{L} = 1600 [\text{fb}^{-1}]$

ILD work in progress

	Signal Events ($ee \rightarrow NN$)				Background Events					
	$M_N=100$	$M_N=150$	$M_N=200$	$M_N=225$	eeqqqq	4f_singleW_ semileptonic	4f_singleZee_ semileptonic	6f_ttbar 2electrons	6f_ttbar 1electron	6f_ttbar 0electron
No cut	554	394	143	45	11898	2825010	699475	16425	129283	11028
$e_{\text{iso}} \# = 2 \ \&\&$ $\gamma_{\text{iso}} \# = 0 \ \&\&$	347	343	79	40	4721	90818	162774	9422	2271	201
Same sign ($e_{\text{iso}1} \times e_{\text{iso}2} = 1$)	176	115	39	12	39	46138	3800	8	439	25
$E_{\text{iso}} < 200$ [GeV]	175	114	39	12	39	41319	3557	8	439	25
$-0.95 <$ $\cos\theta_{\text{iso}e} < 0.95$	156	103	36	11	13	17506	623	4	266	15
IsolatedLepTa gging _{min} > 0.9	94	91	31	10	2	2632	128	1	50	0
$\log_{10}(y_{12}) > -1$	94	90	31	9	2	2632	128	1	50	0
$P_{\text{miss}} < 100 \ \&\&$ ($P_{\text{miss}} < 40 \ \parallel$ $ \cos\theta_{P_{\text{miss}}} >$ 0.95)	84	84	28	9	1	79	30	0	9	0

Cut flow (eLpR)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$ $\mathcal{L} = 1600 [\text{fb}^{-1}]$


ILD work in progress

	Signal Events ($ee \rightarrow NN$)				Background Events					
	$M_N=100$	$M_N=150$	$M_N=200$	$M_N=225$	eeqqqq	4f_singleW _semileptonic	4f_singleZee _semileptonic	6f_ttbar 2electrons	6f_ttbar 1electron	6f_ttbar 0electron
No cut	554	394	143	45	11898	2825010	699475	16425	129283	11028
$e_{\text{iso}} \# = 2 \ \&\&$ $\gamma_{\text{iso}} \# = 0 \ \&\&$	347	343	79	40	4721	90818	162774	9422	2271	201
Same sign ($e_{\text{iso}1} \times e_{\text{iso}2} = 1$)	15	15	5	1	15	17500	625	4	189	25
$E_{\text{iso}} < 200$ [GeV]	15	15	5	1	15	17500	625	4	189	25
$-0.95 <$ $\cos\theta_{\text{iso}e} < 0.95$	15	15	5	1	15	17500	625	4	189	15
IsolatedLepTa gging _{min} > 0.9	94	91	31	10	2	2632	128	1	50	0
$\log_{10}(y_{12}) > -1$	94	90	31	9	2	2632	128	1	50	0
$P_{\text{miss}} < 100 \ \&\&$ ($P_{\text{miss}} < 40 \ \parallel$ $ \cos\theta_{P_{\text{miss}}} >$ 0.95)	84	84	28	9	1	79	30	0	9	0

Signal efficiency \sim **20%**
 Remaining backgrounds events \sim **150 (eLpR), 20 (eRpL)**

Reconstruction methods

After removing isolated electrons force into 4 jets (Durham)



Search for the correct combination of jj and jje

Jet pair 1 $\rightarrow M_{jj1}$, Jet pair 2 $\rightarrow M_{jj2}$

$$F_1 = (M_{jj1} - M_w)^2 + (M_{jj2} - M_w)^2$$

Best jet pair 1 + iso e $\rightarrow M_{jje1}$

Best jet pair 2 + iso e $\rightarrow M_{jje2}$

We expect for " $M_{jje1} = M_{jje2}$ "


$$F_2 = (M_{jje1} - M_{jje2})^2$$

Choose combination with minimum F_1, F_2

Signal mass cut

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
 $\mathcal{L} = 1600 [\text{fb}^{-1}]$

For each M_N , mass window M_N-10, M_N+15 [GeV]




Assume background distribution is flat

20 (eLpR) background events remain in mass window

Signal mass cut

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
 $\mathcal{L} = 1600 [\text{fb}^{-1}]$



Assume background distribution is flat

20 (eLpR) and 3 (eRpL) background events remain in mass window

Less backgrounds thanks to beam polarization

Reduce W contribution


Results

	M_N [GeV]	# of Signal After mass cut	# of BG After cut	σ_0 [fb] Initial benchmark ($ee \rightarrow NN$)	σ^{95} [fb] 95% exclusion limit ($ee \rightarrow NN$)	$\frac{\sigma^{95}}{\sigma_0}$
LR 80,30	100	53.64	20.12	0.35	0.073	0.21
	150	52.73		0.22	0.048	0.21
	200	18.30		0.088	0.054	0.61
	225	5.51		0.029	0.053	1.8
RL 80,30	100	66.75	3.24	0.43	0.040	0.092
	150	63.41		0.27	0.026	0.097
	200	21.23		0.10	0.029	0.29
	225	6.08		0.032	0.032	1

Exclusion plot on cross-section $\sigma(ee \rightarrow NN)$

$$\sigma = \sigma_0 \times \left\{ \frac{2}{N_S} \left(1 + \sqrt{1 + N_B} \right) \right\}$$


Calculate 95% UL on $\sigma(ee \rightarrow NN)$



Exclusion plot on σ/σ_0

Normalised to benchmark cross-section

Calculate 95% UL on σ/σ_0



Exclude benchmark points and cross-sections up to 10x smaller

Summary

Conclusion:

**Can use same sign lepton signature
to set powerful limits on RHN at ILC!**

Current activity & future plan:

- ILC250 case (on going)
 - Try to improve signal efficiency

- Same sign muons
 - Expect smaller backgrounds

Part1: RHN

Model : minimal $U(1)_{B-L}$

Gauged B-L extension of Standard Model (SM)

The unique anomaly free global symmetry in the SM

$$G_{B-L} \equiv SU(3)_C \times SU(2)_L \times U(1)_Y \times U(1)_{B-L}$$

► Anomaly free requirement → **RHNs**

► **Seesaw mechanism** ← automatically included

Gauge boson : Z'

If B-L symmetry breaks spontaneously → Z' becomes **massive**

minimal $U(1)_{B-L}$ model : charge



		$SU(3)_C$	$SU(2)_L$	$U(1)_Y$	$U(1)_{B-L}$
RHN	N_R^i	1	1	0	-1
New Higgs field	Φ	1	1	0	2

$i=1,2,3$


[arXiv\[1812.11931\]](https://arxiv.org/abs/1812.11931)

Arindam Das, Nobuchika Okada, Satomi Okada, Digesh Raut

Signal




Destructive interference




Isolated e, γ, μ

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$


Number of isolated e



Number of isolated γ



Number of isolated μ




- | | |
|-----------------|------------------|
| — $M_N=100$ GeV | — BG_6f_ttbar_2e |
| — $M_N=150$ GeV | — BG_6f_ttbar_1e |
| — $M_N=200$ GeV | — BG_6f_ttbar_0e |
| — $M_N=225$ GeV | — BG_4f_sw_sl |
| | — BG_4f_sze_sl |
| | — BG_eeqqqq |

- Isolated e # = 2 && Isolated $\gamma, \underline{\mu} = 0$

Distribution of Isolated electron energy


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



$E_{\text{iso}} < 200$ [GeV]

Distribution of $\cos\theta_{\text{isoe}}$

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)




$$-0.95 < \cos\theta_{\text{isoe}} < 0.95$$

4 fermions semi leptonic processes in t-channel \rightarrow distributed in $|\cos\theta_{\text{isoe}}| \sim 1$

Distribution of y_{12} (Durham)


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



$\log_{10}(y_{12}) > -1$

$\cos\theta_{P_{\text{miss}}}$ vs Magnitude of missing momentum P_{miss}

- ILC 500 with ISR / BS



$$P_{\text{miss}} < 100 \ \&\& \ (P_{\text{miss}} < 40 \ \&\& \ | \cos\theta_{P_{\text{miss}}} | > 0.95)$$

Cut flow (eRpL)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (+0.8, -0.3)$ $\mathcal{L} = 1600 [\text{fb}^{-1}]$

ILD work in progress


	Expected signal				Expected background					
	$M_N=100$	$M_N=150$	$M_N=200$	$M_N=225$	eeqqqq	4f_singleW semileptonic	4f_singleZee semileptonic	6f_ttbar 2electrons	6f_ttbar 1electron	6f_ttbar 0electron
No cut	558	394	143	45	3925	258648	612455	7100	56233	4894
$e_{\text{iso}} \# = 2 \ \&\&$ $\gamma_{\text{iso}} \# = 0 \ \&\&$	420	343	126	40	1935	9426	249000	6142	1295	127
Same sign ($e_{\text{iso}1} \times e_{\text{iso}2} = 1$)	346	115	81	12	1231	7210	140176	3911	870	94
$E_{\text{iso}} < 200$ [GeV]	171	114	41	12	14	3741	3294	2	177	19
$-0.95 <$ $\cos\theta_{\text{isoe}} < 0.95$	158	103	37	11	3	1324	475	1	113	12
IsolatedLepTa gging _{min} > 0.9	96	91	32	10	0	198	101	0	15	1
$\log_{10}(y_{12}) > -1$	88	90	30	9	0	199	86	0	6	0
$P_{\text{miss}} < 100 \ \&\&$ ($P_{\text{miss}} < 40 \ \parallel$ $ \cos\theta_{P_{\text{miss}}} >$ 0.95)	86	84	29	9	0	4	15	0	2	0

Results

	M_N [GeV]	# of Signal After cut	# of BG After cut	Signal Significance	σ_0 [fb] Initial benchmark	σ^{95} [fb] 95% exclusion limit	$\frac{\sigma^{95}}{\sigma_0}$	α^{95} [TeV ⁻⁴]
LR 80,30	100	53.64	20.12	6.25	0.55	0.12	0.21	3.83E-05
	150	52.73		6.18	0.36	0.076	0.21	2.96E-05
	200	18.30		2.95	0.14	0.086	0.61	7.57E-05
	225	5.51		1.18	0.046	0.085	1.8	2.21E-04
RL 80,30	100	66.75	3.24	7.98	0.71	0.065	0.092	1.69E-05
	150	63.41		7.77	0.45	0.043	0.097	1.35E-05
	200	21.23		4.29	0.16	0.047	0.29	3.57E-05
	225	6.077		1.99	0.052	0.052	1	1.21E-04

Exclusion plot on $U(1)_{B-L}$ parameters

Translate to the $U(1)_{B-L}$ model parameters



The benchmark points isn't excluded only at $M_N = 225$ GeV

eLpR case

Cut flow (eLpR)


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$ $\mathcal{L} = 1600 [\text{fb}^{-1}]$

ILD work in progress


	Signal Entries				Background Entries					
	$M_N=100$	$M_N=150$	$M_N=200$	$M_N=225$	eeqqqq	4f_singleW_semileptonic	4f_singleZee_semileptonic	6f_ttbar_2electrons	6f_ttbar_1electron	6f_ttbar_0electron
No cut	554	394	143	45	11898	2825010	699475	16425	129283	11028
$e_{\text{iso}} \# = 2$ && $\gamma_{\text{iso}} \# = 0$ &&	347	343	79	40	4721	90818	162774	9422	2271	201
Same sign ($e_{\text{iso}1} \times e_{\text{iso}2} = 1$)	176	115	39	12	39	46138	3800	8	439	25
$E_{\text{iso}} < 200$ [GeV]	175	114	39	12	39	41319	3557	8	439	25
$-0.95 <$ $\cos\theta_{\text{iso}e} < 0.95$	156	103	36	11	13	17506	623	4	266	15
IsolatedLepTagging _{min} > 0.9	94	91	31	10	2	2632	128	1	50	0
$\log_{10}(y_{12}) > -1$	94	90	31	9	2	2632	128	1	50	0
$P_{\text{miss}} < 100$ && ($P_{\text{miss}} < 40$ $ \cos\theta_{P_{\text{miss}}} >$ 0.95)	84	84	28	9	1	79	30	0	9	0

Electron Charge

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e^- \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$



Isolated electrons charge




We use only same sign samples $e_1 \times e_2 = 1$

Distribution of IsolatedLepTagging

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated e # = 2 && Isolated γ # = 0 && Isolated μ # = 0
- Isolated e is same sign ($e_1 \times e_2 = 1$)

Minimum of isolated lepton tagging



Isolated lepton tagging
... “output” parameter of MVA
to identify isolated lepton

→ Output for e is **near 1**


**Usually second background
electron is fake**

IsolatedLepTagging_{min} > 0.9


Isolated e, γ, μ

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$


Number of isolated e



Number of isolated γ



Number of isolated μ




- | | |
|-----------------|------------------|
| — $M_N=100$ GeV | — BG_6f_ttbar_2e |
| — $M_N=150$ GeV | — BG_6f_ttbar_1e |
| — $M_N=200$ GeV | — BG_6f_ttbar_0e |
| — $M_N=225$ GeV | — BG_4f_sw_sl |
| | — BG_4f_sze_sl |
| | — BG_eeqqqq |

- Isolated e # = 2 && Isolated $\gamma, \mu = 0$

Distribution of Isolated electron energy


- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



$E_{\text{iso}} < 200$ [GeV]

Distribution of $\cos\theta_{\text{isoe}}$

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)




$$-0.95 < \cos\theta_{\text{isoe}} < 0.95$$

4 fermions semi leptonic processes in t-channel \rightarrow distributed in $|\cos\theta_{\text{isoe}}| \sim 1$

Distribution of y_{12} (Durham)

- ILC 500 with ISR / BS
- $\text{Pol}(e^-, e^+) = (-0.8, +0.3)$
- Isolated $e \# = 2$ && Isolated $\gamma \# = 0$ && Isolated $\mu \# = 0$
- Isolated e is same sign ($e_1 \times e_2 = 1$)



$\log_{10}(y_{12}) > -1$

4f and 6f background information


Cross section — BG

- ILC 500 with ISR / BS

(100%,100%)	eeqqqq			4f_singleW _semileptonic	4f_singleZee _semileptonic
	eexyyx	xxxxee	yyyyee	4f_sw_sl	4f_sze_sl
eLpR	1.64E+01	8.71E-02	1.45E-01	7.81E+03	1.96E+03
eRpL	3.64	4.62E-02	5.31E-02	2.28E+01	1.73E+03
eLpL	6.63	3.38E-02	2.20E-02	7.53E+02	1.78E+03
eRpR	6.61	3.30E-02	1.97E-02	7.50E+02	1.78E+03

Cross section — BG


- ILC 500 with ISR / BS



Information associated
with $U(1)_{B-L}$ model


Current limits - Z' mass

SM like Z' coupling



ATLAS-TDR-LHCC2017-2018

HL-LHC prospects limit for $U(1)_{B-L}$ model




arXiv[1812.11931]

The heavier Z' mass less constrained by LHC

Current limits $|V_{eN}|^2$


$|V_{eN}|^2$: the “light-heavy” neutrino mixing matrix



<https://arxiv.org/pdf/1802.02965.pdf>

Current Limits and prospects - Z' mass, $g_{1'}$

$G_{1'}$: $U(1)_{B-L}$ gauge coupling constant



M_N [GeV]	$M_{Z'}$ [TeV]	$g_{1'}$
100	7	1
200	7	1